their first IPPV episode have subsequently died from their asthma, at median (range) 1.6 years (0.5–8.4) from their first IPPV.

Discussion

The treatment of asthma has changed considerably since 1964, and this was apparent among our ventilated patients. With more effective treatment available, one might expect the numbers of asthmatic children requiring ventilation to decline, but in our regional intensive care unit this is not occurring. This is probably because over the same period, admissions for acute asthma have increased considerably, so that in fact the proportion of admissions requiring ventilation has declined. An increase in tertiary referrals may also have contributed.

Asthmatic children who require IPPV tend to be either the very young or teenagers, to have developed asthma at an early age and required multiple admissions, have an equal sex distribution (in contrast to the overall male predominance in childhood asthma), and have been previously undertreated. In all these characteristics they resemble children who die from acute asthma,^{9 13 14} and indeed our follow up showed a high incidence of further IPPV episodes and of sudden death from asthma. There may be different mechanisms that predispose preschool and teenage asthmatics to respiratory failure. Young children have intrinsically narrower airways, may develop respiratory muscle fatigue more readily, and are less responsive to bronchodilators. Adolescents tend to comply poorly with treatment, may start to smoke, and may deny symptoms and therefore present later to hospital. The three children who presented to hospital after a respiratory arrest were aged 12, 14, and 15 years.

Asthmatics who survive IPPV tend to continue to have daily symptoms and considerable life disturbance, and although many experience further life threatening episodes, few have either a clear plan of action or extra treatment available at home to cope with a severe attack.

In the four large published series (comprising 19 to 30 episodes) of ventilated asthmatic children, all American, the mortality rate ranged from 0% to 5.3% of episodes.^{1–4} In our series mortality was considerably higher, eight of 48 episodes (17%) being fatal. No consistent differences were apparent between the American centres and Liverpool in management during IPPV, but in the American series the threshold for instituting IPPV appeared lower, and hence the ventilated children less severely ill. In Seattle for example, 5.3% of all acute asthma admissions received IPPV (0.6% in Liverpool), mean pH before intubation was 7.22 (7.09 in Liverpool), and 5% of IPPV episodes followed a respiratory arrest (10% in Liverpool).

We are still ventilating two to three asthmatic children per year, and it seems likely that IPPV could be avoided in some of these by improved medical management. Among the children we are ventilating, mortality appears unacceptably high, and is most commonly related to hypoxic-ischaemic encephalopathy before IPPV. Much of this was occurring in hospital, and again

should be preventable. The main inadequacies identified by our study were inconsistency in management (for example, failure to start steroids early), reluctance to use intravenous β agonists, reluctance to measure blood gases, and failure to diagnose respiratory failure early. We have attempted to correct these deficiencies by introducing a written protocol for inpatient management of acute asthma by junior staff. Early clinical diagnosis of hypoxia and respiratory failure is essential but difficult, and to measure arterial blood gases on all asthmatic children admitted would be neither practicable nor desirable. Semiobjective clinical scoring systems have been advocated,¹² but have not been proved to be of value. Pulse oximetry, now becoming more widely available, should allow earlier detection of hypoxia.

Ventilation is still required during some episodes of acute asthma despite improvements in the treatment available. Improved monitoring of children admitted with acute asthma and intensive treatment of early respiratory failure are necessary to reduce the need for IPPV and avoid hypoxic brain damage. Survivors continue to suffer from severe chronic asthma and are at risk of sudden death from their disease. They require close follow up into adult life and must have a clear action plan for acute attacks.

- 1 Woods DW, Downes JJ, Lecks HI. The management of respiratory failure in childhood status asthmaticus. Experience with 30 episodes and the evolution of a technique. *Journal of Allergy* 1968;42:261–8.
- 2 Parry WH, Martorano F, Cotton EK. Management of life-threatening asthma with intravenous isoproterenol infusions. *Am J Dis Child* 1976;130:39–42.
- 3 Simons FER, Pierson WE, Bierman CW. Respiratory failure in childhood status asthmaticus. *Am J Dis Child* 1977;131:1097–101.
- 4 Dworkin G, Kattan M. Mechanical ventilation for status asthmaticus in children. *J Pediatr* 1989;114:545–9.
- 5 Simpson H, Mitchell I, Inglis JM, Grubb DJ. Severe ventilatory failure in asthma in children. *Arch Dis Child* 1978;53:714–21.
- 6 Polgar G, Weng TR. The functional development of the respiratory system. *Am Rev Respir Dis* 1979;120:625–95.
- 7 Cockcroft DW, Killian DN, Mellon JJA, Hargreave FE. Bronchial reactivity to inhaled histamine: a method and clinical survey. *Clin Allergy* 1977;7:235–43.
- 8 Mitchell EA. International trends in hospital admission rates for asthma. *Arch Dis Child* 1985;60:376–8.
- 9 Office of Population Censuses and Surveys. *Mortality statistics series*. (DH2 Nos 1–12.) London: HMSO, 1977–1987.
- 10 Bohn D, Kalloghlian A, Jenkins J, Edmonds J, Barker G. Intravenous salbutamol in the treatment of status asthmaticus in children. *Crit Care Med* 1984;12:892–95.
- 11 Sury MR, Russell GN, Heaf DP. Hydrocortisone myopathy. *Lancet* 1988;ii:515.
- 12 Wood DW, Downes JJ, Lecks HI. A clinical scoring system for the diagnosis of respiratory failure. *Am J Dis Child* 1972;123:227.
- 13 Carswell F. Thirty deaths from asthma. *Arch Dis Child* 1985;60:25–8.
- 14 Speight ANP, Lee DA, Hey EN. Under diagnosis and under treatment of asthma in childhood. *Br Med J* 1983;286:1256–8.

Commentary

Mortality from asthma in the United Kingdom is unchanged or even rising,¹ despite the advent of new drugs, more publicity, greater critical appraisal of treatment policies, earlier hospital admission, and the wider availability of mechanical ventilation (IPPV). The asthmatic process is reversible in a high proportion of patients and mechanical ventilation should, in theory, buy time for other measures to take effect. Nevertheless, most series describing the outcome of IPPV for severe acute asthma in adults

record a significant mortality²⁻⁶ and identify avoidable factors.⁷⁻⁹ Against this background, it is pertinent to examine data for asthmatic children requiring IPPV and reported here from Alder Hey Children's Hospital.

Seddon and Heaf's results make depressing reading. Forty seven episodes of mechanical ventilation were required by 31 children, most around the age of 2 years but with a smaller peak in the teenage years. Approximately two thirds had been admitted more than once in the preceding year and only three children needed mechanical ventilation on their first admission. Despite this, more than half were receiving either no treatment or bronchodilators alone before their first episode of IPPV.

Eight children died during IPPV, five deaths occurring in patients who had required IPPV on an earlier occasion. Six suffered a cardiorespiratory arrest before intubation, three outside hospital but another three who were already in the intensive care unit. Five of these six children died.

Those intubated after an arrest had usually not had blood gas tensions measured. Fourteen children in whom there were no arterial blood gas measurements before intubation progressed to respiratory failure, in hospital, over a median time of six hours. By the time they were intubated, many had evidence of cerebral or cardiac hypoxia. Two suffered a respiratory arrest and one a cardiac arrest during intubation. Six were semiconscious, five unconscious, and two having fits before intubation.

Hypoxic-ischaemic encephalopathy developed during nine episodes of IPPV, three of which had been preceded by a cardiorespiratory arrest and one by hypoxic fits. Six of these nine instances proved fatal. There was evidence of cerebral hypoxia before intubation in five of the six.

Respiratory complications were seen too, lobar collapse in 29 episodes (60%), pneumomediastinum in 13 (27%), and pneumothorax in 10 (21%). Acute circulatory impairment or collapse after the institution of IPPV is the only complication recognised in adults but not included in this series.^{7 8 10-12}

A novel and welcome feature of this report is that the authors explored the subsequent progress of their subjects. Here too their findings are sombre. Four more children have died, all from a rapidly evolving attack of asthma. Sixteen of 17 survivors followed up for more than a

year continued to suffer asthmatic symptoms, which were present every day in more than half of them.

The conclusions are clear. IPPV is needed for a small number of asthmatic children each year. Avoidable deficiencies of management and an unacceptably high incidence of hypoxic complications have been identified. Survivors remain a high risk group in need of long term supervision and counselling, with a clear action plan agreed for handling an acute attack. As a result of these findings, a written protocol has been introduced at Alder Hey to improve the service and reduce mortality in future.

Seddon and Heaf are to be congratulated for their tenacity in gathering this data which covers a 25 year span, for their honest reporting of deficiencies in management, and for reminding us that there is a small, high risk group of asthmatic subjects for whom 'too little, too late' can easily prove fatal. Above all they are to be congratulated for their constructive and practical attempt to identify how services for these vulnerable young patients can be improved in the future.

MARGARET A BRANTHWAITE
Brompton Hospital,
Fulham Road,
London SW3 6HP

- 1 Burney PGJ. Asthma mortality in England and Wales: evidence for a further increase, 1974-84. *Lancet* 1986;ii:323-6.
- 2 Webb AK, Bilton AH, Hanson GC. Severe bronchial asthma requiring ventilation. A review of 20 cases and advice on management. *Postgrad Med J* 1979;55:464-7.
- 3 Westerman DE, Benatar SR, Potgieter PD, Ferguson AD. Identification of the high risk asthmatic patient. *Am J Med* 1979;66:565-72.
- 4 Petheram IS, Branthwaite MA. Mechanical ventilation for pulmonary disease: a six year survey. *Anaesthesia* 1980;35:467-73.
- 5 Picado C, Montserrat J, Roca R, et al. Mechanical ventilation in severe exacerbations of asthma: study by 26 cases with six deaths. *Eur J Respir Dis* 1983;64:102-7.
- 6 Luksza AR, Smith P, Coakley J, Gordan IJ, Atherton ST. Acute severe asthma treated by mechanical ventilation: 10 years' experience from a district general hospital. *Thorax* 1986;41:459-63.
- 7 Darioli R, Perret C. Mechanical controlled hypoventilation in status asthmaticus. *Am Rev Respir Dis* 1984;129:385-7.
- 8 Branthwaite MA. The intensive care of asthma. *Br J Hosp Med* 1985;34:331-8.
- 9 Brear SG, Edwards JD. Intermittent positive pressure hypoventilation and inotropic support in severe acute asthma. *Thorax* 1987;42:731-2.
- 10 Tuxen DV, Lane S. The effects of ventilatory pattern on hyperinflation, airway pressure, and circulation in mechanical ventilation of patients with severe airflow obstruction. *Am Rev Respir Dis* 1987;136:872-9.
- 11 Jardin F, Dubourg O, Margairaz A, Bourdarias J-P. Inspiratory impairment in right ventricular performance during acute asthma. *Chest* 1987;92:789-95.
- 12 Suter PM. Right ventricular pathophysiology in the critically ill. *Clinical Anaesthesiology* 1985;3:899-907.