comply with dietary regimes while actually taking a fraction of their documented intake. The boy was frequently reported in the nursing notes to be 'cheerful and eating well'. The girl underwent precise (and much resented) dietary assessment, showing an actual intake of only 1.25 MJ/day instead of the 5.02 MJ she was thought to be receiving. Attempted referral of both patients to appropriate psychiatric units was refused because of concern about the management of organic disease where there was no resident paediatric cover. The boy in particular was able to use his symptoms to forestall an arranged transfer and both would have been able to use their abdominal symptoms to ensure transfer back to the paediatric unit when they became resentful of the management of their eating disorder.

The features and management of each of these two conditions may clearly affect the other. Two cases of Crohn's disease and bulimia nervosa have been described in which corticosteroid treatment affected the role of the eating disorder.¹² Similarly, in both of our cases, the role of corticosteroid treatment was central. The boy stopped his treatment because of perceived changes in body shape and suffered life threatening consequences. Anorexia nervosa appears to have been precipitated in the girl by the rapid weight gain secondary to high dose steroid treatment. The use of corticosteroids is frequently unavoidable in inflammatory bowel disease, but the risks involved in such treatment at an age when a preoccupation with body shape is common should be clearly understood. The use of alternative management strategies such as elemental diet for small bowel disease should be considered, and patients should be fully counselled about the effects of steroids before they are started. Patients should be followed up regularly and specifically questioned about eating problems. Suspicion of abnormal eating behaviour or unexplained weight loss warrants early formal psychiatric assessment and admission for observation.

We would like to thank Professor JA Walker-Smith for permission to report on his patients and for his kind advice in the preparation of this manuscript.

- 1 Kelts DG, Grand RJ. Inflammatory bowel disease in children and adolescents. *Curr Probl Pediatr* 1980;10:1-40.
- 2 Walker-Smith JA. Crohn's disease and abdominal tuberculosis. *Diseases of the small intestine in childhood*. 3rd ed. London: Butterworths, 1988:328-60.
- 3 Bryant-Waugh R, Knibbs J, Fosson A, Kaminski Z, Lask B. Long term follow up of patients with early onset anorexia nervosa. *Arch Dis Child* 1988;63:5-9.
- 4 Anonymous. Case records of the Massachusetts General Hospital. *N Engl J Med* 1985;312:1175-83.
- 5 Jenkins AP, Treasure J, Thompson RP. Crohn's disease presenting as anorexia nervosa. *Br Med J* 1988;296:699-700.
- 6 Gryboski JD, Katz J, Sangree MH, Herskovic T. Eleven adolescent girls with severe anorexia. Intestinal disease or anorexia nervosa? *Clin Pediatr (Phila)* 1968;7:684-90.
- 7 Hershman MJ, Hershman M. Anorexia nervosa and Crohn's disease. *Br J Clin Pract* 1985;39:157, 159.
- 8 Metcalfe-Gibson C. Anorexia nervosa and Crohn's disease. *Br J Surg* 1978;65:231-3.
- 9 Sreenivasan U. Anorexia nervosa associated with energy-wasting disorders. *Can Med Assoc J* 1984;130:45-6.
- 10 Dally P, Sargent W. Treatment and outcome of anorexia nervosa. *Br Med J* 1966;ii:793-5.
- 11 American Psychiatric Association. *Diagnostic and statistical manual of mental disorders*. 3rd ed, revised. Washington, DC: American Psychiatric Association, 1987.
- 12 Meadows G, Treasure J. Bulimia nervosa and Crohn's disease: two case reports. *Acta Psychiatr Scand* 1989;79:413-4.

Commentary

Childhood onset anorexia nervosa is often a difficult condition to diagnose and it is frequently overlooked. In one study, the average time from onset to diagnosis was nine months.¹ The main reasons for the frequent delay in diagnosis are the child's denial and secretiveness. Thus the pathognomonic symptoms such as fear of fat and distorted body image are commonly denied, and other common features such as *determined* food avoidance, preoccupation with weight and calories, self induced vomiting, exercise and laxative abuse are usually disguised. Anorexia nervosa may all the more easily be overlooked when it complicates inflammatory bowel disease.

In the two patients described by Mallett and Murch, the diagnosis was made on the basis of overt symptoms. In the first case, the boy had a two year history of binge eating, excessive exercising, preoccupation with food intake, weight and body shape, all superimposed upon weight loss. In the second case, the girl had a far shorter history of weight loss, abnormal eating behaviour, and of distorted body image.

How is a diagnosis of anorexia nervosa made in a child with inflammatory bowel disease whose symptoms are far less overt? Firstly, the paediatrician must always be aware of the possibility of a concomitant psychiatric disorder in any child with serious illness. The incidence in such circumstances may be as high as 25%. The regular use of a multiaxial diagnostic scheme such as the International Classification of Diseases² avoids the either/or dichotomy and instead encourages due consideration being given to medical, psychiatric, developmental, intellectual, and social status. Secondly, the paediatrician should have a high index of suspicion when assessing a child with poorly explained weight loss. Certainly the diagnosis should never be excluded on the basis of negative answers to questions about concerns regarding food, weight, and appearance. Thirdly, the diagnosis can almost always be made while the discussion focuses on what would constitute a satisfactory weight, and how much energy should be consumed each day. At this point, children with anorexia nervosa almost always become distressed, argumentative, and negativistic in just the same way as do children whose school phobia has presented with recurrent abdominal pain when told they should now return to school. Finally, if doubt remains, careful observation in hospital will reveal characteristic abnormal eating behaviour and a strong element of secretiveness.

Early diagnosis of anorexia nervosa complicating inflammatory bowel disease is imperative and not only is the condition much harder to treat when treatment is delayed, but as Mallett and Murch have demonstrated it can cause major and even life threatening complications.

- 1 Fosson A, Knibbs J, Bryant-Waugh R, Lask B. Early onset anorexia nervosa. *Arch Dis Child* 1987;62:114-8.
- 2 Rutter M, Shafer S, Sturge C. *A guide to a multi-axial classification scheme for psychiatric disorders in childhood and adolescence*. London: Institute of Psychiatry, 1975.

B LASK
Department of Psychological Medicine,
Hospital for Sick Children,
London