

All medical records and imaging studies were reviewed.

Results Number of Deliveries over the 5-year study period was 25443 of which 4100 (16%) were un-booked. Dilated cistern magna was diagnosed in 26 (0.12%) fetal scans (15 male and 11 females). Post natal scans were done in 20/26 (77%) and not done in 6/26 (23%) of cases.

Of the 20 post natal scans done 12 (60%) were normal and 8 (40%) were abnormal. The following abnormalities were detected: 4 (50%) dandy walker complex, 2 (25%) corpus collasum dysgenesis and 2 (25%) cerebellar hypoplasia. Neuro-developmental assessment was reported as normal in 60% of the isolated mega cistern magna patients.

Conclusion The association of mega cisterna magna with major CNS anomalies is high in our population. Post natal neuroimaging confirmation of all abnormal fetal sonography is required. Long term neurocognitive assessment and follow up is essential for this population.

1057 SPECTRUM OF CRANIAL ULTRASOUND FINDINGS IN NEWBORNS UNDER 26 WEEKS GESTATION OVER 10 YEAR PERIOD IN A TERTIARY NEONATAL UNIT

doi:10.1136/archdischild-2012-302724.1057

¹J Kapur, ²G Cattell, ³SV Rasiah. ¹Department of Neonatology; ²Department of Radiology, Birmingham Women's NHS Foundation Trust, Birmingham, UK

Background and Aims Cranial ultrasonography is an integral part of routine neonatal screening in extreme preterm neonates. To audit the spectrum of cranial ultrasound scan findings in preterms less than 26+0 weeks gestation.

Methods We reviewed all cranial ultrasound findings performed in babies less than 26+0 weeks gestation between 1999–2008. Demographic data was collected using a proforma. Information regarding cranial ultrasound scans on day 1–4, day 7 and day 28 of life for live born babies admitted to the neonatal unit was collected.

Results The results are tabulated in the tables attached. Table 1 shows the demographic details and table 2 shows the spectrum of cranial ultrasound findings.

Abstract 1057 Table 1 Demographic details

Gestation (weeks)	Number of babies	Median birth weight(grams)	Appgar at 1 minute(median)	Appgar at 5 minutes(median)
23 – 23+6	11	630	3	7
24 – 24+6	50	660	4	8
25 – 25+6	69	734	5	8

Conclusion Extremely preterm babies (23 and 24 weeks gestation) had a higher incidence of abnormal cranial ultrasound scans compared to those over 25+0 weeks gestation. This is associated with a high risk of morbidity and mortality. This information is important when counselling parents and for prognosticating outcomes.

1058 NEW SEGMENTATION METHOD SHOWS EFFECTS OF PREMATURITY ON CEREBRAL TISSUE VOLUMES AT TERM

doi:10.1136/archdischild-2012-302724.1058

Abstract 1057 Table 2 Spectrum of cranial ultrasound findings

Gestation	Day 1–4		No results available/died	Day 7		No results available/died	Day 28		No results available/lost to follow up/died
	Normal	Abnormal		Normal	Abnormal		Normal	Abnormal	
23 – 23+6	6	5(45%)	0	4	7(63%)	0	3	7(63%)	1
24 – 24+6	24	19(38%)	7	13	23(46%)	14	15	14(28%)	23
25 – 25+6	44	17(24.6%)	8	39	18(26%)	12	28	12(17.3%)	29

¹L Gui, ²F Lazeyras, ³A Truttman, ¹C Borradori Tolsa, ¹PS Hüppi. ¹Pediatrics, Children's Hospital Geneva; ²Radiology, Geneva University Hospital, Geneva; ³Pediatrics, Children's Hospital Lausanne, Lausanne, Switzerland

Background and Aim Longitudinal studies show that premature birth increases infants' risk for mental and motor development deficits. Our aim was to investigate the influence of prematurity on cerebral tissue volumes at term obtained with a novel fully automatic segmentation method.

Methods 62 preterm infants (GA 27.7±1.3wks) and 15 term-born infants (GA 40±1.1wks) were scanned at term-equivalent age (GA 40.5±1.5wks). T1 and T2 MR images were segmented with a novel atlas-free automatic method based on morphological constraints. Each brain was separated into the two hemispheres, cortical and subcortical gray matter, myelinated and unmyelinated white matter, brainstem, cerebellum and CSF.

Results Linear regression models were fitted to study the dependency of tissue volumes on GA at birth, GA at scan and intracranial volume. Models show significant dependence on GA at birth for cortical gray matter (Beta=0.270, P=0.000, R²=0.818), unmyelinated white matter (Beta=0.196, P=0.03, R²=0.575), cerebellum (Beta=0.348, P=0.000, R²=0.648) and CSF (Beta= -0.329, P=0.000, R²=0.708).

Wilcoxon Signed Ranks tests showed significantly larger unmyelinated white matter volumes in the right hemisphere compared to the left hemisphere (Z= -4.826, P=0.000), and significantly larger total volumes of the right hemisphere compared to the left hemisphere (Z= -3.486, P=0.000).

Conclusions Reliable volume assessments were derived from the new automatic segmentation. CSF volumes at term increased with lower GA at birth, while cortical gray matter, unmyelinated white matter and cerebellum volumes at term increased with GA at birth, suggesting impaired growth of these tissues associated with prematurity. Cerebral asymmetry was present at term for both preterm and term infants.

1059 MYELIN IS DIFFERENTIALLY ASSOCIATED WITH RESTING STATE FUNCTIONAL CONNECTIVITY IN ADULTS WHO WERE BORN VERY PRETERM AND CONTROLS

doi:10.1136/archdischild-2012-302724.1059

¹S Froudish Walsh, ²SH Kolind, ¹KW Nam, ¹M Walshe, ²J O'Muircheartaigh, ²S Deoni, ²SC Williams, ¹RM Murray, ¹MP Allin, ¹C Nosarti. ¹Department of Psychosis Studies; ²Centre for Neuroimaging Sciences, Institute of Psychiatry, King's College London, London, UK

Background and Aims Diffuse white matter injury is the most common form of brain injury following very preterm (VPT) birth. This may reflect altered myelination, which could affect both neurodevelopment and neuronal communication. We investigated whether myelin in the corpus callosum (CC) was associated with functional connectivity; and if these associations differed between young adults born VPT and controls.

Methods 9 VPT-born adults and 13 controls (age 26–28 years) underwent resting state functional MRI (rs-fMRI), diffusion MRI and mcDESPOt, a novel neuroimaging method which provides an *in vivo* estimate of myelin water fraction (MWF). MWF was calculated along the CC. The default mode network (DMN), which