

Insofar, blood pressure and sleep-related breathing should be monitored beginning with an obstructive AHI of 3/hour sleep. Children with OSA should be treated early enough. Prevention and early treatment of obesity as a risk factor for OSA as well as for hypertension is becoming an important social challenge.

135 VENTILATORY SUPPORT IN PATIENTS WITH CENTRAL HYPOVENTILATION SYNDROME (CHS) - RESULTS OF THE EU-CHS NETWORK SURVEY

doi:10.1136/archdischild-2012-302724.0135

¹M Migdal, ²H Trang, ³J Peters, ⁴M Katz-Salamon, ⁵G Ottonello, ⁶M Samuels. ¹Children's Memorial Health Institute, Warsaw, Poland; ²Hopital Robert Debre, Paris, France; ³Klinikum Dritter Orden, Munich, Germany; ⁴Karolinska Institutet, Stockholm, Sweden; ⁵G Gaslini Institute, Genoa, Italy; ⁶University Hospital of North Staffordshire, Stoke on Trent, UK

Background and aim of the study Aim of the study was to identify the European centers for diagnosis and treatment of patients with CHS and to collect information on type of ventilatory support currently used for this group of patients.

Method The survey was performed in countries participating in the EU-CHS network project. Using a special template following information were collected: total number of already diagnosed patients, number of currently treated patients, type of ventilatory support (invasive ventilation via tracheostomy, mask ventilation, diaphragm pacing, negative ventilation).

Results The survey was performed during 2011. Replies were received from 15 centers located in 10 European countries. Number of CHS patients already diagnosed and currently treated varies significantly between centers (from 2 to 112 and from 1 to 70 respectively). Total number of already diagnosed CHS patients were 305 and 199 of them are currently treated. Most frequently used type of ventilator support is ventilation via tracheostomy (52% of patients). Diaphragm pacing alone or diaphragm pacing and ventilation is currently used in 42 patients. Only one CHS patient is currently treated with negative pressure ventilation.

Conclusions Results of this study have shown existing significant differences (mainly number of patients) but also some similarities (type of ventilatory support) between centers for CHS diagnosis and treatment located in 10 different European countries. All of the collected data should be used for further work on the European CHS network of reference centers.

The study was supported by the EU-CHS Network (grant number 2008 12 06).

136 SPONTANEOUS BREATHING PATTERNS OF TRANSITIONING PRETERM INFANTS IN THE DELIVERY ROOM (DR) AND INTERACTIONS WITH MANUAL POSITIVE PRESSURE VENTILATION

doi:10.1136/archdischild-2012-302724.0136

^{1,2,3}G Schmölder, ²J Kaufman, ^{2,4}O Kamlin, ^{2,4}P Davis. ¹Department of Pediatrics, University of Alberta, Edmonton, AB, Canada; ²Neonatal Services, The Royal Women's Hospital, Melbourne, VIC, Australia; ³Department of Paediatrics, Medical University of Graz, Graz, Austria; ⁴Dept. of Obstetrics & Gynaecology, The University of Melbourne, Melbourne, VIC, Australia

Background Positive pressure ventilation (PPV) remains the cornerstone of respiratory support after birth. Effectiveness of PPV in the DR may be compromised by leak, obstruction and inappropriate tidal volume (V_T) delivery.

Aim Describe leak, obstruction and V_T during spontaneous breathing amongst preterm infants transitioning in the DR and interaction of these breaths with mask ventilation.

Methods Patients were enrolled from the control arm of a randomised trial investigating the use of a respiratory function monitor

(RFM) in the DR; infants < 32 weeks who were stabilised by paediatric trainees unaware of RFM data were included. During spontaneous breathing on mask continuous positive airway pressure (CPAP) and the delivery of PPV, airway pressures, gas flow and V_T were recorded. Data were analysed by breath type.

Results In 29 infants, a total of 3864 inflations (mechanical) and inspirations (infant breaths) were analysed (Table 1). Overall mask leak was highest during spontaneous inspirations occurring between manual inflations and those during CPAP. The average V_T was highest during combined inspiration and inflation.

Abstract 136 Table 1 Type of Inflation/Breath

	Number of Inflations	Mask Leak (%)	Tidal volume (mL/kg)
Manual Inflation	2055 (53%)	41 (32–53)	8.2 (6.1–10.7)
Assisted Inflation	335 (9%)	61 (46–81)	9.3 (4.5–12.6)
Spontaneous breath during PPV	181 (5%)	92 (65–100)	2.9 (2.1–4.3)
Spontaneous breath while on CPAP	1285 (33%)	66 (41–100)	4.4 (2.5–6.2)

Conclusion Preterm infants are active participants in stabilisation in the DR. V_T and mask leak vary depending on the interaction between the infant's breathing efforts and the inflations provided by the resuscitator.

137 HIGH FLOW NASAL CANNULAE (HFNC) OR NASAL CONTINUOUS POSITIVE AIRWAY PRESSURE (NCPAP) POST-EXTUBATION IN PREMATURE INFANTS? A RANDOMISED CONTROLLED TRIAL

doi:10.1136/archdischild-2012-302724.0137

¹C Collins, ¹JR Holberton, ¹C Barfield, ²PG Davis. ¹Neonatal Medicine, Mercy Hospital for Women; ²Newborn Research, Royal Women's Hospital, Melbourne, VIC, Australia

Background NCPAP facilitates successful extubation. It is unclear whether HFNC are as effective as NCPAP in preventing extubation failure. In addition to an alternative modality of respiratory support HFNC may result in less nasal trauma than NCPAP.

Methods 132 preterm ventilated infants were randomised and stratified by gestation (< 28 vs 28–32 weeks). Primary outcome was extubation failure defined by a composite of 3 pre-specified failure criteria in the 7 days post-extubation. Individual failure criteria were not mutually exclusive and are defined; Apnoea, > 6 episodes in 6 hours or 1 requiring IPPV, Acidosis, pH < 7.25 & $pCO_2 > 66$ mmHg, and >15% increase in FiO_2 from extubation. A nasal trauma score was adapted from Kaufman [E-PAS 2007:61390].

Results

Abstract 137 Table 1

	HFNC N=67	NCPAP N=65	
Male n (%)	33 (49)	41 (63)	
Birthweight g mean (SD)	1123 (317)	1105 (374)	
Mean Completed Weeks gestation (SD)	27.9 (1.95)	27.6 (1.97)	
Failed Extubation in 1st week by Composite Criteria n (%)	15 (22)	22 (34)	
Apnoea: >6 in 6hrs or 1 needing IPPV n (%)	14 (21)	17(26)	
Acidosis: pH < 7.25 & $pCO_2 > 66$ mmHg n (%)	0	3 (5)	
>15% increase in FiO_2 post-extubation n (%)	7 (10)	12 (18)	
Reintubated in 1st week n (%)	7 (10)	8 (12)	
Nasal Trauma Score 1st week mean (SD)	3.1 (7.2)	11.8 (10.7)	p < 0.001*