

Conclusion In the group of massaged infant seems to obtain better results in motor milestones and in cognitive development, faster maturation of stereopsis, a increase of visual acuity have been also reached.

1737 EPIDEMIOLOGICAL ASPECTS OF STREPTOCOCCAL PHARYNGEAL INFECTIONS IN PEDIATRIC POPULATION

doi:10.1136/archdischild-2012-302724.1737

¹S Iurian, ²L Bera, ³SI Iurian, ¹M Mihut, ³ML Neamtu, ⁴A Muntean. ¹Clinical Laboratory, Pediatric Hospital; ²Statistics Department, Faculty of Medicine, 'Lucian Blaga' University; ³Research Department, Pediatric Clinic, 'Lucian Blaga' University; ⁴Pediatric Clinic, Sibiu, Romania

Background During last 5 years, we noticed an increasing incidence of scarlet fever and streptococcal pharyngitis in our county.

Aims

1. To appreciate the positive results rate for beta-hemolytic pyogenic streptococci in throat specimens (group A streptococci- gAs, group C- gCs, group G- gGs);
2. To evaluate ratio of each streptococcal group pharyngeal infection;
3. To establish correlation between streptococcal infections and diseases that justified throat cultures.

Methods Authors designed a retrospective epidemiological study, analyzing microbiology department data during 14 months period. Inclusion criteria: hospitalized and ambulatory care children aged between 2–18 years (scarlet fever diagnosis, pharyngitis diagnosis, healthy children requesting throat exam). Exclusion criteria: children up to 2 years of age. In order to identify streptococci, authors used Columbia agar with 5% sheep blood, Bacitracin inhibition tests, latex agglutination. Data was statistically analyzed using likelihood ratio.

Results Among 6653 throat cultures, 497 isolates (7.47%) were positive. Ratio for each streptococci group was: group A –88.0%, group C –6.2%, group G –5.8%. Seasonal incidence: higher incidence was reported in February and lower incidence in August. Regarding correlation between patient diagnosis and identified streptococci group in throat specimens (p value =0.000): 112 scarlet fever patients (111 gAs, 1 gGs), 264 pharyngitis patients (234 gAs, 13 gCs, 17 gGs), 121 healthy children (93 gAs, 18 gCs, 10 gGs). Last mentioned patients mean pyogenic streptococci carriers (24.34%).

Conclusions Carriers represents the “infection pool” for community children, maintaining persistence of source infection and explaining diminished efficacy of epidemiological measures and infection outbreaks in pediatric population.

1738 EFFECTS OF FORSYTHIA KOREANA-INCLUDED HERBAL MEDICINE TREATMENT ON UPPER RESPIRATORY TRACT INFECTION IN KOREAN CHILDREN

doi:10.1136/archdischild-2012-302724.1738

¹DY Yoo, ¹SB Yang, ¹JH Park, ¹NG Kim, ¹MA Kim, ²SY Lee, ²HY Choi, ¹H Choi. ¹Department of Research and Development, Hamsoa Oriental Clinic Network; ²Department of Pharmacology and Development, Hamsoa Pharmaceutical, Seoul, Republic of Korea


Background and Aims Korea has known high antibiotics prescription in children upper respiratory tract infections (URIs). Recently, herbal therapy is be magnified instead of unnecessary antibiotics prescription. *Forsythia Koreana* has been extensively used for the treatment of viral and bacterial respiratory tract infections. However, it remains unclear whether the herbal medicine has beneficial effects through clinical control study in URIs. Therefore, to investigate the anti-viral effects of *Forsythia Koreana* included herbal medicine (FKGM) on URIs in children, we conducted a randomized, double-blind, placebo-controlled study.

Methods Participants included 29 children who received either placebo (n=15) or FKGM (n=14) at the onset of URIs symptoms. The therapeutic effects assessed URI symptoms scoring system given by the James A Taylor for clinical studies to identify children with a documented viral URIs. Results were evaluated using student's t-test; the Kaplan-Meier method was used to analyse data regarding symptom duration.

Results

Abstract 1738 Table 1 URI symptom score

Symptom	Placebo (n=15)	FKGM (n=14)	p-value
throat pain	3.27±4.17	1.21±2.42	0.120
expectoration of sputum	4.93±4.04	3.00±2.75	0.146
sneezing	4.13± 4.21	2.21±2.52	0.147
runny nose	9.60±6.67	6.36±4.53	0.140
nasal congestion	8.40±5.10	3.79±2.72	0.006*



Abstract 1738 Figure 1 URI symptom duration between placebo and FKGM

Conclusions FKGM was more effective than the placebo in terms of reducing the duration of URI symptoms and reducing nasal discharge. These findings suggest that FKGM can be used for the replacement of antibiotics.

1739 SOCIOECONOMIC CHARACTERISTICS OF THE CHILDREN WHO NEEDED HOSPITALIZATION IN A PEDIATRIC INTENSIVE CARE UNIT (PICU)-RETROSPECTIVE ANALYSIS

doi:10.1136/archdischild-2012-302724.1739

¹AM Spanaki, ²M Linardakis, ¹D Fitrolaki, ¹T Tavladaki, ¹E Blevrakis, ¹S Ilia, ¹E Vasilaki, ¹E Geromarkaki, ²C Lionis, ²T Philalithis, ¹G Briassoulis. ¹Pediatric Intensive Care Unit, University Hospital of Heraklion; ²Public Health and Health Care Management, Faculty of Medicine, University of Crete, Heraklion, Greece