

10–14% and 10 had $\geq 15\%$. The median age at presentation was 5 days (2–12 days), median sodium level with $>15\%$ weight loss babies was 152 (143–162).

Conclusions A significant number of neonatal re-admissions were due to abnormal weight loss of $\geq 10\%$ and majority were due to failure to establish breastfeeding. An earlier audit in 2009 had identified need for more breast feeding support. Although systems to support breast feeding mothers in community are in place, more support needs to be established including regular assessment of weight to avoid hospital re-admissions.

1761 PITFALLS OF THE NEONATAL SCREENING PROGRAM FOR CONGENITAL ADRENAL HYPERPLASIA (CAH)

doi:10.1136/archdischild-2012-302724.1761

HA Abu Hamdan, F Sabbagh, MM Abou Al-Seoud. *Neonatal Division - Paediatric Department, King Fahad Armed Forces Hospital, Jeddah, Saudi Arabia*

Background Prompt diagnosis and treatment of CAH is essential to prevent mortality and morbidity. The incidence of CAH ranges from 1 in 21270 (New Zealand) to 1 in 5000 (Saudi Arabia).

Our Aim:

Identify the incidence of CAH in our patient population.
Appraise the CAH screening program.

Method Retrospective analysis of filter paper blood samples from infants aged 36 hours or more collected from November 2007 - Sept 2011. The 17-hydroxyprogesterone (17-OHP) tests were part of metabolic screening program at KFAFH. Abnormal values were re-called.

Results 22381 of 22428 (99.7%) births were screened for CAH. The turnaround time was 10 days. 124 infants [males 89 (72%) females 31 (28%)] had abnormal levels and were re-called (recall rate 0.55%). 76 infants had repeated serum 17-OHP concentration (response rate 61%). 7 infants had abnormal elevated levels, 4 had ambiguous genitalia (karyotype female), 3 male infants had symptoms of salt wasting at age of 9.10 and 37 days, turnaround time for the screening results was 11.9 and 16 days respectively. Delayed identification was due to failed contact and response in case 2 and 3 respectively. Recall failure was in 48/124 (39%) reasons were wrong contact numbers in 36 cases (75%) and no show in 10 cases (21%). Failure to repeat in 2 preterm infants died secondary to prematurity.

Conclusion The incidence of CAH is 1 in 3333 in our patient population. Barriers for timely intervention were due to prolonged turnaround time, lack of family education. Strategies to improve the process should be implemented.

1762 CURRENT RESUSCITATION PRACTICES AMONG PEDIATRICIANS IN GUJARAT, INDIA - EFFECT OF 2010 RESUSCITATION GUIDELINES

doi:10.1136/archdischild-2012-302724.1762

¹AS Nimbalkar, ¹SC Bansal, ¹DV Patel, ¹AR Sethi, ^{1,2}SM Nimbalkar. *¹Department of Pediatrics, Pramukhswami Medical College; ²Central Research Services, Charutar Arogya Mandal, Anand, India*

Background and Aims Neonatal Resuscitation Guidelines are updated by international societies every five years. India does not have mandatory certification and hence many pediatricians may not change their practices which may be in variance with current guidelines. We aimed to see the acceptance and awareness of several recently incorporated practices in resuscitation.

Methods Questionnaire based cross-sectional study administered by a web based form the link of which was sent by email and/or social networking sites to 600 pediatricians in state of Gujarat. A total of 25 questions were asked requiring about 10 minutes of the respondents time.

Results 75 pediatricians responded with 62 usable questionnaires. About 83% participants worked in a Level 2/3 NICU. 64% had resuscitated more than 20 neonates and 78% had attended more than 100 deliveries. Hence respondents were attending deliveries and required skills in resuscitation. 81.3% used bag and mask with only 33.3% using room air for resuscitation. 58.5% were using correct rate of ventilation. 76% had saturation monitors but only 17.2% had oxygen blenders. 31.7% approved of self inflating bag for free flow oxygen. 50% judged adequacy of ventilation by chest rise against 41.9% who used rising heart rate. Only about 50% pediatricians had undergone a training course in past 3 years. 41.7% believed in immediate cord cutting for normal deliveries. Only 32.3% responded correctly on the duration of resuscitation for asystole.

Conclusions The knowledge of pediatricians regarding resuscitation of neonates is poor. Only few pediatricians update themselves by attending courses.

1763 NEWBORN HEARING SCREENING: THE EXPERIENCE OF THE UNIVERSITY HOSPITAL OF MODENA

doi:10.1136/archdischild-2012-302724.1763

¹S Palma, ²MF Roversi, ³FM Artioli, ³G Denti, ²F Ferrari, ³E Genovese. *¹Primary Care Department, Azienda ASL; ²Department of Neonatology, Neonatal Intensive Care Unit, Modena University; ³Ent Department, Audiology, Modena University, Modena, Italy*

Background and Aims Newborn hearing screening programs allow to identify infants with bilateral permanent congenital hearing impairment and to facilitate early intervention to minimize the consequences on language development. In our country the concept of implementing universal newborn hearing screening protocols is still a topic of debate. Our aim was to analyze the first data collected in one year of experience since the adoption of the universal newborn hearing screening in Modena University Hospital.

Methods Data were collected during the period from 8th April 2011 to 31st March 2012. The screening was carried out by means of Transient Stimulus Evoked Otoacoustic Emissions, using, for well babies, a two-stage protocol: first screening stage on the second day after birth, followed by re-screening before discharge if a pass response was not obtained from both ears and a second stage follow up screening within 3 weeks later in case of failure. In case of a persistent failure response an audiological evaluation was performed.

Results During the period of the study 3512 babies underwent newborn hearing screening, of these 203 presented increased risk factors. In well babies group, one child will undergo surgery for a cochlear implant, 3 children are receiving a prosthetic-rehabilitative treatment. The prevalence of permanent bilateral hearing loss resulted 1.2:1000.

Conclusion It is important to create an active collaboration between audiologists/ENT specialists and paediatricians to reach the objective of identifying infants with hearing loss as early as possible in order to implement early interventions.

1764 INVESTIGATION OF SOME ASPECTS OF PERCEIVED SOCIAL SUPPORT IN MOTHERS OF HOSPITALIZED CHILDREN IN NEONATAL INTENSIVE CARE UNIT

doi:10.1136/archdischild-2012-302724.1764

¹S Kara, ²S Tan, ²S Aldemir, ³AE Yilmaz, ³MM Tatli, ⁴U Dilmen. *¹Department of Neonatology, Ankara Training and Research Hospital; ²Department of Psychiatry; ³Department of Neonatology, Fatih University Medical Faculty; ⁴Department of Neonatology, Zekai Tahir Burak Maternity and Teaching Hospital, Ankara, Turkey*

Purpose To identify the degree of perceived social support in mothers of hospitalized children in neonatal intensive care unit, and to assess in psychosocial aspects the relationship of the perceived social support with such variables as depression and anxiety level.