



Abstract G199(P) Figure 1

complete the full 2-week course of Omegaven®. All had total bilirubin levels above 80 µmol/l at commencement of Omegaven®. During their episodes of sepsis, bilirubin and CRP rose in all patients. Transaminases were deranged in all. All 7 patients showed improvement in septic markers during Omegaven® treatment. 3 patients showed improvement in bilirubin during treatment, which was maintained in the long term in 2. 1 patient was transferred to another centre for further medical treatment early in her Omegaven® course: her bilirubin was static.

**Conclusion** Use of Omegaven® as a short term rescue ILE in infants with IFALD and sepsis appears safe. The expected deterioration in liver function associated with sepsis was not seen in this series.

#### G200(P) EVALUATION OF SHARED CARE FOR IBD WITHIN A REGIONAL CLINICAL NETWORK

doi:10.1136/archdischild-2013-304107.212

<sup>1</sup>SP Paul, <sup>2</sup>J King, <sup>3</sup>BK Sandhu. <sup>1</sup>Paediatrics, Yeovil District Hospital, Yeovil, UK; <sup>2</sup>Paediatrics, Great Western Hospital, Swindon, UK; <sup>3</sup>Paediatric Gastroenterologist, Bristol Royal Hospital for Children, Bristol, UK

**Introduction** In 2008 BSPGHAN guidelines for Inflammatory Bowel Disease (IBD) were published. The South-West of England paediatricians have developed a regional clinical network whereby children, suspected of having IBD travel to Bristol for full diagnostic work-up. Post diagnosis care is provided by paediatricians at the local hospital with advice from Bristol and a joint three monthly out-patient clinic in Swindon. Children with very severe disease or needing surgery are dealt with by Bristol team. Effectiveness of this model was audited and found to be improved after the introduction of BSPGHAN guidelines

**Aim** This pilot study aimed to assess parent and patient views regarding quality of service provided within Swindon/Bristol regional network.

**Subjects and methods** Thirteen children aged <16 years diagnosed with IBD between 2010–2011, managed by shared care services. A telephone questionnaire survey designed with 12 questions and a free comments section. Questions included length of time to diagnosis, information sharing, satisfaction with services and preferences re further follow-up care.

**Results** 9/13(69%) responded to telephone survey, 3/13(23%) had moved out of area and 1/13(8%) could not be contacted by telephone. Of 9 patients, 4 had Crohn's disease, 3 Ulcerative colitis and 2 Indeterminate colitis. 4/9(45%) parents felt their concerns were adequately addressed initially at Swindon whereas all parents were satisfied with services provided at Bristol for diagnostic work up. 6/9(67%) parents felt they were satisfied with the expertise available locally for post-diagnosis management. However 8/9(89%) parents were happy with follow-up care by the joint care services at Swindon.

**Conclusion** This single centre pilot demonstrated that joint care provided by this model not only leads to care more concordant with BSPGHAN guidelines but is appreciated and valued by parents. There is scope for further improvements. This pilot study provides a template for ensuring and improving parent/patient involvement and satisfaction; there are plans to modify the questionnaire taking into account any suggestions for improvements and roll it out over the whole SW shared care IBD network soon.

#### G201(P) SURVEY OF MANAGEMENT OF IRON DEFICIENCY ANAEMIA IN CHILDREN WITH INFLAMMATORY BOWEL DISEASE IN THE UK

doi:10.1136/archdischild-2013-304107.213

S Ramprakash, A Modi. Department of Paediatrics, Luton and Dunstable Hospital, Luton, UK

**Aim** To study the current practise of management of iron deficiency anaemia in children with inflammatory bowel disease (IBD) in the hospitals across the United Kingdom.

**Methods** We conducted an internet based survey among the Paediatric Gastroenterologist, Paediatrician with interest in Gastroenterology and the Specialist Nurses in Paediatric Gastroenterology using Survey Monkey tool. Survey was conducted over a 3 month period from September 2012 to December 2012. Participants were sent a questionnaire regarding their case load, criteria for investigations for iron deficiency anaemia in IBD and modalities of treatments used for correcting iron deficiency anaemia. A total of one hundred health professionals were invited to participate in the survey.

**Results** The total response rate for the survey was 35%. 57% of the responses were from tertiary care paediatric gastroenterology

consultants who had a case load of more than 100 children with inflammatory bowel disease. Remaining 40% responses were made up by consultant paediatricians with gastroenterology interest and specialist nurses.

Less than half of the participants reported that they do routine screening for iron deficiency as part of IBD management. The lab parameters used to confirm iron deficiency varied significantly. In our survey, 28% of the participants were unsure or did not agree correcting iron deficiency anaemia alone will improve the quality of life in IBD patients if underlying disease control could not be improved. While majority of the participants (48.5%) used oral iron as the first line treatment, some (8.5%) were using IV iron as their first line. Our survey showed that only 11.4% of the participants reported that they had a local or regional guideline.

**Conclusion** Our survey highlights that there is significant variation in practise among the clinicians in methods of diagnosing and treating iron deficiency anaemia in children with IBD. It also underscores the need for evidence based national guidelines in this area.

## G202(P) A SURVEY OF NECROTISING ENTEROCOLITIS AT A TERTIARY NEONATAL UNIT

doi:10.1136/archdischild-2013-304107.214

<sup>1</sup>A Wilcock, <sup>2</sup>S Victor. <sup>1</sup>Manchester Medical School, University of Manchester, Manchester, UK; <sup>2</sup>Faculty of Medical and Human Sciences, University of Manchester, Manchester, UK

**Aims** Necrotising enterocolitis (NEC) is a gastrointestinal emergency occurring in approximately 1–3% of neonates admitted to intensive care units. It carries significant mortality (up to 50%) and an extensive range of short and long term complications.

Despite decades of research, its pathogenesis remains poorly understood, though current understanding suggests a multi-factorial aetiology. Indeed, prematurity, feeding practises, genetics, various maternal factors and certain neonatal morbidities have all been implicated in the pathogenesis of NEC. As there is a paucity of published surveys providing a general overview of NEC in the last decade, we aimed to provide a more current perspective in a tertiary neonatal unit.

**Methods** Using diagnostic criteria outlined in the 11th BPSU Annual Report, 49 infants with NEC within a 13 month period were retrospectively identified. Subsequently, data from patient records (including maternal data) and imaging reports was extracted and analysed. The presence of various risk factors (including those mentioned above) and information regarding the presentation, diagnosis and management of NEC in these infants were recorded along with complication and mortality rates.

**Results** Risk factors including intrauterine growth restriction (18.4%), patent ductus arteriosus (51%) and gastroschisis (10.2%) were highly prevalent within our cohort. Consistent with previous literature, 86% of infants were premature and 71.4% were very low birthweight infants (<1500g). However, 24.5% of infants were from multiple pregnancies which was higher than expected.

Intriguingly, whilst the majority of infants (75.5%) were exclusively fed with breast milk initially, by the time NEC was diagnosed this proportion had reduced dramatically (32.7%).

Gaseous distension was the most commonly seen radiographic feature (87.8%) whilst 36.7% received surgical interventions with nearly all undergoing laparotomy. Gut-related complications were observed in 20.4% of patients at discharge.

**Conclusion** This survey provides a more current insight into the progression of NEC from predisposing factors to current management and complications. In agreement with previous studies, no risk factor was uniformly observed. Of particular interest, however, is the trend regarding breast milk feeding. As breast milk has been shown to play a protective role in NEC, it is therefore imperative that support is offered to mothers to encourage continuation of breast milk feeding.

## G203(P) TRANSIENT TEMPERATURE GEL ELECTROPHORESIS OF STOOL SAMPLES OF PRETERM INFANTS IN A MULTICENTRE OBSERVATIONAL STUDY

doi:10.1136/archdischild-2013-304107.215

<sup>2</sup>K Brunner, <sup>1</sup>LM Beattie, <sup>2</sup>K Gerasimidis, <sup>2</sup>DJ Morrison, <sup>2</sup>CA Edwards. <sup>1</sup>Neonatal Unit, Royal Hospital for Sick Children, Yorkhill, UK; <sup>2</sup>Department of Child Health, University of Glasgow, UK

**Aims** Despite the effect of enteral feeding on the development of intestinal microbiota in preterm infants remaining poorly understood, trials aiming to prevent necrotising enterocolitis (NEC) using probiotics are well-established. Exclusively breast milk fed preterm infants have a reduced risk of developing NEC and this may be linked with a more 'beneficial' gut microbiota.

**Methods** The NAPI Study (see abstract BEAT82431) sequentially recruited infants <32 weeks and <1.5Kg birth weight. Non-meconium faecal samples from the first and fourth weeks of life in 22 infants, 12 with NEC, were analysed by PCR-Transient Temperature Gel Electrophoresis using universal bacterial primers. Species richness and similarities were compared between infants according to feed type: EBM, expressed breast milk, vs Mixed, breast and formula milk.

**Results** There was large variability between number (1–17) and species diversity (25–36 different species). Number of predominant bacterial species did not increase between the 1st and 4th week of life. Bacterial composition varied largely between the 2 sample points. No difference in species richness or similarity within the 2 feeding groups was observed. 4 bands were identified in >50% of infants. Intra-individual similarity varied greatly and ranged from a similarity index (Cs) of 0% to 66.8%. There was no statistical difference between the similarity indices of the feeding groups ( $p = 0.8852$ ) or between those with and without NEC ( $p = 0.1719$ ).

**Conclusion** Microbial community of preterm neonates undergoes several interindividual changes during their first month of life. The feeding mode did not seem to have a major impact on the development of bacterial diversity.

## G204(P) CAN TAUROLIDINE-BASED CATHETER LOCKS REDUCE CENTRAL VENOUS CATHETER RELATED BLOOD STREAM INFECTIONS IN CHILDREN ON LONG-TERM HOME PARENTERAL NUTRITION?

doi:10.1136/archdischild-2013-304107.216

L Howarth, E Gaynor, A Rodrigues, PB Sullivan. Department of Paediatric Gastroenterology, Oxford University Hospitals NHS, Oxford, UK

**Objectives and study** To compare the incidence and characterise the type of catheter-related blood stream infections (CRBSIs) in children with intestinal failure on long-term home parenteral nutrition (PN), using heparin-saline based catheter locks versus those using taurolidine-based catheter locks. There is growing body of evidence that taurolidine-based catheter locks, which have a broad-spectrum antimicrobial and antifungal action, is associated with a decreased incidence of CRBSIs children on home PN.

**Methods** All children referred to a tertiary paediatric gastroenterology service with temporary or on-going intestinal failure requiring long-term PN or preparation for home PN between 2005–2011 were included. Children were given a single-bag system of PN with each infusion via central venous catheter. Parents were formally trained in aseptic techniques and to instil heparin-saline or taurolidine-based solution into the catheter after completion of each infusion. CRBSIs were defined as a laboratory-confirmed blood stream infection from with a peripheral or central venous sample. Results were excluded if evidence that the source of infection was from a second site. All cultures results were confirmed through the microbiology database and clinical records. Research ethics committee approval was sought, but ethical review was not deemed necessary.