

Abstract PS-320 Table 1

Pollutants and Group incidence	Areas with traffic load >1.500	Areas with traffic load <300	P*
	motor vehicles/ hour	motor vehicles/ hour	
SO ₂ (mg/m ³)	0.148 ± 0.104	0.000 ± 0.000	<0.01
NO ₂ (mg/m ³)	0.045 ± 0.016	0.009 ± 0.008	<0.01
Particulate matter (mg/m ³)	0.274 ± 0.082	0.190 ± 0.079	<0.05
CO (mg/m ³)	3.317 ± 0.716	2.275 ± 0.411	<0.01
Annual incidence of croup in children (cases per 1000 child population)	10.78	1.08	<0.01

Table. Traffic load, traffic-dependent pollutants and annual incidence of croup in children (Vinnytsya, Ukraine, 2000–2003)

Conclusions Croup in children is related to traffic load and traffic-dependent pollutants (SO₂, NO₂, particulate matter, CO).

PS-321 BACTERIAL LOAD AND INFLAMMATORY MEDIATORS IN RELATION TO THE SEVERITY OF NECROTIZING PNEUMOCOCCAL PNEUMONIA

¹Y Hsieh, ¹YC Huang, ¹SH Lai, ²CC Liu, ³YC Huang, ⁴HC Lin, ⁵LM Huang, ⁶H Chi. ¹Pediatrics, Chang Gung Memorial Hospital, Taoyuan County, Taiwan; ²Pediatrics, National Chen Kung University Hospital, Tainan, Taiwan; ³Pediatrics, Kaohsiung Chang Gung Memorial Hospital, Kaohsiung, Taiwan; ⁴Pediatrics, China Medical University Hospital, Taichung, Taiwan; ⁵Pediatrics, National Taiwan University Hospital, Taipei, Taiwan; ⁶Pediatrics, Mackay Memorial Hospital, Taipei, Taiwan

10.1136/archdischild-2014-307384.620

Background and aims The incidence of necrotizing pneumococcal pneumonia has increased over the last two decades. We hypothesized that increased pneumococcal load or augmented inflammatory cytokine production may lead to destructive pneumococcal lung disease.

Methods This study prospectively enrolled children aged 0–18 years with a diagnosis of community-acquired pneumonia with pleural effusion admitted to 6 medical centres. Children were diagnosed with pneumococcal empyema if the pleural fluid tested positive for quantitative pneumococcal (lytA) detection by real-time polymerase chain reaction (RT-PCR). Pneumococcal empyema cases were further divided into four groups according to necrosis severity scaled by radiographic image findings: 0) non-necrosis, 1) mild necrosis, 2) cavitation, and 3) broncho-pleural fistula (BPF). Nasopharyngeal and pleural pneumococcal load, as well as proinflammatory cytokines (TNF- α , IL-1 β , IL-6, IL-8), Th1-(IL-2, IFN- γ), Th2-(IL-4, IL-10), and Th17-cytokines (IL-17) in the pleural fluid were measured.

Results Serotypes 19A and 3 accounted for 65.3% and 4.2% (respectively) of 72 cases of pneumococcal empyema. In multivariate analysis, pleural pneumococcal density (adjusted odds ratio [aOR], 1.79; 95% confidence interval [CI], 1.03–3.06), and IL-8 (aOR, 2.64; 95% CI, 1.21–5.75) were independent factors associated with the severity of lung necrosis. There was a good correlation between nasopharyngeal and pleural pneumococcal density ($\rho = 0.42$; $p = 0.001$). A lytART-PCR pleural density $\geq 50,000$ copies/mL had a sensitivity of 88.2% and a specificity of 70.9% for predicting broncho-pleural fistula.

Conclusion Evolution of *S. pneumoniae* toward increased fitness in their interaction with host and exaggerated IL-8 expression

are responsible for the increase of necrotizing pneumococcal pneumonia.

PS-322 A NEW SCREENING APPROACH FOR THE MANAGEMENT OF RSV INFECTION USING INNOVATIVE COMPUTERISED INFERENCE ALGORITHM TECHNOLOGY

¹T Takayama, ²Y Morikawa, ³E Inoue, ⁴I Hokuto, ¹J Ishihara, ⁵N Yahagi. ¹Pediatrics, Yokohama Municipal Citizen's Hospital, Yokohama, Japan; ²Clinical Trial, Tokyo Metropolitan Children's Medical Center, Tokyo, Japan; ³Clinical Research Education, National Center for Child Health and Development, Tokyo, Japan; ⁴Pediatrics, St. Marianna University School of Medicine, Kawasaki, Japan; ⁵Clinical Research Networking, National Center for Child Health and Development, Tokyo, Japan

10.1136/archdischild-2014-307384.621

Background and aims Minimising invasive laboratory testing on children is considered to be the top priority. Respiratory Syncytial Virus (RSV) is the leading cause of lower respiratory tract infection and the hospitalisation in infants. Identifying respiratory pathogens within the population is difficult because numerous invasive sample collections are required. Collecting precise information and estimating the severity of respiratory symptoms using the Innovative Computerised Inference Algorithm (ICIA) technology will minimise RSV screening tests.

Methods Children aged 1 month–15 years at Paediatric Emergency Department (PED) of Yokohama Citizen's Hospital (Yokohama, Japan) who were evaluated with respiratory symptoms had swab samples collected for the RSV test. ICIA prompts the guardian to input symptoms at time of registration, assessing the Disease Severity from 3 levels (mild/ moderate/ serious).

Results There were 23,851 PED visits from January 1, 2012 to December 31, 2013. Of those, 6742 patients had respiratory symptoms and swab samples were collected accordingly. There were 654 RSV infected patients. The severity level, over moderate was 632 and, mild was 22. These results proved that the sensitivity and specificity of ICIA were 96.6% and 87.9% respectively, negative and positive predictive values were 99.6% and 46.3% respectively.

Conclusion ICIA decreases 90% of invasive RSV tests. ICIA supports Paediatricians at each phase of their clinical decision making: i.e., diagnosis, severity assessment and treatment that used to depend on the amount of a paediatrician's knowledge and experience. Thus, ICIA leads to minimising the invasive RSV laboratory tests, in addition to shortening the time of clinical decision making process.

PS-323 THE GENE POLYMORPHISM OF IL-17 G-152A IS ASSOCIATED WITH INCREASED COLONISATION OF STREPTOCOCCUS PNEUMONIAE IN CHILDREN

¹J Vuononvirta, ²V Peltola, ³J Ilonen, ¹Q He. ¹Department of Infectious Disease Surveillance and Control, National Institute for Health and Welfare, Turku, Finland; ²Department of Pediatric, Turku University Hospital, Turku, Finland; ³Department of Medical Microbiology and Immunology, University of Turku, Turku, Finland

10.1136/archdischild-2014-307384.622

Background and aims *Streptococcus pneumoniae* is a common respiratory pathogen and up to 50% of children acquire *S. pneumoniae* in their nasopharynx during the first 12 months of life. The cytokine interleukin-17A (IL-17A) plays an important role in host defense against extracellular bacterial pathogens.