ON GROWTH AND DEVELOPMENT*

PART I. ANATOMICAL CONSIDERATIONS

BY

RUSTIN McINTOSH

From the Babies Hospital, the Pediatric Service of the Presbyterian Hospital, and the Department of Pediatrics, Columbia University College of Physicians and Surgeons, New York

The invitation to give the Sir Leonard Parsons lectures carries high honour. For me the invitation has a particular appeal, the honour a special worth, because I had the privilege of knowing Dr. Parsons for some 20 years. Our first encounter was in Sweden in 1930 when we happened to be making our way toward the International Paediatric Congress by the leisurely means of the Gothenburg-Stockholm canal ship. Professor Smellie remembers it well, too, for he accompanied Dr. Parsons and, like most of the passengers on board, was exploring those unfamiliar Scandinavian waterways for the first time. In that heterogeneous assembly of tourists, professional people, and business men who comprised the passenger list, the customary inclination of the individual traveller to keep to himself, to play his hand cautiously while fixing an appreciative eye on the scenery and awaiting the turn of events, was on its way to establishing standard procedure. Indeed, one might at first have inferred that the only plausible bonds among the various groups were membership in the human race and a common admiration for the beautiful countryside bordering the Gotha Canal—scarcely enough, indeed, to provide footbridges of friendliness across the chasms of reserve which separated one from another. Not so, however, for the Englishman of alert and energetic air, remarkably youthful for one whose hair was snow-white, about whom a cluster of friends seemed to be constantly growing and who had the knack of expressing, or transmitting somehow without verbal expression, a sincere interest in the aims and concerns of even the newest acquaintance. We met by the process of mutual self-introduction, still much less of a shock for an American, I suppose, than for a Briton in spite of the changes in social custom that have taken place over the past generation. In my luggage I had with me a letter of introduction written on my behalf

by my former chief, Dr. E. A. Park, which I had counted on presenting in due course; and on his part Dr. Parsons had received word from some friend of mine, asking him to keep an eye out for me since we might be brought together at the Stockholm meetings. Neither form of introduction proved necessary. Dr. Parsons combined perfect courtesy with an ability to overcome diffidence, in a manner which equipped him admirably for the role which he played so well and which it is quite certain he greatly enjoyed, that of ambassador without portfolio. This was enhanced by the hospitality of his home, over which Lady Parsons presided with grace and warmth, and by the renown of the Children's Hospital, which largely by virtue of his stimulating participation supported an active and highly effective programme of clinical research. Over the years Sir Leonard succeeded in influencing, in helping, in uplifting an extraordinarily large number of people in widely separated areas of the world. My own debt to him is great indeed.

Dr. Parsons's keen interest in problems of growth and development must have guided in some measure the choice of subject which was assigned for these lectures. Indeed, the impetus which he gave to studies of the many-sided evolution of the individual through infancy, childhood and youth toward maturity reflects his appreciation that the effort to understand thoroughly the growing person calls upon every resource of the clinical art. In this conviction, as in many others, he was ahead of his time.

Growth is a conspicuous characteristic of the young of any species, so much so that one whose attention is occupied with infants and children finds it all too easy to take growth for granted. The natural expectation that a young person will continue to increase in stature and weight, and to advance at the same time in maturity, is fulfilled so often in the face of a variety of everyday vicissitudes that one too readily believes that growth will take

* The Leonard Parsons Lectures delivered in Birmingham on May 7 and 8, 1957.

care of itself. All that is required, however, to prick this complacent bubble is to ask oneself whether a given individual, an infant or a child, is growing as well as he can and as he should. Here we come squarely up against a formidable pair of questions, How does one determine optimal growth, and how can it be attained?

At the outset it must be admitted that we have no way of knowing what is best in the matter of growth. We can establish that one rate of growth is faster than another. Admitting the risk of a non-sequitur, we may even assert that a new achievement in growth, being demonstrably faster, is therefore better—an assumption one would find difficult indeed to prove. Even so, it is initially only too clear that in this quest of the absolute we cannot expect in any single step to close more than part of the gap which separates what we know to be attainable from that which we regard as perfection. Only in the course of infinite time can the best in growth be approached as a limit. And the closer we come to our goal, the more difficult it becomes to distinguish the results of our own manipulations from the operation of pure chance. The student of growth depends heavily on biostatistical method and on tests of significance to protect himself from false deductions.

The clinician, appreciating that in the immature animal growth is in some degree a measure of health, seeks a healthy population in which to attempt a quantitative definition of satisfactory or exemplary growth, and in doing so comes up at once against the related question, What is health?—or, specifically, Who is healthy? It is all very well for the World Health Organization to define health as something more positive than the absence of disease; but the identification of the truly healthy child in objective as well as positive terms becomes a formidable task unless one is willing to beg the question, come around full circle, and settle for the answer that he is a child who is growing properly.

Inferences and deductions based on studies of growth in experimental animals or in husbandry have, to be sure, limited direct application to the interpretation of growth in man. At the same time, they have served an important function in defining hypotheses which may then be put to the test in the study of infants and children, either by manipulation of the environment or by analysis of experiments of nature. It will be found that a great many aspects of the vast problem of nutrition in its relation to growth prove downright misleading when transferred from the experimental laboratory animal to man. Certain phases of the relationship of endocrine gland function to growth, although well

defined so far as they apply to experimental animals, have yet to be confirmed in their application to human subjects. A conspicuous example which immediately comes to mind is the huge discrepancy between the experimental demonstration of the role of the pituitary growth hormone in the rat and the inconclusive evidence for a comparable role (at least with the currently available products) when efforts are made to accelerate the growth of a child afflicted with what we regard as hypophyseal dwarfism. In general, then, reliable data on growth in man are to be sought almost exclusively in the study of man.

Linear growth as a biological event has its basis in three fundamental phenomena: cell multiplication, increase in cell size and accumulation of extracellular material. Each of these processes in certain ideal circumstances follows a course which can be described by a relatively simple mathematical equation. Human growth, however, expressing the result of a variety of processes taking place simultaneously, is not easily pinned down in concise mathematical terms. The reasons are several: for one thing, differentiation of tissues begins not long after fertilization of the ovum, and single tissues grow at different rates; again, size and shape of the body are determined only in part by aggregates of cells, since a large and almost continuously variable contribution derives from inert products or accompaniments of cellular activity; and finally, the momentum of growth in the individual is subject to a variety of influences, the effects of which can be anticipated only imperfectly. As a result, the curve of growth as actually observed and recorded reflects the push and pull of a number of forces; and if it happens to coincide fairly well with the plot of an equation of growth or with a succession of such equations, this conformity ought perhaps to be looked on as something of a mathematical accident. One recoils instinctively from the notion that mathematical elegance in studies of human growth is to be purchased only at the price of repeated shifting of formulas as well as of parameters. Such faithless promiscuity indeed departs so far from elegance as to approach vulgarity. Medawar (1945), in one of the essays on growth presented in 1945 to D'Arcy Thompson, struck a proper balance between theory and empiricism in studies of growth when he wrote:

'Those who attach great importance to the concrete forms of growth equations are not as a rule put out when the whole course of growth has to be resolved into two or more separate periods or cycles, each to be described by special functions, or by a single function which abruptly changes the values of its parameters at what can here truly be

called the psychological moment. . . . There is little in the development of the chicken to suggest that its life is really punctuated by these radical metamorphoses.'

What he said of the growth of the chick applies to man also.

The difficulty of formulating any universal mathematical law of human growth is pointed up by consideration of those studies which undertake to compare the growth of any one tissue or portion of the body with that of other segments or of the body as a whole. The fact that differential growth rates of individual tissues appear early and continue throughout foetal life has been well documented by Scammon (1930), by Clatworthy and Anderson (1944) and by others. Plots of the weight of individual foetal organs or parts, measured against foetal age, show little parallelism among the various curves. Similar studies of foetal stores of fat, protein, water and various mineral constituents (Widdowson and Spray, 1951) show comparable divergence. In healthy infants, Forbes and Perley (1951), Ely and Sutow (1952) and Friis-Hansen (1955) have employed a variety of devices for measuring the total water content of the body at different ages and have estimated both intra- and extracellular water in relation to postnatal age. Growth of the various fluid compartments of the body runs fairly parallel to that of length or weight. On the other hand, growth of total body fat follows a curve of its own. Throughout infancy and childhood girls tend to have more subcutaneous fat than do boys of like age (Kornfeld, 1957; Reynolds, 1950), a circumstance for which girls often have reason to be thankful when sea bathing is the popular sport. The diversity of growth trends and growth rates in such selected tissues as haematopoietic marrow (Osgood, 1955), fingernails (Babcock, 1955), pituitary fossa (Acheson, 1954a), head width (Meredith, 1953) and long bones of upper and lower extremities (Maresh, 1955) sounds a warning that each aspect of growth deserves individual study if experience is to have predictive value.

From clinical observations, from the studies of anthropologists, from experiments in biological laboratories or from knowledge of the facts of life as seen on the farm, one can readily assemble a list of factors which are known to influence growth and development or which are suspected of having such an influence. The measurement of these influences, their evaluation in quantitative terms, is of practical significance to clinical medicine. At the same time, the task is one of formidable difficulty in view of the large number of variables which operate simultaneously.

Without attempting at the moment to assign relative values to any of these factors which are known or believed to affect growth, it may be of some use to list them in rather general terms in order to have them in mind for later comment. On a broad scale they can be divided into three groups: (1) Factors which are genetically transmitted; (2) circumstances of the external environment; and (3) conditions of the internal environment. Concerning each one, and indeed concerning subdivisions of each, either there has been evidence advanced indicating an effect on growth, the magnitude of which only remains to be more precisely measured, or there has been fruitful speculation leading at times to the collection of convincing data.

Taking growth determinants in the order in which I have just given them, I must comment first on genetic influences. Phenotypic sex is genetically determined, by and large, save for the few interesting exceptions which have come to light in recent years as a result of applying Barr's methods (1956) to the identification of genotypic sex according to nuclear chromatin pattern. For the most part sex runs true to form, so to speak, being determined by the chromosomal equipment of the particular spermatozoon which fertilized the ovum, whether it contained the X or the Y chromosome. Sex differences in growth rate appear even before birth, male foetuses being heavier and females being more advanced in bone maturation. These sex differences become quite pronounced somewhere around the sixth to eighth year when in many girls the prepubertal spurt begins to make itself known. The earlier onset of the pubertal growth acceleration in girls as compared with boys, and its earlier termination, are well documented phenomena. The greater deposition of subcutaneous fat in girls has already been mentioned. Sex differences are recognized, and there is no risk of their being lost from sight.

Concerning the effect of race, ethnic group, or whatever might be thought of as the human equivalent of what we call breed in domestic animals, it has proved difficult indeed to assemble unequivocal data. The necessary balancing of environmental factors such as food supply, disease experience and general living conditions has been omitted when measurements have been made of a circumscribed population living in its natural surroundings; and when through migration environments have been equalized, or nearly so, inter-marriage over the course of years has almost invariably beclouded genetic interpretations. Many studies of growth carried out over the past half century begin hopefully with the statement that the group of children

studied was selected from families of 'north European stock'. And what, exactly, does that signify in genetic terms? Indeed, in a number of careful growth surveys carried out in the United States (Meredith, 1941, 1951a, 1951b), where the difficulties of defining and interpreting so-called racial traits are as formidable as they are likely to be in any part of the world, the conclusion was reached that differences in mean stature and mean body weight found between contrasted population groups could not be proved to be due to racial, rather than environmental, differences.

Genetically transmitted traits other than racial characteristics may influence growth in other respects. Tall families and short families frequently run true to form—so true, in fact, that the appearance of a deviant of merely average stature will often set unthinking tongues to wagging. Retardation of growth associated with an inherited metabolic defect such as galactosaemia or the hepatic form of glycogen storage disease is well recognized. What is less clear is the extent to which subclinical deviations in functional performance may modify growth capacity. The need for additional study along these lines is obvious, and no student of the problem underrates its difficulty.

Among pertinent factors of the external environment, the first item I have listed is nutrition. In recent years the number of publications describing the impact of dietary factors on human growth has shown a decrease which marks a shift of the focus of interest. Alterations in the growth of children dependent on war or on widespread economic depression are closely correlated with the intensity of the fiscal disturbance, have been well documented, and call for political and social adjustment rather than for special attention from the medical profession (Howe and Schiller, 1952). Of more direct concern to students of physiology are the efforts aimed from time to time at rendering an adequate diet even better by addition of an essential substance which is believed to be present in marginal quantity. For example, claims have been made for a growth-accelerating effect of supplementation of a conventional diet with small increments of vitamin B₁₂ (Larcomb, Perry and Peterman, 1954). In view of the small numbers included in the study, and the probability that differences in stage of maturation at the particular age level of the test subjects complicated the problem of sampling, the conclusions expressed by the authors may well be interpreted with some caution. Comparable scepticism must be applied to claims voiced by Mülhausen (1953) for Goetsch's T-vitamin. In this connexion mention should be made of the demon-

stration by Baptist and de Mel (1955) that Ceylonese children reared on a cereal-legume-vegetable diet grew as satisfactorily as did a test group for whom a supplement of milk protein was provided. Their subjects were between 1 and 6 years of age. Earlier reports of similar studies carried out by Dean (1951) and Dean and Schwartz (1954) both in Germany and in Uganda seem somewhat more candid in recognizing the technical difficulties encountered.

The superiority of an adequate diet over an inadequate one is perhaps less subtly measured in terms of growth than by other criteria, particularly biochemical criteria, which can define the inadequacy more accurately than is feasible by conventional measures of external dimensions.

Macy, whose long experience in meticulous studies of nutrition endows her judgments with unusual claim to respect, has tilted a lance against the instinctivists and has asserted that appetite in children is not to be trusted to assure appropriate growth even in the presence of ample food and in the absence of known infection or other disease. In a series of experiments published in 1951 in association with Hunscher she came to the conclusion that for some subjects the voluntary intakes of food were not providing sufficient calories for satisfactory weight gains; and she stated that 'neither appetite nor any of the known standards can be relied upon to provide adequate food intakes for individual children' (Macy and Hunscher, 1951). In view of the fact that most of her subjects were between 5 and 10 years old and therefore at a highly impressionable age, one cannot refrain from wondering whether any of them were unhappy in the conditions of the experiments. Against these more modern studies may one not cite the observations of Clara Davis (1928) published long ago which led her to exactly the opposite conclusion, namely, that infants and young children who were permitted to select their own diet, qualitatively and quantitatively, from both variety and abundance set before them surpassed in their growth over a year or more the accepted normal standards of those times. Widdowson's observations (1951) in German orphanages showed convincingly that a child's affective reaction to supervisory personnel clearly influenced his weight gain, overbalancing the effect of minor differences in caloric intake. A comparable response in statural growth, although less prominent, was none the less detectable.

The suspicion that geographical location affects growth rates seems implicit in many careful studies of the problem. The device of sending a convalescent off to some remote region for a change of

climate in the interests of his health is firmly established in clinical thinking and practice. A comparable approach to the growth question sounds distinctly plausible. Yet as a thesis it is obstinately resistant to rigorous proof. For some years in the United States there lurked the suggestion that the growth rates found in studies made in communities of the Middle West and the Far West, more rapid than those recorded from cities of the Atlantic seaboard, supported the advice which Horace Greeley had given so often and so emphatically a century ago: 'Go west, young man.' One rather expected a Texan or a Californian to endorse the notion that out in God's country people naturally grew bigger as well as better. But in these comparisons longitude was not the only variable. True enough, when it became apparent that Boston children of 1879 or New York children of 1899 were being matched age for age with groups from other areas, geographic location appeared to exert less influence than did the so-called secular trend, the gradual acceleration of growth over several decades. Meredith, one of the most careful and critical students of growth, reviewed the entire subject in 1941, so far as American data were concerned, and came to the conclusion that no clear difference could be established between the growth rates of children reared in urban as contrasted with rural surroundings, nor between rates from different pairs of urban communities chosen from widely separated and representative parts of the country. More recently Newcomer and Meredith (1951) published observations made on a 1950 sample of 15-year-old boys from Eugene, Oregon, in which the means for both weight and height exceeded those of any other recorded community; but the samples were small, and with the caution that comes of long experience they slyly refrained from expressing an opinion regarding their significance, so far as the geographical factor was concerned.

Variations in growth rates with season of the year have been recorded fairly frequently (Palmer, 1933; Reynolds and Sontag, 1945). In studies carried out in Maryland, gain in weight per calendar month reached a peak in the month of October. Seasonal acceleration of growth in height is more subtle and considerably more difficult to demonstrate. It seems impossible to determine with available evidence whether such fluctuations depend directly on the solar cycle, responding to the influence of daylight, sunshine, environmental temperature, and such; or whether they are indirectly related to the solar angle and reflect the influence of changes in diet, in school demands, in vacation mood. In at least one study from the southern hemisphere, that

of Fitt (1924), the peaks and dips in the curve of weight gain are in reciprocal phase, showing maxima in April, May and June and minima in October, November and December.

Great interest continues to be focused on evaluation of the secular trend, the acceleration of growth and maturation which children of today exhibit in comparison with similarly selected children of past generations. Of the phenomenon itself, the greater height, weight, and other measurements, of children of a given age when matched against the records of children of the same age in their fathers' time or earlier, no serious doubt persists. Whether adult height is likewise increasing with time remains somewhat in dispute, and adult weight has too little finality about it to deserve elaborate statistical analysis. But in respect to stature, even if one concedes for purposes of argument that today's adults are no taller than their forebears, the secular trend in the growth of children is reflected in a progressive shifting to the left of growth curves on the time scale (Meredith, 1941; Weir, 1952; Clements, 1953). Clements (1953) summarizes the trend over a 70-year interval by showing that in height and weight the 6-year-old of 1953 is at about the same stage as the 7½-year-old of 1883. A variety of explanations have been offered. Berry and Cowin (1954) point out that the increase in growth rate is accompanied by slimmer physique. Bennholdt-Thomsen (1942) postulated selective breeding and selective survival of taller, more slender types, and this position has been staunchly supported by Freund and Maier (1952), though one is hard put to it to understand how these ordinarily slow-moving forces can produce in so short a space of time the results actually found. Most students of the secular trend have regarded it as the net expression of a variety of forces including availability of food, understanding of nutrition, relative freedom from significant infectious disease, purchasing power of family units, housing, public sanitation. Nevertheless, the paradox remains that as each single item is individually tested for significance by as careful matching of samples as the data and changing conditions of the times permit, as Meredith (1951a), Weir (1952), Clements (1953) and others have done, a large fraction of growth acceleration remains unexplained. Dissection of the secular trend is, as Meredith (1951a) and Morant (1951) in particular have emphasized, a promising assignment for future studies.

Evaluation of the internal environment in its bearing on growth and development amounts almost to a paraphrasing of the paediatrician's clinical responsibility to his patient. Simply for the purpose

of establishing broad subdivisions I have listed four spacious territories where the hunting is apt to be rewarding, namely, the regions of (1) disturbances of utilization of structural material, (2) endocrine imbalance, (3) malformations of one sort or another, and (4) infections of all kinds. There is little point in attempting to survey in detail the entire area in which the search for causes of disturbed growth must be made, for it is all familiar territory to clinicians and has been well set out in a number of textbooks in common use. A word may be said, however, about trends in the study of specific problems of growth.

Without much question styles in clinical investigation have changed over the past few years; new ideas and a new freedom have been introduced, and the accuracy of identification of causes has increased impressively. Emphasis has shifted from concern over the minutiae of the diet to a more careful appraisal of the patient himself. The trend in feeding practices toward encouraging a child to be omnivorous from an early age has amply demonstrated that excellent growth may be attained with a quite wide variety of menus. A deficient diet can still account for slow growth, but this is seldom encountered nowadays in your country or in mine save when ignorance and superstition have combined with economic restrictions to limit the child's freedom of selection. And with the shifting of focus to the patient and the increasing application of modern diagnostic aids, new causes emerge as the agents responsible for retarded growth: specific enzyme deficiencies such as galactosaemia or hepatic glycogen storage disease; the endocrine imbalance which accompanies the fundamental gonadal disturbance in Turner's syndrome; coarctation of the aorta, pancreatic fibrosis, a variety of anomalies of the kidney; and such vaguely localized functional curiosities as the severe form of idiopathic hypercalcaemia (Schlesinger, Butler and Black, 1956). The only purpose of this fragmentary enumeration is to emphasize the point that over a relatively short space of time such as the past 20 years, means have not only been found but also popularized and made widely available for the accurate identification of responsible factors in problems of growth and development. Save for the reflection that perhaps the richest ores have already been mined and that those who come after may have to dig deeper and employ more elaborate refining processes, there is no reason to anticipate an important slacking off of progress in finding explanations. Finding remedies is, of course, another matter.

Thus far I have chosen to rehearse in summary fashion and in general terms some of the factors

which influence growth—linear growth, increase in weight—without giving due attention to maturation or development. All of the causative agencies just referred to exert an analogous effect on maturation, usually in the same direction or with the same sign.

Maturity has been defined as the state of complete natural development, and degree of maturation may be looked on as a station along the way. Let us first consider skeletal maturation. The task of evaluating it is usually handed over to the radiologist, who studies whatever radiographs he considers appropriate—the hand and wrist, perhaps a knee or foot, or even one-half of the skeleton—and, taking account of the patient's sex, reports back that skeletal maturation is within normal limits for a boy or girl of stated age, or perhaps is abnormal. In the latter event, if the skeletal maturation is alleged to be retarded, or if it is advanced, the question of degree is certain to come up.

Several methods are in common use for evaluating skeletal maturation. In one, films of the patient's hand and wrist are compared with standards contained in an atlas, perhaps Todd's original atlas (1937), or the later modification published by Greulich and Pyle (1950). With either standard of reference, the patient's film is graded as being equivalent to a certain bone age for the corresponding sex. Todd himself had apparently expected two assessors of unequal experience to rate maturation within about six months in a given case. Those who have used his atlas subsequently have often appeared to be somewhat more confident of their ability to pinpoint a maturational rating. In recent studies (Dreizen, Snodgrass, Webb-Peploe, Parker and Spies, 1957) made on white children living in Alabama, it was found that when the two hands were compared, the skeletal age of the right hand exceeded that of the left hand in 52%, was equal in 26%, and was less than that of the left hand in 22%. The greatest disparity was found in the epiphysis of the radius where the two sides got an equal rating in only 22% of the cases. In respect to individual centres in the carpus, the two wrists of a given individual might justify a reading differing by as much as 12 months' bone age or even more.

Mainland (1953, 1954) carried out an important study of both the systematic errors and the variable errors inherent in the method of estimating skeletal maturation by comparison of the patient's radiograph with a standard atlas. Pressing his wife into service as an assessor without previous experience, and therefore presumably without prejudice, he persuaded her to evaluate bone age in several films of the hand, the only information furnished being the sex of the subject. After an interval of several

months, the same films were reassessed in a different sequence. She then compared her readings with those which had previously been made by Todd and his collaborators on these identical radiographs. Later, she repeated the same procedure using the Greulich and Pyle atlas. On the whole, she tended to read skeletal ages lower than did the experts. Her systematic error was quite variable. The conclusion was drawn that the method was better adapted to determination of the average skeletal age of a group of children than to the assessment of bone maturation of an individual child, especially from a single film. The variable error was calculated from the scores made by the same assessor on different occasions, sufficiently separated in time so that memory would play no part. The irregular variation found between independent readings of the same film amounted to a standard deviation of three or four months. This means that, depending upon the age of the subject, the assessor would have to attach an error of ± 8 to 11 months to his estimate of skeletal maturation in order to achieve 95% significance. In other words, one must not look for greater accuracy than the method itself admits; and perhaps some caution is warranted before one accepts a maturation rating at its face value.

Another method consists in the counting of centres of ossification in one-half of the body, a technique popularized by Elgenmark (1946) and by Harding (1952). Essentially it is an extension of earlier techniques in which centres were counted only in the hands and feet. The method has the great advantage of objectivity in scoring, since the presence of a calcified centre, rather than its pattern, constitutes the basis of evaluation. In common with the atlas method, this one has the advantage of making it relatively easy to state whether a given patient's bone maturation is accelerated or retarded, but the disadvantage of making it difficult to express this deviation in maturational units.

The list of procedures for evaluating skeletal maturation could be extended if there were much point in doing so. The large number of different techniques advocated and in use indicates that none has been found wholly satisfactory—an all too common situation in clinical therapeutics. I am well aware that this question has enlisted the interest of Acheson (1954b) and his group (Acheson, Kemp and Parfit, 1955) at the Social Medicine Unit of Oxford University, and it is to be hoped that their exploitation of the system of scoring by 'Oxford maturity units' will overcome the difficulties experienced with other methods of assessment. By scoring both hands for each evaluation they do indeed avoid the irregularity which results from

chance inequalities of bone pattern in the two hands. Careful as they have been in the timing of the ages of their subjects who have served to set the pattern, it is not yet entirely clear whether the increment in maturity from, say, 5 to 6 maturity units, is equivalent to the increment from 20 to 21 units. In other words, has linearity been fully established? Another practical difficulty, soon to be resolved no doubt, is that thus far the published standards extend only up to age 6 years. But the basic principle of scoring maturation in maturity units or in fractions of ultimate maturity, and not in time units, represents a distinct advance over the older and more prevalent device of registering bone maturation in terms of so-called bone age.

Other means of estimating maturation are available besides the pattern of bones as seen in radiographs. The eruption and exfoliation of teeth follow trends which enable one to estimate so-called dental age objectively and quite accurately. The bodily changes which mark the process of sexual maturation normally follow fairly well defined sequences and lend themselves to semi-quantitative evaluation. Likewise, patterns of neuromuscular behaviour and of intellectual attainment ordinarily show a correlation with chronological age which provides a useful tool in the assessment of an individual child's functional capabilities, if one is looking ahead, or of his achievements in neural organization and mental growth, if one is more interested in measurement of past attainment. Independent evaluation of these several functions is of enormous value in the diagnosis of endocrine disturbances like hypothyroidism, and in the titration of results of replacement therapy. Indeed, every well organized endocrine clinic relies heavily on a battery of tests of maturation in following examples of retarded or accelerated growth and development of all sorts, for it is in these endocrine clinics where, rightly or wrongly, such patients usually land. Every clinical investigator who has probed deeply into problems of disturbed growth and development (Prader and Perabo, 1952; Talbot, Sobel, McArthur and Crawford, 1952; Wilkins, 1950) has focused due attention on what might be called differential maturation. Perhaps the odd feature of current practices in this regard is the limited number of fields which the clinician concerned with problems of growth has been willing to cultivate intensively. It seems probable that future efforts will devote more attention than is now customary to the wide variety of functional aspects of development which invite exploration, as contrasted with the rather limited anatomical territory

in which the search for measurable components now tends to operate.

Careful studies of the complex processes of puberty and adolescence, of which a number have been organized in the past few decades, have thrown fresh light on growth and development from several points of view. It has become clear, for one thing, that a secular trend in earlier sexual maturation of girls is discernible over broad spans of time such as 50 or 100 years (Brock, 1954). A comparable acceleration in sexual maturation of boys, although less firmly substantiated, is nevertheless highly probable. With earlier maturation goes earlier appearance of the prepubertal growth spurt; and while this precession of the normal childhood acceleration of linear increase is readily demonstrable in the average behaviour of large populations, its effects are most conspicuous when measured in the individual child. Earlier onset of the growth spurt goes hand in hand with earlier attainment of maximal rate and also with earlier attainment of the limit of linear growth. In other words, the secular trend shows both boys and girls reaching mature height at a younger age: for boys, according to Clements (1954), at age 17 years 9 months on the average, with a standard deviation of close to 10 months; for girls, at a mean age of about 16 years 3 months with a standard deviation of 13.6 months. Evaluation of skeletal maturation with relation to age and attained height has permitted certain students of growth problems (Bayley, 1956; Bayley and Pinneau, 1952; Tanner, 1955) to predict ultimate height with greater accuracy than would be possible on the basis of age and height alone. In the management of problems of endocrine imbalance associated with dwarfism, maturational aspects of development are accorded fully as much attention in the assessment of treatment as are the simple linear aspects of growth. These considerations have become all the more important since pharmacological agents now available have sufficient potency to accelerate skeletal maturation to a harmful degree in that, by speeding up the process, they can actually cause fusion of epiphyses, and therefore cessation of growth, before the patient's inherent growth potential has been realized.

Now, although most of the factors which influence growth have been touched only lightly and many have not been mentioned at all, it is appropriate to return to the basic clinical situation, the physician confronted with a patient, a growing patient. With respect to the patient's growth status the physician formulates a hypothesis which takes one of three forms: (1) Up to now the child has grown

adequately; or (2) his growth has been insufficient; or (3) it appears to have been excessive. The hypothesis is now to be tested by measurement of the patient and matching his measurements against some standard of reference. Most doctors have messed around sufficiently with tape measures, scales and clocks to understand the technique of mensuration and to know when and how mistakes are commonly made. They realize that every measurement involves an error of mensuration, and they understand fairly well that when two comparable measurements are added together, the error of the sum is somewhere between the smaller of the two individual errors and twice the larger of the two. I suspect it is not so clearly appreciated that when one measurement is divided by another, in order to obtain a ratio, the error of the quotient may be very much greater than the error of the sum. This risk of self-deception applies to such apparently innocent procedures as obtaining a cardiothoracic index, or the ratio of the linear measurements crown-to-pubis over pubis-to-heel.

At this early stage of evaluation, in the first-order testing of the three simple, basic hypotheses just mentioned, what dimensions need to be measured? Is it enough to weigh the patient and to record a single linear index such as standing height or recumbent length? Or should one follow the lead of the students of growth and include sitting height, span, head width, head circumference, shoulder width, intercrystal diameter, intertrochanteric diameter, thickness of subcutaneous tissues, and so on? Practices do indeed vary widely. On the whole, I would defend the simplest sort of evaluation at this stage, leaving until later the more meticulous and detailed measurement in the event that examination did actually bring out evidence of growth disturbance. Every physical examination of an infant or child should comprise a sort of battery of assessments—of body build, organ size, dental maturation, neuromuscular development, even in a rough way mental competence, all with relation to attained age. Any unusual findings turning up in the course of a thorough clinical evaluation will then point the way toward those specific dimensions which one will later want to watch with particular care. As I review my own sins of omission I have no inclination to regret that I have refrained from subjecting hospital and clinic patients to detailed routine measurements merely for the sake of accumulating data, but I do wish that some of the time thus saved had been devoted to a more detailed exploration of the course of growth in those patients whose clinical problem was found to include impairment or acceleration of growth.

What standard of reference is now to be used in judging the adequacy of growth? Ideally, of course, the frame of reference for the individual patient ought to be a healthy population of his own general group. In most situations no such standard is available, and the clinician must then ask himself how much difference it makes in actual practice to refer to whatever growth tables are at hand. If the inquiring physician happens to be over-conscientious and consults more than one set of tables, he is almost certain to find divergent answers and to become progressively more confused and unhappy the farther he searches. At this stage of the assessment of a clinical problem it seems to me to make little difference whether the frame of reference used is new or somewhat less than new, provided it has been so designed as to afford a convenient way of estimating, either through calculated percentiles or through multiples of the standard deviation, the degree to which the patient under scrutiny varies from the given mean for his age and sex. Tanner (1952) has put the problem in proper perspective by pointing out that one's criteria for separating normal from abnormal depend largely on what is planned as the next step: if the next step is to be closer study of the patient, then one will be far more willing to classify a measurement as abnormal than if the decision is tantamount to relegating the patient to some corner of limbo. In this respect it is a fortunate accident of history that in growth matters the standards of reference, the available tables of height and weight for age and sex, have usually lagged some years behind the uses to which they are put; and in view of the secular trend toward acceleration, the net effect has been to provide a somewhat spurious satisfaction with modest achievement. When the practical outcome of the evaluation of new measurements by old criteria has been to allay suspicion in respect to marginally retarded growth, the resulting tranquillity has no doubt been all to the good, so long as really pressing problems are not entirely overlooked. And on the other side of the coin, no great harm is done when the practitioner pats himself on the back because the great majority of patients under his care are above average in their growth attainment, again by comparison with outdated standards of reference.

Once it has been admitted that the individual child presents a problem in growth, because his achievement thus far leaves something to be desired, the next questions, How fast is he growing, and Will his rate of growth enable him to close the gap eventually? are of a quite different order of technical complexity. Students of growth (Meredith and Meredith, 1950; Shuttleworth, 1951; Tanner,

1952) extol the study of growth rates and point out how much more is to be learned from a knowledge of the speed of growth at a given age than from the mere attainment of length or whatever other dimension in relation to chronological age. At the same time it must be kept in mind that with inaccurately measured linear dimensions separated by too short a time interval the error of measurement can easily lead to absurdity. While I am quite prepared to agree that the most accurate evaluations of growth progress are attainable through careful determinations of rate in relation to stage of maturation, it is perhaps permissible to express a word of caution to the clinician who might be tempted to approach this assignment lightly.

Rate of growth in relation to stage of maturation—this is really the gist of the problem of assessment of the individual. It sounds easy, actually it is difficult, and yet even with all its possible sources of error it can be rewarding in its prediction of later performance. The accuracy with which chronological age can be measured, to four, five, or even six significant figures, must not be allowed to lure us into excessive faith in the value of the time dimension when we are trying to learn whether the patient before us has a favourable chance of growing into a pleasing and effective adult. At any given age the range of acceptable variation about a mean value may be quite large—more for some dimensions than for others. For boys 3 years of age, for example, the coefficient of variation $\left(\frac{100 \text{ S.D.}}{\text{Mean}} \right)$

for height is of the order of 3, for weight of the order of 10 (Thomson, 1956), and for skeletal maturation by the Elgenmark technique of the order of 30. The paradox is that in our actual evaluation of the patient, especially if he shows evidence of endocrine imbalance, our predictions and sometimes our diagnostic judgments rely fully as much on our estimate of maturation, the most variable of the dimensions measured, as on any of the other parameters. Again, one must plead for more intensive study of all aspects of maturation in order that our tools of measurement may be brought to ever finer calibration. In the assessment of the problems of adolescents, in whom the vagaries of sexual maturation are closely bound up with emotional adjustments as well as with simple physical growth, we have to recognize that much remains to be accomplished. In spite of the meticulous studies of Bryan and Greenberg (1952), of Hogben, Waterhouse and Hogben (1948), of Shuttleworth (1951) and others, it would seem premature to claim that the measurement of sexual maturation has been put on an altogether satisfactory quantitative basis.

Growth and development encompass too large an area to permit one to summarize the current state of our information with any sense of contentment. In a condensation such as I have attempted, omissions are unavoidable, and distortions are certain to intrude. I have tried to bring to the fore those features which seem to me of relative significance in the clinical evaluation of a patient's status and prospects—in his anatomical assessment at a time when he is still short of full maturity. In the course of the past several years improvements have been made in the techniques of measurement, though most of these are not of major importance. A more significant advance lies in greater acceptance of the range of expected variation consistent with health, in willingness to consider a variety of possible causes when growth disturbance is recognized, and, finally, in appreciation that the patient himself deserves intensive study as well as his background and his environment.

REFERENCES

- Acheson, R. M. (1954a). *Brit. J. Radiol.*, 27, 298.
 — (1954b). *J. Anat. (Lond.)*, 88, 498.
 —, Kemp, F. H. and Parfit, J. (1955). *Lancet*, 1, 691.
 Babcock, M. J. (1955). *J. Nutr.*, 55, 323.
 Baptist, N. G. and de Mel, B. V. (1955). *Brit. J. Nutr.*, 9, 156.
 Barr, M. L. (1956). *Canad. med. Ass. J.*, 74, 419.
 Bayley, N. (1956). *J. Pediat.*, 48, 187.
 — and Pinneau, S. R. (1952). *Ibid.*, 40, 423.
 Bennholdt-Thomsen, C. (1942). *Ergebn. inn. Med. Kinderheilk.*, 62, 1153.
 Berry, W. T. C. and Cowin, P. J. (1954). *Brit. med. J.*, 1, 847.
 Brock, J. (1954). *Biologische Daten für den Kinderarzt*, 2nd ed., Vol. 1, p. 86. Berlin.
 Bryan, A. H. and Greenberg, B. G. (1952). *Hum. Biol.*, 24, 117.
 Clatworthy, H. W., Jr. and Anderson, R. G. (1944). *Amer. J. Dis. Child.*, 67, 167.
 Clements, E. M. B. (1953). *Brit. med. J.*, 2, 897.
 — (1954). *Archives of Disease in Childhood*, 29, 147.
 Davis, C. M. (1928). *Amer. J. Dis. Child.*, 36, 651.
 Dean, R. F. A. (1951). *Brit. J. Nutr.*, 5, 269.
 — and Schwartz, R. (1954). *Courrier*, 4, 293.
 Dreizen, S., Snodgrass, R. M., Webb-Peploe, H., Parker, G. S. and Spies, T. D. (1957). *A.M.A. J. Dis. Child.*, 93, 122.
 Elgenmark, O. (1946). *Acta paediat. (Uppsala)*, 33, Suppl. 1.
 Ely, R. S. and Sutow, W. W. (1952). *Pediatrics*, 10, 115.
 Fitt, A. B. (1924). *Rep. Aust. Ass. Adv. Sci.*, 16, 704.
 Forbes, G. B. and Perley, A. (1951). *J. clin. Invest.*, 30, 558.
 Freund, J. and Maier, E. H. (1952). *Z. Kinderheilk.*, 71, 1, 79.
 Friis-Hansen, B. (1955). *Helv. paediat. Acta*, 10, 12.
 Greulich, W. W. and Pyle, S. I. (1950). *Radiographic Atlas of Skeletal Development of the Hand and Wrist*. Stanford University Press.
 Harding, V. S. V. (1952). *Child Developm.*, 23, 247.
 Hogben, H., Waterhouse, J. A. H. and Hogben, L. (1948). *Brit. J. soc. Med.*, 2, 29.
 Howe, P. E. and Schiller, M. (1952). *J. appl. Physiol.*, 5, 51.
 Kornfeld, W. (1957). *Hum. Biol.*, 29, 62.
 Larcomb, J. W., Perry, C. S. and Peterman, R. A. (1954). *J. Pediat.*, 45, 70.
 Macy, I. G. and Hunscher, H. A. (1951). *J. Nutr.*, 45, 189.
 Mainland, D. (1953). *Pediatrics*, 12, 114.
 — (1954). *Ibid.*, 13, 165.
 Maresh, M. M. (1955). *A.M.A. J. Dis. Child.*, 89, 725.
 Medawar, P. B. (1945). In *Essays on Growth and Form, presented to D'Arcy W. Thompson*, ed. Clark, W. E. Le Gros and Medawar, P. B., p. 157. Oxford.
 Meredith, H. V. (1941). *Amer. J. Dis. Child.*, 62, 909.
 — (1951a). *A.M.A. J. Dis. Child.*, 82, 702.
 — (1951b). *Growth*, 15, 39.
 — (1953). *Pediatrics*, 12, 411.
 — and Meredith, E. M. (1950). *Child Developm.*, 21, 141.
 Morant, G. M. (1951). *Brit. med. Bull.*, 7, 316.
 Mülhausen, W. (1953). *Msehr. Kinderheilk.*, 101, 13.
 Newcomer, E. O. and Meredith, H. V. (1951). *Hum. Biol.*, 23, 24.
 Osgood, E. E. (1955). *Pediatrics*, 15, 733.
 Palmer, C. E. (1933). *Publ. Hlth Rep. (Wash.)*, 48, 211.
 Prader, A. and Perabo, F. (1952). *Helv. paediat. Acta*, 7, 517.
 Reynolds, E. L. (1950). *Monogr. Soc. Res. Child. Dev.*, 15, No. 2. Child Development Publications.
 — and Sontag, L. W. (1945). *J. Pediat.*, 26, 336.
 Scammon, R. E. (1930). In *The Measurement of Man*, by Harris, J. A., Jackson, C. M., Paterson, D. G. and Scammon, R. E., p. 173. Univ. of Minnesota.
 Schlesinger, B. E., Butler, N. R. and Black, J. A. (1956). *Brit. med. J.*, 1, 127.
 Shuttleworth, F. K. (1951). *Monogr. Soc. Res. Child. Dev.*, 14, Nos. 1, 2. [1949.]
 Talbot, N. B., Sobel, E. H., McArthur, J. W. and Crawford, J. D. (1952). *Functional Endocrinology from Birth through Adolescence*. Harvard University Press.
 Tanner, J. M. (1952). *Archives of Disease in Childhood*, 27, 10.
 — (1955). *Growth at Adolescence*. Oxford.
 Thomson, J. (1956). *Brit. J. prev. soc. Med.*, 10, 128.
 Todd, T. W. (1937). *Atlas of Skeletal Maturation*. St. Louis.
 Weir, J. B. de V. (1952). *Brit. J. Nutr.*, 6, 19.
 Widdowson, E. M. (1951). *Lancet*, 1, 1316.
 — and Spray, C. M. (1951). *Archives of Disease in Childhood*, 26, 205.
 Wilkins, L. (1950). *The Diagnosis and Treatment of Endocrine Disorders in Childhood and Adolescence*. Springfield, Illinois.

PART II. PHYSIOLOGICAL CONSIDERATIONS

Physiological maturation represents in some degree the process of adaptation to the new requirements of a changing environment.

Perhaps the first task of the infant at birth is to insure for himself a proper supply of oxygen in the shift from placental to pulmonary respiration. By and large, this adjustment is one which he achieves by his own efforts. What we can do for him remains rather tentative in spite of the vagaries of fashion with respect to techniques of what we are still pleased to call resuscitation.

Great uncertainty still surrounds the concept of what actually represents a proper oxygen supply for a newborn infant. There is good reason to believe that the oxygen requirement of the newborn

infant is not by any means as immediately pressing as would be that of an older individual in comparable circumstances. Indeed, studies of postnatal oxygen saturation of arterial or capillary blood (Apgar, Girdany, McIntosh and Taylor, 1955) have disclosed a wide range of variation in the time required for individual infants to attain an arterial oxygen pressure believed to be necessary for transport of oxygen in adequate quantity to the tissues. Some of the newborn infants whose blood oxygen we followed at intervals over the course of the first three hours after delivery were still showing blood oxygen levels of less than 5 volumes per 100 ml. as late as 15 minutes after birth, and a few for even longer; yet in subsequent evaluation three to five

years later they exhibited no demonstrable residua of their earlier hypoxaemia. The battery of tests to which these subjects were exposed included a careful history to uncover any episodes which might be classed as seizures, a detailed neurological examination and formal psychometric evaluation by a modification of the Stanford-Binet test. On comparison with controls, the patients who in early infancy had been exposed to low blood oxygen pressures had progressed as satisfactorily as had their more briskly oxygenated fellows in the observational group. These somewhat reassuring results stand in sharp contrast to the prevalent tendency among neuropathologists to attribute a number of clinical syndromes, including cerebral lesions, to the action of postnatal anoxia (Wolf and Cowen, 1954), even when the historical evidence for anoxaemia is unimpressive. While the biochemical studies referred to have served to remove some of the blame from the long-suffering whipping boy, postnatal anoxaemia, too readily accused of responsibility for such sequelae as mental retardation, cerebral palsy, or epileptic seizures, the role of prenatal anoxia remains largely unexplored, unmeasured. Uncertainty in regard to the oxygen requirement of premature infants is even more formidable, for in this area of the special biochemistry of the immature subject we are still groping for fundamental information.

Tolerance of the foetal and newborn mammal for low oxygen pressures at the tissue level has long been established for experimental animals and seems to be related to an alternative capacity to derive energy from carbohydrate breakdown through a process of anaerobic glycolysis (Wilson, Reardon and Murayama, 1948). Evidence that the human foetus possesses this capacity for anaerobic glycolysis has only recently been obtained (Villev, 1957). How long such a capacity for biochemical legerdemain persists, how long after the infant's birth it constitutes an available resource, is not yet certain. While it lasts, it apparently purchases energy at the cost of accumulation of non-volatile metabolic breakdown products, especially lactic and pyruvic acids, which cannot be got rid of as rapidly as can carbon dioxide and water. One result is the temporary establishment in the newborn infant of a metabolic acidosis which is expressed in the plasma by depression of the pH and of the concentration of bicarbonate ions. For some time following birth many premature infants exhibit periodic breathing in room air but can be made to breathe regularly when placed in an oxygen-enriched atmosphere. Wilson and his colleagues (1948) interpreted this phenomenon as evidence that the premature infant

is apt to be incapable of meeting his full oxygen requirements by breathing ordinary room air. Nowadays, on the other hand, such evidence which might make one wish to enrich the oxygen content of the atmosphere to which premature infants are exposed is heavily counterbalanced by the risk of inducing retrolental fibroplasia (Crosse and Evans, 1952). An interesting observation on the developmental aspects of susceptibility to retrolental fibroplasia is the demonstration that, when once the retinal vessels have grown far enough to reach the ora serrata, they are no longer readily vulnerable to high oxygen injury (Ashton, Ward and Serpell, 1954).

The role of foetal haemoglobin in determining the transient peculiarities of oxygen utilization of the newborn infant has not yet been fully elucidated. The blood of premature infants has been shown by Jonxis (1949) and by Cook, Brodie and Allen (1957) to contain relatively large proportions of foetal haemoglobin, averaging close to 90% of total haemoglobin, while the blood of mature newborn infants contains somewhat less, averaging about 72%. Although foetal haemoglobin is known to take up oxygen in the alveoli more readily than does adult haemoglobin, at the same time it gives up oxygen less readily to the tissues. The latter characteristic has been offered as a partial explanation of the relative anoxia of premature infants. Whilst the concentration of carbonic anhydrase in the newborn infant's blood is only about half that of the adult, and in the premature infant as little as one-quarter (Stevenson, 1943), there is no proof that these reduced quantities of the enzyme constitute a physiological handicap (Day and Du Pan, 1949). In other words, shortage of carbonic anhydrase cannot be offered as an explanation of inadequate pulmonary alveolar exchange of CO₂ for O₂. Save for the accurate mapping of the disappearance of foetal haemoglobin from the infant's blood with time and growth (Jonxis, 1949), most of the biochemical and physiological changes just described which have to do with oxygen supplies have not been fully exploited as maturational phenomena.

The infant's ability to regulate his body temperature is dependent in part on his state of maturity. With falling body temperature the point at which shivering begins is lower in the premature infant than in one born at term; some very small infants in fact fail to show reflex shivering at all. With rising body temperature the premature infant is again at a disadvantage in that his deficient sweat production (Day, Curtis and Kelly, 1943) deprives him of the ability to lose heat by evaporation of

water from the skin surface. His relative poverty in subcutaneous fat adds to his difficulties in that, lacking its insulating properties, his tissues as a whole have a relatively high thermal conductance. The smaller the subject, the higher is the ratio of surface area to body weight, a potentially important aspect of heat gain or loss by convection and conduction. The practical significance of relatively large surface area, however, in the maintenance of steady temperature must in point of fact be of secondary order, for it is common to observe a gradual stabilization of a premature infant's body temperature, showing as a decrease in the excursion from highest to lowest point in a 24-hour cycle, long before gain in weight has accounted for more than a minute drop in the ratio of surface area to mass.

Observations documenting the development of the functions just mentioned are fragmentary in comparison with studies of renal regulatory capacity in early life, most of which have been carried out in the past decade. Only a brief and quite inadequate summary can be offered in this context. The ingenious investigations of McCance and Widdowson (1953) throw light on renal performance before birth as well as afterward. Urine collected from infants at the moment of birth, and therefore formed some time before, was compared with maternal urine and with urine excreted later by the infant. Prenatal urine was characterized by low osmolality, less than half that of the cord serum, to which NaCl contributed a relatively large fraction and urea a small fraction; it usually contained a little more Na than Cl, comparatively little K, and only traces of inorganic P. These findings provoke interesting speculation as to the source of excess water over and above foetal growth requirements, and one must wonder also whether the foetal kidney is capable of responding to the antidiuretic hormone of posterior pituitary or to parathyroid hormone.

In order to explore the transition from intra-uterine to extra-uterine life, McCance and Widdowson (1954) then studied the urine excreted in the first two days of life by infants who during this period were given no food and little water, in all, only about 25 ml. of water over the course of 48 hours. In these circumstances the substances presenting for renal excretion are almost exclusively those which the infant brings with him into the extra-uterine environment or produces in the course of his own energy release. Although the volume of urine passed was small, as would be expected with essentially no intake of fluid, its osmolar concentration rose only moderately from the relatively low level of solutes found in prenatally formed urine. Indeed, its average osmolar concentration during

the first two days after birth was only about one-third that which would be attained in the urine of adult subjects in comparable conditions of fasting and thirsting. Both urea clearance and glomerular filtration rates were at this stage low in comparison with values which one might expect of an older subject in relation to the volume of body water.

These and other findings may be looked on as manifestations of relative immaturity of renal function in the immediately postnatal period and in the first few weeks of extra-uterine life. Impressive ingenuity has been expended by a number of investigators (Rubin, Bruck and Rapoport, 1949; Vesterdal and Tudvad, 1949; McCance, 1950; Barnett, Vesterdal, McNamara and Lauson, 1952; McCance, Naylor and Widdowson, 1954) in defining with ever greater precision the quantitative aspects of this feature of physiological maturation. In view of the variety of methods used, substances measured and conditions of experiment, the coincidence of results and the agreement in interpretations appear little short of remarkable. Whether glomerular filtration rate is estimated by clearance of endogenous creatinine, mannitol, or inulin, all workers concede that immediately after birth the rate is relatively low, that it begins to rise almost at once, and that in the course of several days or weeks, possibly as long as a few months, it reaches values corresponding to those found in older subjects. The performance of the infant's kidney, if it is to be compared with that of an older, larger, and more mature person, must be brought into some common frame of reference such as weight, height or surface area. By general agreement all earlier workers evaluated their results after calculating equivalence with a subject benevolently endowed with the arbitrary standard surface area of 1.73 square metres. McCance and his colleagues (1954) introduced the device of referring urine flow and other renal functions to total body water, rather than to surface area. Although this calculation involves certain assumptions which may not be strictly valid for the individual infant, it does strive to relate renal performance to a bodily dimension more closely identified with what one expects of the kidney than is surface area. We do ask the kidney to maintain homeostasis with respect to water and electrolytes and other dissolved substances occupying the fluid compartments of the internal environment, and body water seems to be the kidney's direct business. Application of these calculations has the effect of bringing the infant's glomerular filtration rate close to the adult level at a much earlier age, within a month after birth, for instance, rather than in four months to a year as some earlier

estimates suggested. The use of calculated total body water as a frame of reference for evaluating renal functional maturation is not necessarily justified because it appears to accelerate adjustment to the circumstances of extra-uterine existence. Indeed, the possibility of an even more satisfactory criterion being exposed by closer study of the problem cannot be excluded from our speculations.

Another feature of renal function, clearance of those particular substances like para-aminohippurate and diodrast ('diodone') which are dealt with primarily in the tubules, is likewise found to be relatively low at birth, increasing with age but at a slower rate than the spontaneous rise in glomerular filtration rate previously mentioned. One practical result of relative immaturity of the young infant's kidney is his sluggishness in excreting certain substances used in treatment of infections, like penicillin, which leave the body chiefly by tubular excretion. It thus is easier with small infants to sustain a given plasma concentration of penicillin with doses usually used in therapy.

The ability of young infants to excrete a large dose of water administered by stomach tube has been studied by McCance *et al.* (1954). Again it becomes difficult to know precisely how to put small infants on the same footing as the adults with whom they are to be compared; but when subjects of any age were given over a brief interval a dose of water amounting to 5% of their body weight, its excretion was appreciably less efficient on the part of infants younger than 3 weeks of age. Whereas an adult could regularly excrete 90% of the administered dose within four hours, infants managed in the same period of time to get rid of an average of only 57% of the test load. In addition, the infant was incapable of excreting urine of quite as low osmolar concentration as could the adult at the peak of diuresis. It should be added that other workers, whose results in general were in agreement with those of McCance and his group just quoted, differed in regard to the last point in that they succeeded in causing young infants, even premature infants weighing less than 2,500 g., to excrete urine fully as dilute as that produced by more mature subjects. Barnett and his colleagues (1952), for example, by imposing a water load of 40 ml. per kg. of body weight in premature infants reduced the urine concentration of their subjects to the level of 40 to 60 m.osmol./l. at the moment of maximal diuresis, which is the same as saying that they were unable to bring out evidence of renal functional immaturity even in their very small patients, so far as the ability to eliminate superfluous water was concerned. They made the point,

however, that the ability to excrete urine of low osmolar concentration is a function of postnatal age, rather than of body weight. Thus, their patients aged 5 to 8 days showed a limited capacity to excrete dilute urine, while another subject, a premature infant 36 days of age whose body weight had reached only 2,460 g. at the time of the test, reacted to the water load in a manner comparable to that of an adult.

In this connexion the investigations of Ames (1953) should be mentioned. Although her primary goal was to explore the function of the posterior hypophysis in the first days after birth and to learn whether the newborn infant's kidney would respond to antidiuretic hormone, her studies brought out interesting aspects of the process of maturation. Water administration comparable to that used in the experiments referred to a moment ago disclosed that, in general, infants less than 3 days of age exhibit little, if any, diuresis in response to a water load. This situation changes rapidly, however, with increasing postnatal age and by one month after birth all infants in her study were capable of excreting 85% or more of the administered load within the course of two hours. One very small infant of birth weight 1,200 g. excreted only 50% of the administered water load within two hours at the age of 4 days, but by 12 days of age accounted for 100%. Since no water diuresis could be demonstrated in infants younger than 3 days of age, no response to injected antidiuretic hormone was expected or found; but in older infants, that is, older than 3 days of age, who were showing a brisk diuresis in response to administered water, injection of as little as 0.5 milli-unit of pitressin intravenously invariably checked diuresis promptly. When infants less than 3 days of age were subjected to a period of thirsting of six or eight hours' duration and their urine collected and analysed for anti-diuretic hormone, relatively large amounts of the hormone were demonstrable, the average concentration being 1.25 ± 0.25 milli-units per ml. This proves that the posterior hypophysis is indeed capable of functioning at this age and of responding to the stimulus of mild thirst.

Liver function is difficult to measure at any age, and few investigators have been bold enough to attempt to observe maturation of any single component of this complicated field. Mollison and Cutbush (1949) took advantage of the accessibility of the umbilical vein in infants less than six hours old to inject bromsulphalein and measure the speed of its disappearance from circulation. In general, the rate of clearance was very much slower in these infants than it was in adults to whom an analogous

dose of 5 mg./kg. body weight had been given. The difference was ascribed to immaturity of hepatic excretory function. Their patients were followed long enough to observe the severity of physiological jaundice appearing in the course of the next few days, and no gross difference was found, in regard to bromsulphalein clearance, between those who later became jaundiced and those with lower degrees of bilirubinaemia. A somewhat similar study was carried out by Yudkin, Gellis and Lappen (1949) in which, again, bromsulphalein was administered intravenously to infants at various ages up to 9 days. Excretion was delayed in all. Although the degree of delay diminished with increasing age of the subjects, the rate of excretion was still sluggish in a 9-day-old veteran, the oldest one included, in comparison with adult standards. It would seem obviously desirable to pursue such studies long enough to learn when the mature level of bromsulphalein excretion is reached.

Excretion of bilirubin by young infants was studied by Fashena (1948) in relation to the occurrence of physiological jaundice of the newborn. Functional immaturity of the liver had long ago been postulated on theoretical considerations as one of the plausible determinants of neonatal jaundice, since no satisfactory correlation could be found between the degree of bilirubinaemia and the postnatal fall of haemoglobin concentration, red blood cell count or haematocrit level in the peripheral blood. Determination of the velocity of excretion of bilirubin (Weech, Vann and Grillo, 1941) as a liver function test has the double advantage that it employs in the load test a natural substance, rather than a totally extraneous one like bromsulphalein, and that it can be carried out without risk in the presence of hyperbilirubinaemia and clinical jaundice. All infants included in the study were tested during the first week of life. Those whose blood before loading showed a serum bilirubin level higher than 2.4 mg./100 ml. invariably displayed sluggish excretion of bilirubin, with a velocity constant lower than 2.0×10^{-3} mg. units. (The smaller the constant, the more limited is excretory function.) Those whose serum bilirubin level was lower than 1 mg./100 ml. before application of the test had a velocity constant equal to or greater than 2.8×10^{-3} , reflecting satisfactory excretory function. In four infants who exhibited impaired excretion during the course of hyperbilirubinaemia and physiological jaundice, a second test repeated after the lapse of four to eight days showed that the velocity constant had now risen to normal level. The inference seems clear that many, not all, infants have immature hepatic excretory function for

bilirubin during the first few days following birth, and that in the course of time mature function can be expected to develop spontaneously.

Many liver function tests which can be employed in older subjects in the study of disease have yielded little more than confusion when applied to young infants in an effort to measure the maturational process, so that useful information extends but little farther than the brief summary of work just cited. A new area has recently been opened up, however, which bids fair to add greatly to our knowledge of physiological development. I refer to the maturation of hepatic cellular enzyme systems. An early lead was given by the observation of Levine, Gordon and Marples (1941) that premature infants fed a high-protein diet without ascorbic acid excreted large amounts of hydroxyphenyl compounds in their urine, that mature infants would excrete these substances only if fed rather large amounts of the precursor amino-acids D, L-phenylalanine or L-tyrosine, and that in either premature or mature infants who showed the trait the excretion of these unusual hydroxyphenyl compounds could be promptly abolished by administration of large amounts of ascorbic acid. The phenomenon when it was first studied was recognized as having attributes of a maturational episode, but little attention was for a time attached to the possibility of its being localized in hepatic cells. Quite recently Kretchmer, Levine, McNamara and Barnett (1956) have devised techniques for studying soluble enzyme systems extracted from hepatic cells obtained from human livers, and have been able to measure the progressive development of specific enzyme constellations such as the tyrosine oxidizing system. Premature infants, full-term infants and adults formed three contrasting groups, with enzyme activity increasing in proportion to the figures 1 for prematures, 5 for mature infants and 20 for adults, whether comparison was made on the basis of wet weight of liver tissue originally used as source, milligrams of nitrogen in the enzyme solution, desoxyribonucleic acid content of the homogenate of liver cells or surface area of the subject. These initial studies explored only the tyrosine oxidizing system. In a subsequent extension of the work Kretchmer, Levine and McNamara (1957) demonstrated a relative lack of p-hydroxyphenylpyruvate oxidase activity in the livers of small infants, which could be partly overcome *in vitro* by the addition of relatively large quantities of ascorbic acid. The response of this particular enzyme system to ascorbic acid seemed to be dependent upon the weight of the infant, again suggesting a maturational phenomenon.

The fact that immaturity of these highly specific

hepatic cell enzyme systems results in urinary excretion of substances not ordinarily found there affords certain promise of an impending flood of new discoveries in the general picture of maturation of liver function. The element of chance ordinarily inherent in divination of the future is entirely removed from this particular prophecy by recent demonstration of the pattern of urinary amino-acid excretion in foetuses, premature infants, and mature infants of various ages. Norton, Pratt and Hassel-meyer (1954) have confined their study almost entirely to premature infants, while Woolf and Norman (1957) extended theirs to include a variety of ages and stages of maturity. The pattern is a complicated one but at the same time strongly consistent in the appearance and disappearance of individual components. This consistency applies not only to the excretion of amino-acids but to that of other substances as well—creatine, taurine, and various sugars, for example. Doubtless not all of them are controlled by enzyme systems residing exclusively in the liver, but it seems a reasonable prediction that most of them are and that closer study of liver function will eventually clarify the developmental picture. The fact that the waxing and waning pattern of urinary amino-acids represents a feature of developmental maturation is borne out by their spontaneous disappearance in the course of a few months.

The prothrombin level of the young infant's plasma follows a course which suggests a developmental evolution influenced in part by liver function, in part by the ingestion of prothrombin precursors in the food, and in considerable part also by the formation of precursors within the alimentary tract as a by-product of growth and metabolism of contaminating bacteria. The vagaries of the prothrombin story in early life depend a great deal on methods used in analysis, and complete agreement has not yet been reached among all authorities. Nevertheless, the general outlines are clear and I shall take the liberty of presenting them more or less in Quick, Murat, Hussey and Burgess's (1952) terms. When the mother has received no vitamin K during pregnancy beyond that contained in the diet, the prothrombin as measured by the one-stage method is found to be at about the same level at term in the mother's blood, the cord blood, and the infant's blood on the day of birth. On application of the two-stage method, which reveals not only active and available prothrombin but also reserves of prothrombinogen in inactive form, the mother's blood is found to have a considerable stock of prothrombinogen, a reserve which can easily be augmented still further by giving her additional

vitamin K. In the newborn infant, however, administration of vitamin K on the day of birth has no immediate effect on his prothrombin level, just as administration to the mother before delivery has no influence on the active prothrombin content of the infant. He appears incapable of manufacturing prothrombinogen within the first two to four days after birth, and his active prothrombin concentration may even fall to a lower level, reaching a nadir generally on the third day of life. Thereafter it begins to rise and by the end of the first week or a little longer has once more reached the level which was present at birth. Reserves of prothrombinogen build up more gradually in the infant's blood, and as much as a year may elapse before the level has reached that which an adult usually shows. The pace of accumulation of reserves of inactive prothrombin may be retarded by liver trouble, by obstructive lesions of the upper alimentary tract which interfere with bacterial colonization of the lower tract and by persistent administration of antibiotics which likewise inhibit the growth of a normal intestinal flora.

Mention of infection, in this instance a beneficent one, leads naturally to discussion of developmental immunology, the acquisition of defence against harmful infections. The subject may be divided into three sections: mechanisms of defence transferred from mother to child; mechanisms produced by the infant independently of contact with specific infecting agents; and, finally, development of resistance orientated specifically toward a particular agent of disease with which the infant or child has come in contact in some form—dead (vaccine) or alive (infection).

A good deal of accurate quantitative information is available concerning transplacental conveyance of humoral immunity. It has long been known that diphtheria antitoxin concentrations in the blood of a mother and in the cord blood of her infant at term are essentially equal. The premature infant is not always quite so well off; and if he is very young and immature, his mother's blood will usually show a considerable excess of antitoxin. Osborn, Dancis and Rosenberg (1952) set the crucial maturity point at a birth weight of roughly 2 kg., in that infants smaller than this limit were more apt than not to have an antitoxin titre lower than that of their mothers. After delivery, the infant receives no significant increment of diphtheria antitoxin from his mother, regardless of whether he nurses her colostrum; and from birth on, there begins a logarithmic descent of the antitoxin level which leaves him devoid of adequate protection from this source from the age of about 4 to 6 months onward, if no additional steps are taken.

Placental transfer of other specific defence agents has been similarly documented. A well known example is the postnatal curve of blood concentration of bactericidal antibodies against *Haemophilus influenzae*, plotted against age, which was long ago published by Fothergill and Wright (1933). The newborn infant has a significant antibody titre, which then falls rapidly so that by the age of 3 to 6 months specific antibody can no longer be measured. Not until the age of 3 years does the curve begin to rise again, and the authors brought out clearly in their publication the fact that the age incidence of *H. influenzae* meningitis reaches its peak during the period between 3 months and 3 years of age when specific antibody against this organism is most difficult to detect.

The age incidence of measles clearly suggests that immunity is passively transferred from the mother before the infant is born, and that the substance responsible for immunity gradually diminishes by attrition, becoming unreliable as an effective defence from about the age of 6 months onward. This interpretation was widely accepted on epidemiological evidence long before titration of measles antibody became technically feasible. Not all specific infections behave alike, however. Chicken-pox and whooping cough frequently occur within a few weeks after birth, while mumps in infants less than 1 year of age is distinctly rare. We still lack accurate information regarding the defences against a number of infections, and in many instances we are quite uncertain how much resistance depends on humoral as opposed to non-humoral agencies.

In our search for information we must of necessity resort to certain indirect methods of investigation, less closely orientated toward specific infections. While it seems unlikely that the gamma globulin fraction of plasma is composed entirely of specific antibodies, a degree of parallelism undeniably exists between serum gamma globulin concentration and resistance to a variety of infections. Studies of the gamma globulin fraction of plasma in mother and foetus indicate that from an early stage of pregnancy, perhaps even as early as the beginning of the second trimester, the concentrations in the two bloods are virtually identical. At term the concentrations have been shown to be essentially the same in cord blood and in maternal blood (Orlandini, Sass-Kortsak and Ebbs, 1955). From this point on, the infant's plasma gamma globulin level commences a gradual fall which reflects the general rate of attrition of plasma protein components which are not being renewed by active production. Gitlin (1957) gave the half-life of a preparation of gamma globulin as 20 days when

radioactivity was the basis of measurement, 45 days when immunochemical methods were used. It seems reasonable to expect that, because of known variations in the physical and chemical behaviour of individual plasma proteins, different rates of spontaneous attrition of individual specific antibodies will be found. In general, what has been learned in the laboratory concerning the disappearance of gamma globulin from an infant's blood in the early weeks of his independent existence following birth is entirely consistent with what is known of his gradual loss of resistance against certain infections.

Even the newborn infant possesses some humoral defence against infection which is independent of specific antibodies. The all-purpose defence on which most attention has recently been focused is the properdin system, initially described by Pillemer, Blum, Lepow, Ross, Todd and Wardlaw (1954). According to Hinz (1956) properdin, itself a protein of high molecular weight, is demonstrable in plasma in approximately equal concentration at all ages. The properdin system, to be effective in controlling infection, must include interaction of properdin, complement and Mg^{++} ion. Although the exact role of the properdin system in early life has not yet been fully assessed, some explanation is required for the asymptomatic course which most infants succeed in following between the time when the antibodies they have received from their mother give out and the time when their own antibody production has reached an adequate level. It would seem highly probable that mechanisms of defence are operating on the young infant's behalf, of which we have as yet only imperfect knowledge, such as the circumstances which govern phagocytosis, or about which we know nothing at all. We lack a satisfactory explanation of the paradox that young infants from an early age commonly harbour virulent strains of staphylococci in the nasopharynx, and yet the frequency of significant staphylococcal infections remains on the whole surprisingly low.

Chemical analyses of plasma for its total content of gamma globulin, whether the analytic method is that of electrophoresis or the cruder but nevertheless serviceable titration of zinc turbidity (Harris, 1952; Orlandini *et al.*, 1955), show that the concentration reaches its low point at the age of about 3 months, after which it starts a gradual rise. The slope of the ascending curve of gamma globulin concentration has not been shown to be clearly dependent on the number of recognized infections experienced by the individual subject. Raffen (1953) cites anatomical evidence that bone marrow of infants does not contain plasma cells until the age

of 3 months. This finding is consistent with the commonly accepted belief that the plasma cell is the site of production of specific antibodies, and with the timing of the point on the age axis at which plasma gamma globulin begins to be renewed from autochthonous sources. It leaves unexplained, however, the demonstrated ability of an infant much younger than 3 months of age to respond to contact with an appropriate antigen by production of antibody in significant quantities, as will be described later.

Studies of specific antibody formation in response to spontaneously acquired infections, whether clinical or subclinical, are surprisingly few. Fothergill and Wright's (1933) measurements of the curve of antibody titre against *H. influenzae* in relation to age have already been mentioned, with their rise in titre beginning at age 3 years. To judge from the information gathered on this subject regarding antibodies to the three chief types of poliomyelitis virus, before the widespread use of vaccine, the likelihood that any given individual would show significant levels of one, two, or all three antibodies depended on a host of variables—his age, the number of people in the family, rural or urban setting, school contacts, housing standards and standards of general hygiene, as well as exposure to epidemic infection (Melnick, Paul and Walton, 1955; Sabin, 1955). Important background information of this kind in relation to poliomyelitis will obviously become entirely unattainable as soon as prophylactic administration of vaccine becomes universal. The chances of obtaining comparable information concerning humoral antibodies to various bacterial infections seem likewise to be evaporating in the face of the widespread application of interfering factors in the form of effective antibacterial agents.

A vast field remains open for future study, of course, in the measurement of the infant's or child's response to vaccines and other agents given for prophylactic purposes. Results published thus far would seem to indicate that when a three-pronged agent like the popular combination of tetanus and diphtheria toxoids and pertussis bacterial vaccine is injected into young infants, beginning at 1 week of age, an apparently different response is obtained to each of the three components (di Sant'Agnese, 1949). Successful production of antibodies against tetanus toxin indicates that even at this early age active immunization is feasible. On the other hand, the less impressive development of measurable antibodies to diphtheria toxin and to pertussis leaves unanswered whether the fault lies with the infant, in an immature mechanism for responding to a particular agent, or in the nature of the antigen used,

or possibly in the interfering effect of residual specific antibody originally transmitted *in utero*. At a later age, antibody production in response to the same agent is much more active.

Certain infections evoke different responses at different ages, usually producing more serious disease the younger the subject. A number of examples come to mind. Congenital toxoplasmosis, transmitted from mother to infant *in utero*, is almost invariably more serious than the disease acquired in later life. This difference may be one of dosage, but thus far proof is lacking. Congenital syphilis in the pre-penicillin era was far more lethal to the foetus and infant than to the mother. Among severe infections caused by *H. influenzae*, purulent meningitis is most frequent between the ages of 2 months and 2 years, almost 80% of all examples of influenzal meningitis falling in this sharply limited age zone. Acute epiglottitis or infection of the supraglottic structures caused by the same organism is even more frequently associated with bacteraemia, can also be fatal all too frequently, and yet is hardly ever encountered in a patient younger than 2 years of age. Infection with Coxsackie B virus, according to Gard (1954) and Johnsson (1954), has produced three distinct syndromes, depending on the age of the patient: a fatal myocarditis in very young infants, a form of 'aseptic meningitis' which leaves no permanent residua in older infants and children, and in adults pleurodynia or so-called epidemic myalgia. The Sendai virus, first isolated by Kuroya, Ishida and Shiratori (1953) in Japan in 1951, was originally recovered from the lungs in fatal cases of pneumonitis which occurred in newborn infants. Later, an apparently identical virus was grown from the respiratory tract of adults suffering from a non-fatal form of an influenza-like syndrome, and Matsuzawa and his colleagues (1953) still later isolated the same virus from a child who recovered after a 40-day illness with respiratory symptoms. In the last instance a rise in antibody titre was demonstrated for the first time in the convalescent patient. Jensen, Minuse and Ackermann (1955), to whom a sample of the virus was sent soon after its initial isolation, found significant levels of antibody in a large number of sera collected from various parts of the State of Michigan. Some of these American sera were fresh, others had been preserved for as long as 19 years, and in no instance was historical evidence obtainable of a corresponding clinical infection. One is left to speculate whether the diverse patterns of clinical disease produced at different ages by what are believed to be identical infecting agents depend upon insidiously acquired specific immunity, which must of necessity have been

inadequate to achieve full protection, or whether they reflect different levels of immunological maturation.

In bringing this fragmentary and tentative consideration of growth and development to a close, it is conceivably more prudent as well as more informative to take note of known omissions than to attempt to survey the ground covered. Nothing has been said of intellectual growth, of emotional maturation, or social adaptation. Important as these are, all of them, in the assessment of a clinical situation, they represent areas of concern in which the physician's responsibility is subordinate to that of the family unit, the school, the community. The broad field of symptomatic expression with relation to age of patient has been almost completely ignored, and that companion territory—the age incidence of specific diseases and syndromes, which goes so far to justify paediatrics as a specialty within the broad boundaries of general practice—has received but scant mention. Only a beginning has been made in indicating lines of investigation which seem to hold promise for the near future. There is no need to dwell on the diminutive scope of what we now know in comparison with what is yet to be learned. Growth and development, with their vagaries and their ever-changing manifestations, keep a physician on the alert and add flavour of their own to the pursuit of the Hippocratic tradition among those patients who have not ceased to be children.

I would express my deep obligation to my paediatric colleagues on the staff of Babies Hospital in New York for the help and inspiration they have given over the past several months in the shaping of these thoughts. The task of setting them down in organized form has been a happy one, constantly lightened by awareness of their intent, to pay tribute to the memory of the man whom these lectures honour, Sir Leonard Parsons.

REFERENCES

- Ames, R. G. (1953). *Pediatrics*, 12, 272.
 Appgar, V., Girdany, B. R., McIntosh, R. and Taylor, H. C. (1955). *Ibid.*, 15, 653.
 Ashton, N., Ward, B. and Serpell, G. (1954). *Brit. J. Ophthalm.*, 38, 397.
 Barnett, H. L., Vesterdal, J., McNamara, H. and Lauson, H. D. (1952). *J. clin. Invest.*, 31, 1069.
 Cook, C. D., Brodie, H. R. and Allen, D. W. (1957). *A.M.A. J. Dis. Child.*, 93, 9.
 Crosse, V. M. and Evans, P. J. (1952). *A.M.A. Arch. Ophthalm.*, 48, 83.
 Day, R. and Du Pan, R. M. (1949). *Amer. J. Dis. Child.*, 77, 109.
 —, Curtis, J. and Kelly, M. (1943). *Ibid.*, 65, 376.
 di Sant'Agnese, P. A. (1949). *Pediatrics*, 3, 20.
 Fashena, G. L. (1948). *Amer. J. Dis. Child.*, 76, 196.
 Fothergill, L. D. and Wright, J. (1933). *J. Immunol.*, 24, 273.
 Gard, S. (1954). *Acta paediat. (Uppsala)*, 43, suppl. 100, p. 54.
 Gitlin, D. (1957). *Pediatrics*, 19, 657.
 Harris, R. C. (1952). *Bull. N.Y. Acad. Med.*, 28, 721.
 Hinz, C. F. (1956). *Ann. N.Y. Acad. Sci.*, 66, 268.
 Jensen, K. E., Minuse, E. and Ackermann, W. W. (1955). *J. Immunol.*, 75, 71.
 Johnson, T. (1954). *Arch. ges. Virusforsch.*, 5, 384.
 Jonxis, J. H. P. (1949). In *Haemoglobin* [Barcroft Memorial Conference], p. 261. F. J. W. Roughton and K. C. Kendrew, editors. London.
 Kretschmer, N., Levine, S. Z. and McNamara, H. (1957). *A.M.A. J. Dis. Child.*, 93, 19.
 —, —, — and Barnett, H. L. (1956). *J. clin. Invest.*, 35, 236.
 Kuroya, M., Ishida, N. and Shiratori, T. (1953). *Tohoku J. exp. Med.*, 58, 62.
 Levine, S. Z., Gordon, H. H. and Marples, E. (1941). *J. clin. Invest.*, 20, 209.
 Matsuzawa, M., et al. (1953). *Virus (Osaka)*, 3, 336.
 McCance, R. A. (1950). *Amer. J. Med.*, 9, 229.
 — and Widdowson, E. M. (1953). *Proc. roy. Soc. B*, 147, 488.
 —, — (1954). *Archives of Disease in Childhood*, 29, 488.
 —, Naylor, N. J. B. and Widdowson, E. B. (1954). *Archives of Disease in Childhood*, 29, 104.
 Melnick, J. L., Paul, J. R. and Walton, M. (1955). *Amer. J. Publ. Hlth.*, 45, 429.
 Mollison, P. L. and Cutbush, M. (1949). *Archives of Disease in Childhood*, 24, 7.
 Norton, P. M., Pratt, E. L. and Hasselmeier, E. (1954). *A.M.A. J. Dis. Child.*, 88, 659.
 Orlandini, T., Sass-Kortsak, A. and Ebbs, J. H. (1955). *Pediatrics*, 16, 575.
 Osborn, J. J., Dancis, J. and Rosenberg, B. V. (1952). *Ibid.*, 10, 450.
 Pillemer, L., Blum, L., Lepow, I. H., Ross, O. A., Todd, E. W. and Wardlaw, A. C. (1954). *Science*, 120, 279.
 Quick, A. J., Murat, L. G., Hussey, C. V. and Burgess, G. F. (1952). *Surg. Gynec. Obstet.*, 95, 671.
 Raffel, S. (1953). *Immunity, Hypersensitivity, Serology*, p. 72. New York.
 Rubin, M. I., Bruck, E. and Rapoport, M. (1949). *J. clin. Invest.*, 28, 1144.
 Sabin, A. B. (1955). *Wld Hlth Org.: Monogr. Ser. No. 26*, p. 297. Geneva.
 Stevenson, S. S. (1943). *J. clin. Invest.*, 22, 403.
 Vesterdal, J. and Tudvad, F. (1949). *Acta paediat. (Uppsala)*, 37, 429.
 Villee, C. C. A. (1957). *Josiah Macy, Jr., Foundation Conference on Prematurity*, Princeton.
 Weech, A. A., Vann, D. and Grillo, R. A. (1941). *J. clin. Invest.*, 20, 323.
 Wilson, J. L., Reardon, H. S. and Murayama, M. (1948). *Pediatrics*, 1, 581.
 Wolf, A. and Cowen, D. (1954). In *Neurology and Psychiatry in Childhood*, R. McIntosh and C. C. Hare, editors. *Res. Publ. Ass. nerv. ment. Dis.*, vol. 34, p. 199. Baltimore.
 Woolf, L. I. and Norman, A. P. (1957). *J. Pediat.*, 50, 271.
 Yudkin, S., Gellis, S. S. and Lappen, F. (1949). *Archives of Disease in Childhood*, 24, 12.