Serum Siderophilin in Kwashiorkor

A. U. ANTIA, H. McFARLANE, and J. F. SOOTHILL

From the Departments of Paediatrics and Chemical Pathology, University College Hospital, Ibadan, Nigeria; and the Department of Immunology, Institute of Child Health, Guilford Street, London, W.C.1

Kwashiorkor is known to be associated with low plasma proteins (Edozien, 1960) and low serum iron-binding capacity (Edozien and Udeozo, 1960). Unpublished observations of Neale and Soothill (1958) confirmed that the iron-binding capacity was down to about one-third of a normal level, but showed that the siderophilin (syn. transferrin) was much more profoundly depressed—down to 6 mg./100 ml., compared with a normal European level of about 200. We therefore report a study of the serum siderophilin concentration estimated immunochemically in a series of children with typical kwashiorkor and of some control data. These values are correlated with a clinical grading of the severity of the disease and the effect of treatment.

Materials and Methods

The patients were 24 random attenders at the Nutrition Clinic of the Department of Paediatrics at the University College Hospital, Ibadan in whom the diagnosis of kwashiorkor was made on standard clinical grounds. These were classified clinically into three groups—mild, moderate, and severe. Criteria on which the classification was based were oedema, mental status, skin changes, and state of mobility. Serum siderophilin was estimated by the double gel diffusion precipitin technique (Soothill, 1962). Serum pooled from Nigerian healthy adults was used as standard. Results are presented in terms of this standard (% PS), which was estimated in terms of a purified siderophilin preparation kindly supplied by Dr. Tristram Freeman and found to contain 150 mg./100 ml. of siderophilin. It was also compared with the reference normal serum standard used by Soothill (1962) for which some data from a healthy European population were reported. The Nigerian standard was 90% of the European. Total serum proteins were estimated by the technique of Gornall, Bardawill, and David (1949).

Comparable data were obtained from four breast-fed infants who had marasmus due to breast milk insufficiency, and from 10 children with other diseases—infective, developmental, and neoplastic—not obviously associated with malnutrition.

Results

The results for the kwashiorkor series are shown in Table I, and for the marasmic and the control series in Table II. The mean values for siderophilin, total protein, and packed cell volume (PCV) for the three series are given. The siderophilin concentrations of the kwashiorkor series are significantly lower than either the marasmus series or the miscellaneous control series ($t = 2.26$, $p < 0.05$, and $t = 6.09$, $p < 0.001$, respectively on log data). The values for the control and the marasmic series and those from patients with kwashiorkor, classified both according to the clinical severity and to the course and management adopted (i.e. those who died, and those who received in-patient or out-patient treatment), are shown in Fig. 1. All the fatal cases were receiving in-patient treatment.

A 3×3 χ^2 analysis shows a highly significant relation between the levels of the siderophilin grouped into 3 equal batches: less than 13, 13 to 39, and more than 39, and the clinical course of the patients ($\chi^2 = 14$, $p < 0.01$), but not with the clinical assessment of the severity of the disease ($\chi^2 = 3.7$, $p > 0.4$), though the severe group includes the lowest values detected and the only case classified as mild was the highest. Serum total protein, and PCV fitted little better, even with the outcome classification (Fig. 2). Repeated values on some of the patients, treated either as in-patient or as out-patient by dietary measures alone, are given in Fig. 3.

Discussion

The low levels of siderophilin in kwashiorkor observed by Neale and Soothill (1958) are confirmed by these results. This may well be the most profound defect known of acquired disturbance of production of a plasma protein. Though there is a significant correlation between serum total protein and the siderophilin levels ($r = 0.5115$, $p < 0.02$), the proportional depression of the siderophilin concentration is considerably greater than that of

Received January 10, 1968.

TABLE I
Serum Siderophilin in Kwashiorkor

Age (yr.)	Sex	Duration of Symptoms (wk.)	Clinical Features Laboratory Data					
			Weight (kg.)	Clinical Grading	Clinical Outcome	PCV (%)	Total Protein (g./100 ml.)	Siderophilin % of PS
5	F		9.5	Severe	Died		3.5	4
1½	M	6	10.9	Severe	Died	17	4.3	6
1½	M	22	11.9	Moderate	Out-patient	24	3.4	7
1½	F	1		Moderate	Out-patient	22	4.3	8
2	M	2	8.6	Severe	Died	33		12
2	M	4	7.7	Severe	Died		4.4	12.5
3½	F	8	9.1	Severe	Died	28	3.8	12.5
5	F	10	9.5	Severe	Died	24	4.7	12.5
1½	F	6	5.9	Severe	In-patient	27	4.9	17
1½	F	3	6.4	Severe	Out-patient	27	4.9	25
5	F	4	9.1	Severe	In-patient	28	3.4	27
2½	F	12	10	Moderate	Out-patient	31	5.7	28
3	F	8	7.3	Moderate	In-patient	20	3.9	29
2	M	3	6.8	Severe	Out-patient	33	4.4	30
2½	F	8	6.8	Moderate	In-patient		4.2	33
2	M	2	14.1	Moderate	Out-patient			34
1½	M	4	7.7	Moderate	Out-patient	34	5.1	40
2	F	3	8.6	Moderate	Out-patient	32	4.2	50
1½	F	2	5.9	Moderate	Out-patient	31	5.4	50
5	F	9	8.2	Severe	In-patient		4	56
2½	M	3	8.2				4.2	59
5	F	8	17.7	Moderate	Out-patient	25	4.2	67
5	F	3	6.8	Severe	Out-patient	29	5.9	71
2	M	4	7.3	Mild	Out-patient	24	6.5	125
Kwashiorkor mean			8.9			27	4.5	34

the total plasma protein, which is never below half of the lowest value of the control series, though the siderophilin value fell to one-fifth of the lowest control value. The considerably higher values in

the severely marasmic children, in the control series containing a number of diagnoses, and in healthy newborns and adult Nigerians (McFarlane and Udeozo, 1968) and our experience in a wide

TABLE II
Serum Siderophilin in Nigerian Children with Miscellaneous Diseases and with Marasmus

Diagnosis	Age (yr.)	Sex	Weight (kg.)	Siderophilin % of PS	Total Protein (g./100 ml.)
Tonsillitis	2½	F	11.6	67	6.3
Tuberculosis	5	F	14.3	500	6.4
Bronchiolitis	½		6.9	200	6.9
Tuberculosis	7	F	14.1	286	7.1
Empyema	1½	F	10.9	57	6.6
Tuberculosis	3	M	14.1	200	7.2
Tuberculosis	2	F	10	200	6.2
Ventricular septal defect	1½	F	9.7	100	6.3
Osteomyelitis	2	M	10.9	125	—
Burkitt's tumour	—	—	—	200	—
Miscellaneous	mean		11.4	194	6.6
Marasmus	5/12	F	3.6	33	7.0
Marasmus	10/12	F	4.7	100	7.6
Marasmus	9/12	M	3.5	67	7.6
Marasmus	9/12	F	3.5	84	7.8
Marasmus	mean		3.8	71	7.5

FIG. 1.—Serum siderophilin levels in 24 patients with kwashiorkor, classified clinically into severe, moderate, and mild, and into those who died and those who were treated as in-patients or out-patients, in 4 children with marasmus, and in 10 children with miscellaneous diseases.

FIG. 2.—Serum total protein and packed cell volume (PCV) in patients with kwashiorkor, classified according to whether they died or whether they were treated as in-patients or out-patients.

range of disease processes in Great Britain (Neale and Soothill, 1958), suggests that the very low levels of siderophilin found in kwashiorkor may well be a diagnostic abnormality of this disease with the possible exception of asiderophilinaemia. The defect is almost certainly one of production, and it would be interesting to know whether it would affect the genetically controlled various siderophilin types differently. It is probable that these differences have little effect on the immunochemical assay. It is interesting that the grading, derived from outcome, and from the accustomed decision making entailed in clinical management, namely whether the patients died or were treated as in-patients or out-patients, fitted the siderophilin values far more closely than

FIG. 3.—Sequential values of serum siderophilin in 8 patients with kwashiorkor treated with diet; ● = out-patients and ○ = in-patients.

the attempt at a clinical grading of the severity of the condition when first seen. Clearly the test provides an objective assessment of the severity of the disease, of use for both treatment and research purposes. The remarkable response of the siderophilin level to a high protein diet (see Fig. 2—in one patient it rose nearly tenfold in a week) suggests that it would be useful for control of the efficacy of different treatment regimens. Such immunochemical determinations are easy and cheap.

If the observation is confirmed that the serum iron-binding capacity is not so profoundly depressed as the siderophilin (Neale and Soothill, 1958), it may be that this defect is less relevant in handling iron than might be expected. Indeed, it seems likely that the serum siderophilin has a considerable

functional reserve. There is little correlation between the siderophilin level and the packed cell volume in our series ($r = 0.508$, $p > 0.05$), suggesting that other factors are probably more relevant in the anaemia. More detailed knowledge of the amino acid requirement for production of each individual protein is required before one can speculate on the mechanism of this extreme and highly reversible defect.

Summary

Serum siderophilin levels were profoundly depressed in a series of 21 patients with kwashiorkor.

The level correlates with independent grading of the severity of the disease.

Siderophilin level was far more profoundly depressed than the total serum protein level. It correlated poorly with the packed cell volume.

The defect was rapidly reversible by high protein diet.

The possible usefulness of this simple test in

diagnosis, in grading the severity of the disease, and in research on means of treatment is discussed.

We are grateful to Professor R. G. Hendrickse for advice and help in this study, to Dr. H. C. Goodman of the World Health Organization Immunology Division for support to one of us (J.F.S.) as Temporary Consultant during the period of this study, and to Mr. A. O. Bameke, A.M.I.L.T. of the Department of Chemical Pathology for technical assistance.

REFERENCES

- Edozien, J. C. (1960). The serum proteins in kwashiorkor. *J. Pediatr.*, **57**, 594.
- , and Udeozo, I. O. K. (1960). Serum copper, iron and iron binding capacity in kwashiorkor. *J. trop. Pediatr.*, **6**, 60.
- Gornall, A. G., Bardawill, C. J., and David, M. M. (1949). Determination of serum proteins by means of the biuret reaction. *J. biol. Chem.*, **177**, 751.
- McFarlane, H., and Udeozo, I. O. K. (1968). Immunochemical estimation of some proteins in Nigerian paired maternal and fetal blood. *Arch. Dis. Childh.*, **43**, 42.
- Neale, F. C., and Soothill, J. F. (1958). Unpublished data. Quoted by Soothill, J. F. (1967). Quantitative disturbances of plasma proteins in disease. *Sci. Basis Med. ann. Rev.*, 276.
- Soothill, J. F. (1962). Estimation of 8 serum proteins by a gel diffusion precipitin technique. *J. Lab. clin. Med.*, **59**, 859.

The following articles will appear in future issues of this Journal:

- Review Article: Disorders of the Serum Lipoproteins. II: Hyperlipoproteinaemic States.* By June K. Lloyd.
- Effect on Height of Corticosteroid Therapy in Asthmatic Children.* By K. F. Kerrebijn and J. P. M. de Kroon.
- Long-term Dwarfing Effects of Corticosteroid Treatment for Childhood Nephrosis.* By C. N. Lam and G. C. Arneil.
- Malaria Prophylaxis and Immunosuppressant Therapy in Management of Nephrotic Syndrome Associated with Quartan Malaria.* By J. W. Kibukamusoke.
- Diagnosis and Treatment of Tyrosinosis in a Girl.* By A. Fairney, D. Francis, R. S. Ersser, J. W. T. Seakins, and D. Cottom.
- Hypertension and Excretion of 1-oxygenated Steroids.* By R. W. H. Edwards, D. R. Harvey, and E. Knight-Jones.
- Growth Hormone Secretion in Subjects with Ovarian Dysgenesis and Turner's Syndrome.* By S. R. Meadow, B. J. Boucher, K. Mashiter, M. N. R. King, and L. Stimmler.
- Familial Haemophagocytic Reticulosis.* By R. J. M. Bell, A. J. E. Brafield, N. D. Barnes, and N. E. France.
- Primary Congenital Hydrocephalus. Long-term Results of Controlled Therapeutic Trial.* By J. Lorber and R. B. Zachary.
- Extensive Vascular Calcification in Association with Juvenile Rheumatoid Arthritis and Amyloidosis.* By M. McC. Reid and T. F. Fannin.
- Glycogen Storage Disease Type IV, Amylopectinosis.* By B. Levin, E. A. Burgess, and P. E. Mortimer.
- Clinico-pathological Study of Thymic Dysplasia.* By C. L. Berry and E. N. Thompson.
- Simple Tests of Ventilatory Capacity in Children with Cystic Fibrosis. Part I: Clinical and Radiological Findings in 85 Cases. Part II: Three-year Follow-up on 50 Patients.* By M. B. Mearns.
- A Working Population Model for Haemolytic Disease of the Newborn.* By E. G. Knox and W. Walker.
- Internal Anal Sphincter. Observations on the Development and Mechanism of Inhibitory Responses in Premature Infants and Children with Hirschsprung's Disease.* By E. R. Howard and H. H. Nixon.