2SD for breast fed) and for magnesium as 1.42. All cases had one or both ions below these levels. 39 of the 75 cases showed hypomagnesaemia with hypocalcaemia and 5 had hypomagnesaemia alone. 33 were treated with magnesium and calcium and 9 with magnesium alone. There were 9 infants who showed the hypotonic normo-alerted clinical picture of hypermagnesaemia as a result of therapy. CSF calcium and magnesium levels were significantly lowered ($p < 0.005$ Mg, < 0.0025 Ca).

The infants were fed on full-cream Regal milk which has a high phosphate (90 mg./100 ml.) and a high osmolality.

OFD Syndrome. J. A. Dodge. Oral-facial-digital syndrome (OFD) is one of a few inherited disorders which are virtually limited to the female sex, and at least 2 of the 7 reported male cases had XXY karyotypes. Rimoin and Edgerton (1967) drew attention to a very similar disorder (OFD II, or Mohr syndrome) which is inherited as an autosomal recessive and may, therefore, occur in males. Examples of both syndromes have been encountered in Northern Ireland. 'Classical' OFD was seen in the first and third generations of a pedigree, but the single living patient in the second generation had very minor dental and dermatoglyphic abnormalities which would have been overlooked or disregarded but for the manifestations in her daughters. She is, therefore, regarded as a true '*forme fruste*'. Her mother and four daughters all had lobulated tongues with hamartomata, supernumerary frenula, and gross dental malformations. Another female in the second generation is reported to have had typical OFD features, but she died at the age of 8 months; she had the same mother as the '*forme fruste*', but a different father. This is further evidence for genetic transmission as a dominant characteristic, through the female—a rare mode of inheritance which may be explained by postulating that the gene is X-linked, lethal when hemizygous. This hypothesis is supported by the finding of Klinefelter's syndrome in some of the affected males. In order to investigate further the suggestion of X-linkage, the Xg blood groups of the pedigree were determined. The findings indicated that recombination had occurred in 3 out of 4 patients in the third generation. If the OFD gene is carried on the X chromosome, it is likely to be situated a long way from the Xg locus.

REFERENCE

- Rimoin, D. L., and Edgerton, M. T. (1967). Genetic and clinical heterogeneity in the oral-facial-digital syndromes. *Journal of Pediatrics*, 71, 94.

Vertebral Abnormalities in First Degree Relatives of Cases of Spina Bifida and of Anencephaly. K. M. Laurence. The parents and sibs aged over 5 years of 100 cases of anencephaly, spina bifida cystica, and encephalocele were x-rayed to determine spinal abnormalities. All these individuals were matched for age and sex with cases seen in the accident unit. Verte-

bral abnormalities found were divided into the mild and severer groups.

Sibs showed 54% mild and 26% severer forms of spina bifida occulta, as compared with 36% and 8% respectively, amongst the controls under 20 years of age. Parents had 20% mild and 4% severer abnormalities, and adult controls had 20% and 1%, respectively.

The results suggest that the severer variety of spina bifida occulta only is probably part of the dysraphic neural tube syndrome.

Hand and Foot Preference in Thalidomide Children. R. W. Smithells. Of 197 thalidomide children, 34% were left-handed and 37% were left-footed. Of those with no arms but normal legs, 70% were left-footed. There was no increase in left-side preference amongst their parents or sibs. The reason for this is obscure, but the observations support the view that a wholly genetic explanation for hand and foot preference is untenable.

Measurement of Adipose Cells in Children. C. G. D. Brook (introduced by June Lloyd). Early nutritional experiences may have lasting consequences on cell growth and behaviour. Animal experiments have demonstrated that different weaning diets have permanent effects on adipose cells in the rat. This paper presents preliminary observations on the measurement of adipose cells in children.

Subcutaneous adipose tissue was obtained at the time of operation from 34 children, aged 3 months to 42 years and of normal weight, who were undergoing elective surgery for various conditions. In 12 of these children additional samples were obtained from deep (intra-abdominal) sites. The lipid content and fatty acid composition were estimated by quantitative gas-liquid chromatography in all samples. In 21 of the children the number of cells was counted by method III of Hirsch and Gallian (1968), and total adipose tissue mass estimated from skin-fold thicknesses using the regression equations of Durnin and Rahaman (1967).

There was a small but statistically significant difference between the mean lipid content of adipose tissue (% wet weight) taken from subcutaneous and deep sites, though the fatty acid composition did not differ. In those children in whom biopsies were taken from both subcutaneous and deep sites the cell size of the deep adipose tissue was considerably less than that of the subcutaneous tissue. However, because deep fat forms only a minor part of the total adipose tissue mass the difference in cell size was ignored in estimating the total number of adipose cells in the body. This has been calculated from the mean cell size of subcutaneous adipose tissue; other workers have shown that subcutaneous cells from different parts of the body are of similar size (Hirsch, Knittle, and Salans, 1966), and this has been confirmed. Results so far obtained indicate a linear relation between total cell number and chronological age in children of normal weight. Preliminary observations in a few obese children from whom

adipose tissue was obtained by subcutaneous aspiration indicate that the cell size is greatly increased and that total cell number may also be increased.

REFERENCES

- Durnin, J. V. G. A., and Rahaman, M. M. (1967). Assessment of amount of fat in the human body from measurements of skinfold thickness. *British Journal of Nutrition*, **21** 681.
- Hirsch, J., and Gallian, E. (1968). Methods for the determination of adipose cell size in man and animals. *Journal of Lipid Research*, **9**, 110.
- , Knittle, J. L., and Salans, L. B. (1966). Cell lipid content and cell number in obese and non-obese human adipose tissue (abst.). *Journal of Clinical Investigation*, **45**, 1023.
- with adrenocortical carcinomas (ages 1 and 27 years), and one with the 11-hydroxylase defect form of congenital adrenal hyperplasia. 97 normotensive patients and healthy persons did not show 1-oxygenated steroids on systematic analysis of urinary steroids by two-dimensional chromatography.

Growth Hormone Production During Puberty in Hypopituitarism. P. H. W. Rayner. The pubertal growth spurt has in the past been attributed to gonadal and adrenal androgens and oestrogens. Though it is recognized that the pituitary output of growth hormone (GH) after provocative stimulation is increased during puberty, the relative importance of GH and sex hormones in linear growth promotion remains undecided. Several workers have shown that testosterone administration promotes an increased GH production in sexually retarded males, and pre-treatment with oestrogen is known to enhance the pituitary GH response to arginine infusion.

We have assessed GH production after pituitary stimulation in 5 patients with hypopituitarism (4 males and 1 female), both before and during puberty. In 3, the aetiology of the pituitary defect was unknown, one followed head injury and one had a chromophobe adenoma. In 3 cases an isolated deficiency of GH was present and in 2 it was combined with thyrotrophin deficiency.

When assessed before puberty between 10 and 13½ years all were growing at less than 3.0 cm./year, and all produced levels of plasma GH measured by radio-immunoassay of less than 9 ng./ml. Puberty occurred spontaneously with accelerated rates of linear growth ranging from 5 to 9 cm./year. Further assessment during puberty showed increased levels of plasma GH in all cases ranging from 10 to 50 ng./ml.

These findings confirm that there is a direct interplay between the gonadal hormones and the pituitary GH-releasing system and suggest that the secretory pattern of GH is dependent on the concentration of circulating gonadal hormones. Gonadal hormones may promote linear growth by increasing the production of pituitary growth hormone.

Concentration of IgE Immunoglobulin in Serum of Children and its Clinical Significance. C. B. S. Wood. The single radial diffusion method of measuring protein concentration has been adapted for the measurement of IgE immunoglobulin in serum, by increasing its sensitivity using a sandwich technique. Sera from a series of allergic and non-allergic children have been studied and the concentrations of IgE in the sera of allergic children were found to be significantly raised. Atopic asthmatics who were below average in weight tended to have high IgE concentrations. The majority of asthmatic children who responded to disodium cromoglycate had high serum IgE concentrations, but a small minority were identified in whom response was good but the IgE concentration was not detectably raised.

Absorption of Antibiotics in Cystic Fibrosis.

H. B. Valman (introduced by B. Wharton). 40 children with cystic fibrosis were given cloxacillin or erythromycin, and the serum protein unbound levels, i.e. the biologically active fractions, were determined 1½ and 4 hours after the administration of cloxacillin and 2 and 4 hours after erythromycin. These levels were compared with the concentration of antibiotics required to inhibit the growth of 7 strains of *Staphylococcus pyogenes* isolated from the same patients. The average serum levels did not differ significantly from those previously reported in normal children receiving similar doses, indicating that there was no malabsorption of the antibiotics in the group of patients with cystic fibrosis. However, in some individuals the serum unbound levels did not exceed the level required to inhibit the growth of the patient's own staphylococci *in vitro*. This suggests that some children with cystic fibrosis need higher than conventional doses of oral antibiotics for elimination of their staphylococci.

Changes in Major Vessels During Growth and After Operation.

C. L. Berry.

Plasma Insulin, Glucose, and Cortisol Levels During and After Exchange Transfusion.

H. V. Price (introduced by O. P. Gray). Hypoglycaemia is a recognized complication of erythroblastosis fetalis. 21 exchange transfusions were studied. The group as a whole showed a steady rise in plasma glucose and insulin levels. Response in individual babies varied widely, particularly the insulin. These changes could not be correlated with plasma cortisol, birthweight, gestation, jaundice, or severity of the haemolytic disease. Plasma cortisol levels varied with each infant but no consistent rise was seen.

Steroid-Endocrine Hypertension.

R. W. H. Edwards. The occurrence of hypertension as a facet of Cushing's syndrome has often been noted and remarked upon, and generally ascribed to over-secretion of mineralocorticoid. It is now interesting to rediscuss the matter because of the recent association of urinary 1-oxygenated steroids with hypertension. Patients showing hypertension and excretion of 1-oxygenated steroids include two who might be said to suffer from idiopathic hypertension (ages 1 week and 2 years), two