- Dartois, A. M., and Lestrade, H. (1966). Food consumption of diabetic children at summer camps in France. *Journal of the American Dietetic Association*, **48**, 32.
- Dorchy, H., and Loeb, H. (1975). Valeur comparée de différents tests semi-quantitatifs pour la détermination de la glucosurie chez les diabétiques. *Diabète et Métabolisme (Paris)*, **1**, 87-90.
- Dorchy, H., Mozin, M. J., and Loeb, H. (1975). Alimentation des enfants et adolescents diabétiques. *Revue Médicale de Bruxelles*, **31**, 391-392.
- Dorchy, H., Devroede, M., Toussaint, D., and Ernould, C. (1976). Angiofluorescein studies in infantile diabetic retinopathy. *Proceedings of the 2nd Meeting of the International Study Group of Diabetes in Children and Adolescents* (in press).
- Drash, A. (1976). The control of diabetes mellitus. Is it achievable? Is it desirable? *Journal of Pediatrics*, **88**, 1074-1076.
- Ernould, C., Dorchy, H., Francois, B., Geussens, H., Mozin, M. J., and Loeb, H. (1973). *Guide du Jeune Diabétique*, p. 51. Revue Médicale de Liège, Liège.
- Ernould, C., Devroede, M., Dorchy, H., and Loeb, H. (1976). Observation of growth in diabetic Belgian children and adolescents. *The Balance of Diabetes in Juveniles*. Ed. by Z. Laron. Karger, Basel (in press).
- François, R. (1976). Réflexions sur le traitement du diabète infantile insulino-dépendant. *Archives Françaises de Pédiatrie*, **33**, 425-430.
- Lestrade, H., and Billaud, L. (1968). Complications dégénératives après 15 ans de diabète infantile. *Presse Médicale*, **76**, 303-306.
- Lestrade, H., Dartois, A. M., and Machinot, S. (1974). L'alimentation spontanée de l'enfant et de l'adolescent diabétiques traités par l'insuline. *Annales de Pédiatrie*, **21**, 667-676.
- Mozin, M. J., Dorchy, H., Ernould, C., Francois, B., Geussens, H., and Loeb, H. (1975). Observations concerning the spontaneous feeding of a group of diabetic adolescents in a vacation camp. *Diabetes in Juveniles*, p. 385. Modern Problems in Paediatrics, Vol. 12. Ed. by Z. Laron. Karger, Basel.
- Mozin, M. J., Dorchy, H., Smets, P., Ernould, C., and Loeb, H. (1976). Spontaneous variations of intakes and balance of diabetes in diabetic adolescents. *Acta Paediatrica Belgica* (in press).
- Rimoin, D. L., and Schimke, R. N. (1971). *Genetic Disorders of the Endocrine Glands*. Mosby, St. Louis.

Achondrogenesis type I

Sir,

We read with great interest the article by Lauder *et al.* (*Archives*, 1976, **51**, 550). Their observation is correct in that their 2 cases are subvariants of achondrogenesis type I according to the McKusick classification.

We would like to point out that two distinct subtypes of that condition have already been described (Spranger *et al.*, 1974; Yang *et al.*, 1974, 1975). According to the criteria developed in our studies, the 2 cases reported by Lauder *et al.* should be classified into type I of lethal achondrogenesis (Yang *et al.*, 1974, 1975, 1976, 1977), which is characterized by deficient cranial ossification, multiple rib fractures, shorter appendicular bones,

sufficient cartilage matrix in the epiphyseal cartilage, higher incidence of familial occurrence, etc. Recently we have observed intracytoplasmic inclusion bodies in the resting chondrocytes (Yang *et al.*, 1976, 1977). We suggest that Dr. Lauder and colleagues re-evaluate their histological sections of cartilage, which seem in the low magnification photomicrographs to contain several inclusions.

Type II lethal achondrogenesis differs roentgenographically and morphologically from type I. Grebe achondrogenesis, formerly type II, is nonlethal and appears to be an unrelated disease.

SHENG-S. YANG and
JAY BERNSTEIN

William Beaumont Hospital,
Royal Oak, Michigan 48072,
USA

Drs. Lauder and Ellis reply as follows:

We are grateful to Drs. Yang and Bernstein for their comments on our paper on achondrogenesis type I. At the time of preparation of our manuscript we were not aware of their recent paper (Yang *et al.*, 1974) in which they report 2 cases, one of which exhibits similar clinical and pathological features to our own 2 cases. We agree with them that the available evidence seems to suggest two distinct forms of achondrogenesis type I. It remains our view that until the precise biochemical defects have been elucidated, no definitive classification is possible. This is also the view expressed by Wiedemann *et al.* (1974) whose collection of 6 cases is still the largest reported. The difficulties in attempting a classification based on 2 cases plus a review of previous studies are exemplified by the necessity for a 'readjustment of eponyms' which Yang *et al.* (1975) subsequently reported. As our paper points out, many of the studies are unsatisfactory in that no comments are made regarding the appearances in undecalcified material. The absence of adequate biochemical and histochemical data in these reports also makes distinction from conditions such as severe hypophosphatasia extremely difficult. It was also our aim to stress the importance of pulmonary hypoplasia as a cause of death in these infants.

We were most interested in the observation by Yang *et al.* (1976) of intracytoplasmic inclusion in resting chondrocytes. We have not had the opportunity to study their report or any illustrations of these bodies. We have not as yet been able to convince ourselves of the presence of inclusion bodies in our material.

I. LAUDER and H. A. ELLIS
Department of Histopathology,
Royal Victoria Infirmary,
Queen Victoria Road,
Newcastle upon Tyne NE1 4LP.

References

- Spranger, J. W., Langer, L. O., Jr., and Wiedemann, H.-R. (1974). *Bone Dysplasias*, p. 24. Saunders, Philadelphia.
- Wiedemann, H. R., Remagen, W., Hienz, H. A., Gorlin, R. J., and Maroteaux, P. (1974). Achondrogenesis within the scope of connately manifested generalized skeletal dysplasias. *Zeitschrift für Kinderheilkunde*, **116**, 223-225.