dependent on systemic blood pressure. It is this possible link between blood pressure and cerebral bleeds that makes the work of Fujimura *et al.* of such interest.

References

- ¹ Cross K W, Dear P R F, Warner Ruth M, Watling G B. An attempt to measure cerebral blood flow in the newborn infant. *J Physiol (Lond)* 1976; **260**: 42–3P.
- ² Cross K W, Dear P R F, Hathorn M K S, *et al.* An estimation of intracranial blood flow in the newborn infant. *J Physiol (Lond)* 1979; **289**: 329–45.
- ³ Rahilly P M. The responses of cranial blood flow in the newborn infant to some common stimuli. MD thesis, London University, 1979: 82–9.
- ⁴ Lou H C, Lassen N A, Friis-Hansen B. Low cerebral blood flow in hypotensive perinatal distress. *Acta Neurol Scand* 1977; **65**: 343–52.

P M RAHILLY
Department of Medicine,
Royal Alexandra Hospital for Children,
Pyrmont Bridge Road,
Camperdown,
NSW 2050,
Australia

Giant exomphalos—conservative or operative treatment?

Sir,

I would like to comment on the successful use of conservative treatment of giant exomphalos by Gough and Auld (Archives, 1979, **54**, 441). Their 6 patients on conservative treatment had the intact sac painted daily with 2% aqueous mercurochrome. Two of these patients had complications of mercury intoxication, one with failure to thrive at age 8 weeks and one who became mentally retarded with microcephaly. The risk of mercury poisoning in conservative treatment of giant exomphalos can be avoided by using 1% aqueous gentian violet. I successfully treated 6 infants with exomphalos measuring

more than 5 cm in diameter in Singapore between 1969 and 1972. Three had the intact sac painted with 1% aqueous gentian violet and 3 with 2% aqueous mercurochrome. All survived and none developed complications during the 2 years that they were followed up. The infants treated with 1% gentian violet paint responded with epithelialisation and healing of the abdominal defect within 12 weeks while those treated with mercurochrome did not heal until 20 weeks. The only disadvantage of using gentian violet was that it stained the linen.

All my patients required secondary surgery to the ventral hernia which replaced the exomphalos.

M C K CHAN
Department of Tropical Paediatrics,
Liverpool School of Tropical Medicine,
Pembroke Place, Liverpool L3 5QA

Mr Gough comments:

I agree that mercury intoxication remains a significant risk when mercurochrome applications are used daily in the management of large omphaloceles. However, one of our patients had 3 exchange transfusions for hyperbilirubinaemia and it is difficult to know what part this played in his subsequent poor progress. The second case illustrated in the article has developed normally despite high urine mercury excretion when aged 8 weeks, and thus the risks may have been overstated.

I have recently treated a patient with gentian violet applications, as Dr Chan describes, and can confirm the favourable progress which he noted, and his letter gives further support to the concept that conservative treatment is both effective and safe.

D C S GOUGH FRCS
Royal Manchester Children's Hospital,
Pendlebury,
Manchester M27 1HA