cally incorporated into 'standard practice' as the sine qua non of abuse. Equal open mindedness to the interpretation of social factors in such cases should be maintained until a definite diagnosis has been made.

Like any other haemorrhage, those in the optic fundus may be traumatic or spontaneous. Among the spontaneous causes, a bleeding diathesis should be considered.¹ Diagnostic evaluation by computed tomography of the brain is also important. A localised intracerebral haematoma is most unlikely to be traumatic if there is no other evidence of parenchymal injury. Shaking may cause generalised brain contusion.³ Alternatively, a subdural haemorrhage may be induced and a posterior interhemispheric distribution is particularly characteristic of shaking injury (Fig. 2).⁴

Recognition of a subdural haemorrhage by aspiration through the anterior fontanelle should also be considered critically. Any intracranial haemorrhage may leak into the subarachnoid space and a subdural haematoma can develop if an aneurysm or arteriovenous malformation ruptures through the arachnoid mater. Such a finding has been described in a 3 month old child with an intracerebral aneurysm.⁵ Arteriovenous malformations may also occur within the meninges and so bleed directly into the subdural space. We can see little value in performing diagnostic subdural taps when access to computed tomography facilities are available.

We suggest that a computed tomogram of the brain should be considered urgently in the evaluation of cases of possible shaking injury or unexplained fundal haemorrhages. This should clarify the appropriate course to follow and prevent misinterpretation. A ruptured intracerebral arterial aneurysm may be eminently suitable for curative neurosurgery.⁶

We thank Dr Ian Holland for his neuroradiological expertise.

References

- ¹ Caffey J. The whiplash shaken infant syndrome: manual shaking by the extremities with whiplash induced intracranial and intraocular bleedings, linked with residual permanent brain damage and mental retardation. *Pediatrics* 1974;**54**:396-403.
- ² Ludwig S, Warman M. Shaken baby syndrome: a review of 20 cases. *Ann Emerg Med* 1984;**13**:104-7.
- ³ Ellison PH, Tsai FY, Largent JA. Computed tomography in child abuse and cerebral contusion. *Pediatrics* 1978;**62**:151-4.
- ⁴ Zimmerman RA, Bilaniuk LT, Bruce D, *et al.* Computed tomography of craniocerebral injury in the abused child. *Radiology* 1979;**130**:687-90.
- ⁵ Vapalanti PM, Schugk P, Tarkkanen L, Af-Björkstén G. Intracranial arterial aneurysm in a 3 month old infant. *J Neurosurg* 1969;**30**:169-71.
- ⁶ Crisostomo EA, Leaton E, Rosenblum EL. Features of intracranial aneurysms in infants and report of a case. *Dev Med Child Neurol* 1986;**28**:68-72.

Correspondence to Dr N McLellan, Consultant Paediatrician, Selly Oak Hospital, Raddlebarn Road, Birmingham B29 6JD, England.

Received 19 May 1986

Topical anaesthesia for venepuncture

S CLARKE AND M RADFORD

Department of Child Health, Southampton General Hospital

SUMMARY A topical anaesthetic cream was tested in a randomised, double blind, placebo controlled trial of 15 children. The severity of pain experienced during venepuncture was assessed, using visual analogue and verbal rating scales. The topical anaesthetic cream was found to be significantly superior to placebo using each form of assessment.

Venepuncture is probably the commonest procedure carried out on paediatric patients requiring medical treatment. Many children tolerate the procedure well, but for some it is a major ordeal. A topical preparation that could be applied without

discomfort and alleviate the pain of venepuncture would be particularly useful for children receiving repeated venepunctures.

To penetrate intact skin a topical anaesthetic cream should contain local anaesthetic in high concentrations. This has hitherto been difficult to achieve because most anaesthetic bases are crystalline at room temperature and unstable in high concentrations. By mixing two anaesthetic bases, lignocaine and prilocaïne, in equal proportions, an eutectic mixture is obtained that is liquid at room temperature. This mixture, known as EMLA (Eutectic Mixture of Local Anaesthetics), can be incorporated into an emulsion without first dissolving the bases in oil, making it possible to attain a

The aim of the present study was to evaluate the analgesic effect of the anaesthetic cream with respect to venepuncture pain in children with leukaemia who were attending for repeated venepunctures.

Fifteen children receiving maintenance treatment for lymphoblastic leukaemia were asked to take part in the trial. Informed consent was obtained from patients and parents and the trial was approved by the local ethical committee.

Each patient was assigned a pack of four tubes. These were labelled 1-4 in order of administration and contained the treatment sequences A₁ P₂ A₃ P₄ or P₁ A₂ P₃ A₄ (A=active, P=placebo) as specified by a randomisation list. The trial was conducted double blind. The first application for each child was made in the clinic by the investigator, after which tubes 2-4 were taken home for the cream to be applied by a parent before subsequent clinic visits. The cream was always applied in a thick layer (roughly 2 ml each application) under an occlusive dressing, Tegaderm 3M, for a period of at least 60 minutes before performing the venepuncture. At the end of this time the dressing and cream were removed and the site cleansed with 0.5% chlorhexidine in 70% ethanol. Venepunctures were performed on the dorsum of the hand or the antecubital fossa, using a 21 or 23 gauge needle. A total of seven clinicians and nursing staff were involved in performing the venepunctures.

made by the child. In the case of younger children the assessment was made with the aid of the parent.

At the end of the trial the value of the active cream was assessed by asking the parents whether they found the cream useful and worth while.

Results

Severe pain

Venepuncture pain (mm)

Mean —

No pain Placebo Active

p<0.0005, Friedman non-parametric two way ANOVA.

The distribution of pain scores on the visual analogue scales is shown in the Figure. There was a highly significant difference between the values for the active cream and the placebo ($p<0.0005$).

The results on the four point verbal response scale of venepuncture pain also showed that the active cream was significantly superior to placebo. (Fisher's exact test, $p<0.01$) There was no difference between the two treatments with respect to ease of venepuncture.

Most parents and patients considered the use of the active cream worth while. Parents were able to apply the cream to a suitable vein at home without difficulty. There were several occasions when patients had to wait for the cream to take effect, but this inconvenience was considered to be offset by the advantage of less painful venepuncture. Eleven of the 15 children or their parents said they would like to continue using the active cream.

There were occasional skin reactions, consisting of pallor or redness, to both active cream and placebo. These were transient and clinically unimportant.

Discussion

The results of this study clearly indicate that an anaesthetic cream is an effective agent for alleviation of venepuncture pain. It is slightly inconvenient to use, but many children requiring regular venepuncture seem to be sufficiently motivated to accept this. Most children in this study elected to continue using the active cream for subsequent venepuncture.

The factors affecting absorption of local anaesthetic through intact skin have been studied by Evers, who found it necessary to use a thick layer of cream as emulsion droplets in contact with the skin readily became depleted of local anaesthetic.³ In the same study percutaneous absorption of lignocaine and prilocaine into the systemic circulation was found to be measurable but slight.

As in all local anaesthetic preparations it is necessary to consider the potential for hypersensitive reactions. There is an extensive clinical experience with lignocaine and prilocaine given by injection, and true allergic reactions are extremely rare. There is no reason to believe that these agents applied topically will behave differently.

We acknowledge the help of Astra Pharmaceuticals Ltd.

References

- 1 Ehrenstrom-Reiz GMS, Reiz SLA. EMLA—a eutectic mixture of local anaesthetics for topical anaesthesia. *Acta Anaesthesiol Scand* 1982;26:596.
- 2 Tregear RT. Molecular movement. The permeability of the skin. In: Tregear RT, ed. *Physical functions of skin*. Academic Press, 1966:1–52.
- 3 Evers H, von Dardel O, Juhlin L, Ohlsen L, Vinnars E. Dermal effects of compositions based on the eutectic mixture of lignocaine and prilocaine (EMLA). *Br J Anaesth* 1985;57: 997–1005.

Correspondence to Dr M Radford, Department of Child Health, Centre Block, Southampton General Hospital, Southampton SO9 4XY, England.

Received 30 June 1986.

Meningitis due to *Haemophilus influenzae* resistant to ampicillin and chloramphenicol

A P FRAISE, A C G MEEKS, AND J E RICHARDS

Department of Microbiology/Public Health Laboratory and Department of Child Health, Leicester Royal Infirmary

SUMMARY A case of ampicillin and chloramphenicol resistant *Haemophilus influenzae* meningitis successfully treated with cefotaxime is described.

Meningitis due to *Haemophilus influenzae* resistant to chloramphenicol and ampicillin is a recognised

problem in the United States¹ but has only been reported once in the United Kingdom.² To our knowledge the case reported here is the first British case to be treated with cefotaxime.

Case report

A previously healthy 7 month old boy presented in