British paediatrics

Perinatal care

The British Paediatric Association (BPA) has recently issued its responses to two official reports relating to the care of the newborn baby—namely, the report of the Social Services Committee (SSC) on perinatal and neonatal mortality and the report of the Maternity Services Advisory Committee (MSAC) on postnatal and neonatal care. Both reports are welcomed by the BPA but, as might be expected, not all the specific recommendations meet with the Association's approval.

According to the SSC the only body in a position to produce generally acceptable national minimum standards is the MSAC. The BPA strongly disagrees with this view and refers to the reports it has prepared in conjunction with the British Association for Perinatal Paediatrics and the Royal College of Obstetricians and Gynaecologists (RCOG), in which widely recognised minimum standards have been drawn up. The minimum standards for nursing and medical staff in neonatal units as proposed by the MSAC are not accepted by the BPA. The recommended level of nursing staff in neonatal units is totally inadequate, and the Association has strong reservations about the statement that in every neonatal unit there should always be a nurse in charge who has at least one year's experience in neonatal nursing. In a neonatal unit providing intensive care it is not acceptable to have a nurse at less than the grade of sister in charge. Furthermore, the recommendations for paediatric medical staff are not sufficiently strong. For example, the MSAC states that 24 hour medical cover for the care of the newborn should be provided by a paediatrician 'ideally on site' and that in regional perinatal centres there should be 'at least one consultant perinatal paediatrician'. The BPA considers that 24 hour cover provided by a paediatrician on site is not just ideal but essential and that there should be an absolute minimum of two consultant perinatal paediatricians in a regional centre.

Although both committees support the principle of 24 hour paediatric cover, which implies the provision of adequate numbers of junior posts in paediatrics, the government has stated that there can be no increase in such posts as this would mean training more doctors than there are likely to be

career outlets available. The BPA points out that this view does not take account of the fact that junior posts in paediatrics are suitable for training general practitioners and consultants in other specialties and therefore do not need to be rigidly related to higher grade posts in paediatrics alone.

The two reports recommend that each region should make adequate provision for neonatal intensive care, but only the SSC is prepared to place the responsibility for making available the necessary resources on the Secretary of State. Regional health authorities will clearly have to determine the proper distribution of intensive care cots throughout their regions, set up regional referral units, and ensure that districts have the facilities to undertake short term ventilation before transfer. The BPA has evidence from at least two regions that babies are being refused admission to intensive care units through lack of beds. It is important, therefore, that the scale of this problem should be assessed. It would also be helpful if information were gathered about refusals of antenatal transfer. According to the BPA there should be at the very least one intensive care cot per 1000 annual deliveries, but the MSAC was not prepared to recommend a minimum number of cots, in spite of the advice of the paediatricians on the committee that this was essential. The strategy of the MSAC is that regions should plan to make up deficiencies and develop their neonatal services over a period of 10 years, but this is seen by the BPA as too slow. There should be a two year planning and consultation period with three more years for implementation.

A policy statement in the MSAC report states that babies who need a measure of extra care short of continuous skilled supervision should be nursed alongside their mothers, but no mention is made of intermediate or transitional care, which enables nursing skills to be concentrated on a single postnatal ward. Wherever they are nursed babies needing extra care will require a more generous allocation of nurses than on a normal postnatal ward. The SSC recommends that more National Board Courses in special and intensive neonatal nursing be established and that qualifications gained in these courses be recognised in pay levels. The

BPA is in favour of this but considers that only those nurses actually working in special and intensive care units should receive the extra payments for these qualifications.

A recommendation of the SSC, which the BPA considers crucially important, is that the long term health of babies treated in neonatal intensive care units should be monitored. The Oxford Region's Child Development Project could, in the Association's view, be a model for such a monitoring system, and the increasing number of consultant paediatricians with an interest in child health could help its implementation.

The MSAC report includes a list of recommended equipment for neonatal units that, it says, will have to be regularly reviewed in the light of up to date information from the BPA and other appropriate professional bodies. While approving this list, the BPA regrets the omission of a statement on the need for a budgeted replacement programme for obsolete or worn out equipment. Replacement of such expensive items is a constant concern to paediatricians, who are forced to spend an increasing amount of time seeking charitable funds for this purpose. Reliance on charity money is, in any case, of questionable propriety in a centrally funded health service.

There are a number of recommendations that the BPA warmly welcomes. For example, the SSC recommends that the recognised nursing establish-

ments in neonatal special and intensive care units should be at the level proposed by the joint committee of the BPA and RCOG and that there should be a greatly increased rate of consultant expansion in obstetrics and paediatrics. The BPA also supports the recommendations of both committees that there should be individual records for newborn babies, that regional plans should incorporate provision for neonatal surgery in regional perinatal centres, and that full time perinatal pathologists should be appointed in all regions. The SSC is strongly supported by the BPA in its encouragement of both the Department of Health and Social Security and the Medical Research Council to give high priority to the funding of research in prenatal, perinatal, and neonatal medicine.

Both reports were, on the whole, well received by the BPA. The MSAC report was considered to be a useful guide to good practice, but it was thought that its advice should have been couched in stronger terms. The SSC report echoes precisely the views of the BPA when it concludes that 'the downward trend in perinatal and neonatal mortality can only be sustained if resources are directed towards improving services in areas of deprivation and if authorities devote more effort to the better organisation of services'.

A D M JACKSON