- ² Creighton SJ. *Child abuse in 1985: initial findings from NSPCC register research*. Research Briefing No 7. London: NSPCC, 1986.
- ³ Krugman RD. Recognition of sexual abuse in children. *Pediatrics in Review* 1986;**8**:25–30.
- ⁴ Hunter RS, Kilstrom N, Loda F. Sexually abused children: identifying masked presentations in a medical setting. *Child Abuse Negl* 1985;**9**:17–26.
- ⁵ Heger A. *Response child sexual abuse: a medical view*. Los Angeles: United Way and Children's Institute International, 1985.
- ⁶ Hobbs CJ, Wynne JM. Buggery in childhood – a common syndrome of child abuse. *Lancet* 1986;**ii**:792–6.
- ⁷ Enos WF, Conrath TB, Byer JC. Forensic evaluation of the sexually abused child. *Pediatrics* 1986;**78**:385–8.
- ⁸ McCauley J, Gorman RL, Guzinski G. Toluidene blue in the detection of perineal lacerations in pediatric and adolescent sexual abuse victims. *Pediatrics* 1986;**78**:1039–43.
- ⁹ Hammerschlag MR, Cummings M, Doraiswamy B, Cox P, McCormack WM. Nonspecific vaginitis following sexual abuse in children. *Pediatrics* 1985;**75**:1028–31.
- ¹⁰ Hammerschlag MR, Doraiswamy B, Alexander ER, Cox P, Price W, Gleyzer A. Are recto-genital chlamydial infections a marker of sexual abuse in children? *Pediatr Infect Dis* 1984;**3**:100–4.
- ¹¹ Vizard E. Interviewing young, sexually abused children – assessment techniques. *Family Law* 1987;**17**:28–33.
- ¹² Summit RC. The child sexual abuse accommodation syndrome. *Child Abuse Negl* 1983;**7**:177–93.
- ¹³ Racusin RJ, Felsman JK. Reporting child abuse: the ethical obligations to inform parents. *J Am Acad Child Psychiatry* 1986;**25**:485–9.
- ¹⁴ Department of Health and Social Security. *Child abuse – working together*. London: HMSO, 1986.
- ¹⁵ Jones DPH, McGraw JM. Reliable and fictitious accounts of sexual abuse in children. *Journal of Interpersonal Violence* 1987;**2**:25–45.
- ¹⁶ Peters JJ. Children who are victims of sexual assault and the psychology of offenders. *Am J Psychother* 1976;**30**:398–421.
- ¹⁷ Leventhal JM, Hamilton J, Rekedal S, Tebano-Micci A, Eyster C. Results of a diagnostic interview in young children suspected of being sexually abused. Submitted for publication.
- ¹⁸ Fallor KC. Is the child victim of sexual abuse telling the truth? *Child Abuse Negl* 1984;**8**:473–81.
- ¹⁹ de Young M. A conceptual model for judging the truthfulness of a young child's allegation of sexual abuse. *Am J Orthopsychiatry* 1986;**56**:550–9.
- ²⁰ Green AH. True and false allegations of sexual abuse in child custody disputes. *J Am Acad Child Psychiatry* 1986;**25**:449–56.
- ²¹ Benedek EP, Schetky DH. Allegation of sexual abuse in child custody and visitation disputes. In: Schetky DH, Benedek EP, eds. *Emerging issues in child psychiatry and the law*. New York: Brunner/Mazel, 1985:145–56.

Correspondence to Dr J M Leventhal, Yale University School of Medicine, 333 Cedar St, New Haven, Connecticut 06510, USA.

Received 22 June 1987

Commentary

R E I ROBERTS

Manchester

Much of what Hobbs and Wynne say is uncontroversial and merely describes in detail the accepted practice succinctly laid down in the current Department of Health and Social Security (DHSS) guidelines for dealing with child sexual abuse—a multi-disciplinary investigation should take place and the child should be talked to and examined in a sensitive manner in a clinical setting. It is the specific aspects of the medical examination and the conclusions that they draw which are extremely controversial: they state that 40% to 50% of boys and girls have anal abnormalities.

Four women doctors in the Greater Manchester area doing examinations at the request of police, social workers, the National Society for the Prevention of Cruelty to Children (NSPCC), and paediatricians saw approximately 500 children in 1986 (Robinson, Jacobs and Welliver, personal communication). They always take a full history and talk to the child and carer about medical problems as well as any history of abuse. A history of possible anal interference is obtained from a much smaller proportion of these children than from the cases reported by Hobbs and Wynne, and in only a

handful has definite medical evidence in the anal area been found. Precise figures are not yet available but they are in the process of being prepared for publication. Police surgeons throughout the country have made the same observations and the subject was exhaustively discussed at the recent Association of Police Surgeons of Great Britain annual meeting. None of the experienced police surgeons, many of them having the Diploma in Medical Jurisprudence and most of them experienced general practitioners with considerable experience of children, has found anything more than an extremely small proportion of children with anal evidence of sexual abuse.

Hobbs and Wynne say that they can tell merely by inspecting an anus that it is lax. They use the subjective and misleading test of so called 'reflex dilatation' on skin traction on the buttocks to make a definitive diagnosis of 'sexual abuse.' The opening of the anus on inspection—which should be done without traction by gently placing the hands on the cheeks of the buttocks—is variable, misleading, and present in a number of other conditions including chronic constipation, thread-worms, and anal thrush

following courses of antibiotics.¹ Even if it is present in some sexually abused children (and more than half of them have no abnormal medical findings) it cannot be used as evidence (except as possible support to other relevant evidence) because it is also present in some children with other diseases who have not been abused. The same considerations apply precisely to anal twitchiness and dilated veins.

A few years ago I believed that flattening of the labia around the vaginal opening was a good sign of persistent attempted penile penetration, but I am now much more cautious in interpreting this sign because it can occur as part of normal development (and for other reasons) and is not reliable enough as a sign of abuse per se, even though it may have been caused by abuse. The anal findings described by Hobbs and Wynne are subject to the same reservations and much more caution is needed in their interpretation than has been demonstrated hitherto.

The vaginal findings in girls can be difficult to interpret. Cantwell's paper, on which Hobbs and Wynne base their opinion (that 0.5 cm is the accepted upper limit of normal for the size of the hymen), is open to many criticisms.² After the child's hymen was measured (inaccurately) by placing a tape or ruler against the vulva, leading questions were asked of the child. Even after this quite unacceptable practice only 74% of the children with larger openings were diagnosed as having been sexually abused. Is the size to be the same for a babe in arms and a big girl of 11 or 12? There is considerable risk that when paediatricians (and other doctors) look at a child's genitalia they will mistake the vestibule for the hymen and erroneously state that the vaginal opening is too large. A recent paper reported that only 59.1% of doctors including paediatricians correctly identified the hymen on a colour photograph.³ Errors are easy to make when there is a congenital frilly hymen that may be mistaken for tears, and where the doctor's knowledge of anatomy and pathology is inadequate; 'heaped-up scars' of the hymen have been described.

Reddening and bruising around the anus and vagina are of importance where there has been recent traumatic interference but there is great danger in paediatricians with no forensic training and no experience of the aging of bruises and other injuries giving opinions as to the likely cause of such findings. Hobbs and Wynne give no evidence that they know the proper techniques for determining that a discolouration is a bruise, and not, for instance, a birth mark; Reid has emphasised the pitfalls for paediatricians in this area.⁴ When giving evidence in court a doctor who has examined a child

must be able to give an opinion as to what may have been the cause of the findings, how many times it has occurred, what has been used and with what degree of force. It is not enough to say 'consistent with sexual abuse' without being able to say whether a penis has entered the anus or whether it is the tip of a finger. Case conferences and proceedings in juvenile courts for care orders have been satisfied with a much lower standard of evidence than that acceptable in a criminal case, but the consequences for the child and the family may be just as serious and damaging. We must therefore strive to improve the quality of the evidence at all levels and make sure that the diagnosis is based on sound medical and medicolegal principles.

Dr Leventhal gives useful advice about talking to children, but there is increasing concern among doctors and lawyers about the use of the techniques described. Used by a kindly, skilled, and experienced interviewer they can be helpful, but there is a serious danger of them being used inappropriately. 'How much questioning is necessary before being reasonably certain that nothing has happened (given that it may be impossible to exclude completely such a diagnosis?).' Many interviewers using these techniques seem to work on the basis that abuse has taken place and will never take 'no' for an answer. If the child denies abuse it is said to be 'blocking' and must continue to be interviewed until the 'blocking' is released. In my experience children find the repeated interviewing and continual reference to the same subject distressing or boring. Under continual pressure they can become confused and having learnt which answers appear to be acceptable may eventually give the answer sought.^{5 6} Continued interviewing of the child can in some cases constitute an abuse in itself.

Both these papers, though containing much that is useful and helpful also contain dangerous and unfounded assumptions: the uncritical acceptance of these has already led to enormous difficulties for many people trying to help sexually abused children and not least for many children and their families.

References

- 1 Clayden G. Anal appearances and child sex abuse. *Lancet* 1987;i:621.
- 2 Cantwell HB. Vaginal inspection as it relates to child sexual abuse in girls under 13. *Child Abuse Negl* 1983;7:171-6.
- 3 Ladson S, Johnson CF, Doty RE. Do physicians recognise sexual abuse? *Am J Dis Child* 1987;141:411-5.
- 4 Reid DHS. Pitfalls in determining child abuse. *Lancet* 1985;i:1316-7.
- 5 Scobie W. The child abuse scandal that never was. *The Observer* 4 January 1987.
- 6 Collier C, Szwed E. Prompting tales of abuse. *The Guardian* 28 July 1987.