(1) Birth order had no significant influence on the incidence of the disease in boys or girls. Birth order distribution (first born, 42.9%; second, 31.8%, third, 17.9%; fourth, or more 7.4%) did not differ significantly from census data.³

(2) There were significantly more infants with low birth weight (<2500 g) (9.3%) in the group with pyloric stenosis than in the control population (4.9%).⁴

(3) Short gestation pregnancies (<38 weeks) occurred in 16% of mothers of patients with pyloric stenosis (excluding multiple births); this compares with 9% in the general Western Australian population. (Hitchcock NE. Unpublished data.)

(4) Several Aboriginal patients had the disease. To our knowledge, pyloric stenosis has not previously been reported in Australian Aborigines. Unfortunately, the hospital records did not describe how many of these Aboriginal patients were of full descent or were part Aboriginal.

(5) A family history of pyloric stenosis was present in 65 patients (11%), just over half being in the father's family, one quarter in the mother's family, and the rest in siblings. There were 13 multiple births involved. In only two instances were both infants affected, and in one set of all male triplets only one infant was affected. Pyloric stenosis has been shown, in other studies, to be significantly more common in children of affected parents,

especially if the affected parent is the mother^{5 6}—perhaps because the gene responsible for the disease in girls has greater penetrance than that responsible for the disease in boys. This was not our experience in Perth.

We are grateful to the medical records staff of Princess Margaret Hospital for their cooperation and to the Health Department of Western Australia and the Australian Bureau of Statistics for statistical information.

References

- 1 Anonymous. Incidence of infantile hypertrophic pyloric stenosis [Editorial]. *Lancet*. 1984;i:888-9.
- 2 Webb AR, Lari J, Dodge JA. Infantile hypertrophic pyloric stenosis in South Glamorgan 1970-9. Effects of changes in feeding practice. *Arch Dis Child* 1983;**58**:586-90.
- 3 Australian Bureau of Statistics. *Australia Year Book No 66*. Canberra: Australian Bureau of Statistics, 1982.
- 4 Western Australian Midwives Notification System. *Perinatal statistics in Western Australia*. Second annual report of the Western Australian Midwives Notification System for 1984. Perth: Health Department of Western Australia, 1986.
- 5 McKeown T, MacMahon B. Infantile hypertrophic pyloric stenosis in parent and child. *Arch Dis Child* 1955;**30**:497-500.
- 6 Carter CO. The inheritance of congenital pyloric stenosis. *Br Med Bull* 1961;**17**:251-4.

Correspondence to Mrs N E Hitchcock, GPO Box D184, Perth, 6001, Australia.

Received 14 November 1986

Pulmonary haemosiderosis and gluten

R READING, J G WATSON, J W PLATT, AND A G BIRD

Department of Paediatrics, West Cumberland Hospital, Whitehaven, and Departments of Paediatrics and Immunology, Newcastle General Hospital, Newcastle upon Tyne

SUMMARY A child with idiopathic pulmonary haemosiderosis for three years required three monthly transfusions. Circulating avian, gliadin, and reticulin antibodies suggested the diagnosis of gluten enteropathy, and jejunal biopsy showed subtotal villous atrophy. During 15 months on a gluten free diet his growth and behaviour improved and he required no transfusions.

Pulmonary haemosiderosis is characterised by recurrent pulmonary infiltrates due to alveolar haemorrhage, anaemia, and finding haemosiderin laden macrophages in the sputum. In adults certain systemic disorders may be associated, but in chil-

dren no specific cause is usually found. A proportion, however, have circulating antibodies to cow's milk proteins, and withdrawal of cow's milk from the diet may be beneficial.¹

Case report

The boy was born at term weighing 2490 g and was formula fed. At 2 years of age his height and weight lay just below the 3rd centile.

He presented aged 4.5 years with a few days' history of pallor and decline in his chronically poor appetite. His height (1.06 m) and weight (15.5 kg) still lay just below the 3rd centile. An initial haemoglobin concentration of 5.2 g/dl was attributed to iron deficiency, but an erythrocyte sedi-

mentation rate of 50 mm Hg in the first hour was not explained. He was re-admitted three days later with a haemoglobin concentration of 3.9 g/dl, a cough, scattered crepitations, mild eczema, small cervical lymph nodes, and a palpable spleen tip. Investigations showed 5% reticulocytes; iron 3 $\mu\text{mol/l}$, and total iron binding capacity 81 $\mu\text{mol/l}$, a Coombs' test yielded negative results, and stools were occult blood positive three times. He was thought initially to have cardiac failure, but a chest x ray film showed unexpected bilateral pulmonary infiltrates, which ultimately led to the diagnosis of idiopathic pulmonary haemosiderosis. He was transfused and the lung signs and chest x ray film cleared in four weeks.

Over the next 30 months he had 10 similar episodes of cough and pallor associated with infiltrates on his chest x ray film. These no longer cleared completely between episodes and showed an increasing profusion of micronodular shadows. He required repeated blood transfusions (Figure). His height continued parallel to the 3rd centile but his weight fell away slightly. Inhaled disodium cromoglycate for three months had no effect.²

Re-investigation at 7 years gave similar haematological results and folate, vitamin B₁₂, albumin, globulin, zinc, IgG, IgM, IgE, and complement concentrations were normal. No abnormal circulatory autoantibodies were detected; in particular, antglomerular basement membrane antibody was not found. Other investigations are shown in the Table. Macrophages laden with haemosiderin were present in gastric washings. Radioallergosorbent tests against aspergillus, cladosporium, and cow's milk yielded negative results and skin prick tests showed +++ to house dust mite and + to mixed feathers only. IgG precipitins were present against pigeon serum, budgerigar serum, chicken serum, and egg yolk but not against milk proteins. This pattern of precipitins was not that associated with bird fancier's lung; it was, however, typical of the pattern reported in coeliac disease.³ Gliadin antibodies (++++) were present, jejunal biopsy examination showed subtotal villous atrophy, and a gluten free diet was therefore begun.

Over the subsequent 15 months his weight and height velocities have increased modestly and from

Figure Haemoglobin concentrations over three years in a case of idiopathic pulmonary haemosiderosis who required transfusions (T) every three months until he was put on treatment with a gluten free diet. An unsuccessful trial with disodium cromoglycate (DSG) is also shown.

Table Results of investigations before and seven months after starting gluten free diet

	Before starting diet	Seven months after starting diet	Normal range/reference values
Iron ($\mu\text{mol/l}$)	9	21	13-32
Total iron binding capacity ($\mu\text{mol/l}$)	68	46	43-73
Immunoglobulin A (g/l)	3.3	0.81	0.5-2.3
Reticulin antibody	1/80	Negative	Negative
Precipitins to:			
Chicken serum	++	-	
Budgerigar serum	++	-	
Pigeon serum	+	$\pm$	
Egg yolk	++	-	
Forced expiratory volume in one second/forced vital capacity	0.6/0.6	0.79/0.87	0.95/1.25

one month after starting his diet he has required no further transfusions (Figure). Re-investigation after seven months showed a haemoglobin concentration of 12.6 g/dl, with 1% reticulocytes. The results of other investigations are shown in the Table. Chest x ray film showed unchanged widespread micro-nodular shadowing. Jejunal biopsy examination showed mildly stunted villi with some epithelial stratification and an increased epithelial lymphocyte count. Gliadin antibodies remain strongly positive.

Discussion

Reports of an association between pulmonary haemosiderosis and villous atrophy are uncommon, with no previous report of gluten withdrawal leading to an improvement in the pulmonary disorder. Three single case reports document a young adult with pulmonary haemosiderosis who had coeliac disease in childhood,¹ a young adult with coincident pulmonary haemosiderosis and coeliac disease,⁴ and an 18 year old with a probable 11 year history of both conditions.⁵ The only reported series is of seven young adults with pulmonary haemosiderosis who underwent jejunal biopsy examination. Only three, of whom only one had bowel symptoms, had any villous atrophy.⁶ Although treated with a gluten free diet, no follow up data were presented.

Pulmonary haemosiderosis has not been reported previously in association with antibodies against food antigens other than milk proteins. Our patient is unusual in having antibodies against gliadin and egg yolk but not against milk. We believe this to be the first report of clinical features of pulmonary haemorrhage responding dramatically to a gluten free diet.

The response of the chronic blood loss to a gluten free diet suggests this association is not coincidental but is secondary to the gluten enteropathy. Another possibility would be a common primary defect, but the response of both conditions to the diet does not support this. Furthermore, there was no clinical or serological evidence of autoimmune disease.

A reasonable suggestion is that dietary antigen, probably gluten, caused the villous atrophy, leading to antigen penetration and systemic immunisation, resulting in IgG antibodies and subsequent formation of immune complexes. Pulmonary damage would follow their deposition in the lung and demonstration of tissue fixed immune complexes would have supported this theory had a lung biopsy examination been undertaken.

Although the second biopsy examination did not yield entirely normal results, we think the clinical and histological response to a gluten free diet was sufficient provisionally to diagnose gluten enteropathy. In view of the severity of his previous symptoms, gluten challenge may be contraindicated. Whether withdrawal of other dietary antigens would cause further improvement in his growth and villous architecture and the use of steroids for his pulmonary haemosiderosis are speculative.

We suggest that all children with idiopathic pulmonary haemosiderosis should undergo an intestinal biopsy examination. Gliadin antibody and precipitins to other food antigens should be measured at intervals. There is no prima facie reason why milk should contain the only food antigens associated with pulmonary haemosiderosis.

References

- Heiner DC, Sears JW, Kniker WT. Multiple precipitins to cow's milk in chronic respiratory disease. *Am J Dis Child* 1962;103:634-54.
- Dolan J, McGuire S, Sweeny E, Bourke J, Ward OC. Mast cells in pulmonary haemosiderosis. *Arch Dis Child* 1984;59:276-8.
- Hendrick DJ, Faux JA, Anand B, Piris J, Marshall R. Is bird fancier's lung associated with coeliac disease? *Thorax* 1978;33:425-8.
- Lane DJ, Hamilton WS. Idiopathic steatorrhoea and idiopathic pulmonary haemosiderosis. *Br Med J* 1971;ii:89-90.
- Bailey P, Groden BM. Idiopathic pulmonary haemosiderosis, report of two cases and review of the literature. *Postgrad Med J* 1979;55:266-72.
- Wright PH, Menzies IS, Pounder RE, Keeling PWN. Adult idiopathic pulmonary haemosiderosis and coeliac disease. *Q J Med* 1981;197:95-102.

Correspondence to Dr R F Reading, Department of Child Health, Medical School, Fourth Floor, Framlington Place, Newcastle upon Tyne, Tyne and Wear NE2 4BA, England.

Received 7 November 1986