endotoxin damages the cochlea, children with meningococcal disease would be a suitable population to study. Children present predominantly with septicaemia, meningitis, or a mixed picture of septicaemia and meningitis.³ Deafness in survivors of fulminant septicaemia, meningococcaemia without meningitis, and meningitis alone might be commoner in all groups than in a control population. A case-control follow up study of a cohort of children with meningococcal disease would answer this question.

The practical conclusion of such a study might be that clinicians should be ordering routine audiological follow up for patients with meningococcal septicaemia, as well as for those with meningitis.

ALISTAIR THOMSON
OMNIA MARZOUK
Institute of Child Health,
Royal Liverpool Children's
Hospital (Alder Hey),
Eaton Road,
Liverpool L12 2AP

- 1 Tarlow MJ, Comis SD, Osborne MP. Endotoxin induced damage to the cochlea in guinea pigs. *Arch Dis Child* 1991;66:181-4.
- 2 Brandtzaeg P, Kierulf P, Gaustad P, Dobloug J, Mollnes TE, Sirnes K. Systemic meningococcal disease: a model infection to study acute endotoxaemia in man. *Prog Clin Biol Res* 1988;272:263-71.
- 3 Thomson APJ, Sills J, Hart CA. Meningococcal disease in Liverpool 1977-1987: mode of presentation. *Pediatric Reviews and Communications* 1990;5:109-16.

Head injury—how community paediatricians can help

SIR,—Head injury must be the major health issue among children in the United Kingdom, accounting for 15% of deaths in children aged 1-15 years.¹ However, we question the admission rate of 45 per 1000 children per year after head injury quoted by Dr Crouchman in her recent article.² We have recently conducted a retrospective review of all children aged less than 16 years who died or were admitted to hospital after head injury in the Northern region during the eight year period 1979-86.^{1,3,4} During this period head injury was the single major cause of admission to hospital, however, the annual admission rate was only a tenth of that quoted by Dr Crouchman, being 4.9 per 1000 children. In view of the importance of accurate epidemiological data in planning services for head injured children, we believe that it is imperative that this issue is clarified.

We would also like to suggest that community paediatricians should have a leading role in the prevention of head injury and not confine their activities simply to the provision of rehabilitation services. The recent evidence that serious head injury occurs more frequently at particular times of day and in areas of high material deprivation⁵ should surely stimulate community paediatricians to take the lead in initiating appropriately targeted preventive measures.

P M SHARPLES
J A EYRE
Department of Child Health,
The Medical School,
Framlington Place,
Newcastle upon Tyne NE2 4HH

- 1 Sharples PM, Storey A, Aynsley-Green A, Eyre JA. Avoidable factors contributing to death of children with head injury. *BMJ* 1990;300:87-91.
- 2 Crouchman M. Head injury—how community paediatricians can help. *Arch Dis Child* 1990;65:1286-7.
- 3 Sharples PM, Storey A, Aynsley-Green A, Eyre JA. Causes of fatal childhood accidents involving head injury in the Northern region, 1979-86. *BMJ* 1990;301:1193-7.
- 4 Sharples PM, Aynsley-Green A, Eyre JA. The management of children with head injuries in district general hospitals. *Arch Dis Child* 1988;63:570.
- 5 Kraus JF, Rock A, Hemyari P. Brain injuries among infants, children, adolescents and young adults. *Am J Dis Child* 1990;144:684-91.

Dr Crouchman comments:

I must thank Drs Sharples and Eyre for pointing out the typing error in the first sentence of my paper. It should, of course, have read 45 per 10 000 children per year.

I agree that there is an important role for the consultant community paediatrician in the prevention of childhood accidents. My own view is that health education has only a limited place, and that every district needs a multi-agency group with the teeth to influence local housing, street planning, and the provision of safe play space.¹ It would seem very appropriate that consultant community paediatricians should contribute to this group as part of their essential public health function.²

- 1 Crouchman M. Children with head injuries. *BMJ* 1990;301:1289-90.
- 2 Discussion Document from Joint Working Party of the Faculty of Public Health Medicine and the British Paediatric Association. *Working together for tomorrow's children*. London: BPA, 1990:9-11.

Doppler assessment of pulmonary artery pressure in acute phase of hyaline membrane disease

SIR,—I read with interest the paper by Evans and Archer.¹ The study is well described with clear presentation of results.

I would, however, take issue with the definition of right ventricular ejection time (RVET). The authors define RVET as the time interval between the systolic waveform leaving its peak velocity and returning to the baseline. The total RVET is more appropriately defined as the time from the onset of ejection to that of zero flow. The latter definition is that used by Kitabatake *et al* and this is the main reference article by the authors.²

Any error in the measurement of total RVET will be reflected in the ratio time to peak velocity: right ventricular ejection time (TPV: RVET). The ratio as defined by Kitabatake *et al* was shown to have a linear inverse relationship to log₁₀ of the mean pulmonary artery pressure.² Any relationship to pulmonary artery pressure by the ratio TPV: RVET as described by the authors has not been verified.

B G CRAIG
Royal Belfast Hospital for Sick Children,
180-184 Falls Road,
Belfast BT12 6BE

- 1 Evans NJ, Archer LNJ. Doppler assessment of pulmonary artery pressure and extrapulmonary

- shunting in the acute phase of hyaline membrane disease. *Arch Dis Child* 1991;66:6-11.
- 2 Kitabatake A, Inoue M, Asao M, *et al*. Non-invasive evaluation of pulmonary hypertension by a pulsed Doppler technique. *Circulation* 1983;68:302-9.

Drs Evans and Archer comment:

We are grateful to Dr Craig for drawing our attention to an error in the text which we failed to notice at proof reading. The correct definition of right ventricular ejection time (RVET) and the one which we did indeed use is shown in the following phrase which is a corrected version of the text: 'RVET was the time interval between the systolic wave form leaving and returning to the baseline'. This is consistent with Dr Craig's definition.

Photic sneezing

SIR,—Photic sneezing is well documented in the ophthalmological¹ and neurological² literature. Light induced sneezing appears to be more common in male than female, white than black, and in those with a positive family history. Photic sneezing, an uncontrollable paroxysm of sneezing provoked by sudden exposure to intense bright light, is an occupational hazard for the ophthalmologist.

An enhanced photic sneeze reflex has recently been reported in patients with cystinosis.^{3,4} The mechanism of the reflex is obviously complex involving optic, oculomotor, and trigeminal nerves, autonomic pathways, and central brainstem structures. I am a 'sufferer' from photic sneezing, which is most noticeably inconvenient when driving into bright sunlight, and I have two boys who also demonstrate the reflex. I note, with interest, how frequently I have observed the reflex in babies.

I wonder about the evolutionary process behind this bizarre reflex? The reflex appears to be less marked as the child matures. Is this because it serves no useful purpose and is a 'vestigial' reflex?

BARBARA BUCKLEY
Department of Child Health,
Royal Free Hospital,
Pond Street, Hampstead,
London NW3 2QG

- 1 Lewkonia I. An infrequent response to slit lamp examination. *Br J Ophthalmol* 1969;53:493-5.
- 2 Everett HC. Sneezing in response to light. *Neurology* 1964;14:483-90.
- 3 Smith R. Editorial. *Br J Ophthalmol* 1990;74:705.
- 4 Katz B, Melles RB, Wenson MRS, Schneider JA. Photic sneeze reflex in nephropathic cystinosis. *Br J Ophthalmol* 1990;74:706-8.

Duplicate publication

SIR,—I read your recent editorial on duplicate publication with interest and some wry amusement.¹ This is not a new problem and has exercised the minds of editors of general as well as specialist journals.² It is surprising that the editors, all distinguished clinicians, should after an accurate diagnosis spend their efforts discussing control of symptoms with no reference to treating the underlying cause.

Clinicians will continue to submit multiple papers so long as appointment boards contain individuals who believe that a doctor who has published 10 papers must be twice as good as one who has published five. Why a good clinician, who wishes to practice as such,

should also be expected to be a good research worker has never been explained. These are different jobs requiring different skills and interests. Animal physiologists seeking promotion would look askance if rejected on the grounds that they were poor veterinarians—and yet good clinicians are frequently rejected on the grounds they are poor researchers. No wonder they retaliate by producing a large quantity of second rate work that clogs up the journals. It has been argued that the number of publications is one way of distinguishing one candidate from another in a close field. Why not use another irrelevant measure instead: skill at tightrope walking for instance? This would reduce the present volume of substandard publications, save the time of authors, editors and readers, and might even lead to an improvement in the quality of consultant appointments.

In the meantime you have the task of distinguishing between those papers that are written with genuine scientific endeavour and the desire to inform and those written merely to improve a thin looking curriculum vitae. In this you have my sympathies.

IAN G VERBER
Department of Paediatrics,
Memorial Hospital,
Hollyhurst Road,
Darlington DL3 6HX

- 1 Bier DM, Fulginiti VA, Garfunkel JM, *et al.* Duplicate publications and other related problems. *Arch Dis Child* 1990;65:1289–90.
- 2 Lock S. Repetitive publications: a waste that must stop. *BMJ* 1984;288:661–2.

The editors comment:

It is not the role of editors to influence medical appointment boards and we do not mind what motivates people towards research. The need to improve a curriculum vitae may lead to important scientific findings and conversely an altruistic desire to break down the barriers of science may result in substandard work. Our main job is to produce a high quality journal and for this we rely to a large extent on peer review. We publish only one third of the papers submitted to the *Archives of Disease in Childhood* and if we were really clogging the journal up with second rate work our readers would vote with their feet. Instead our circulation is increasing.

SOFTWARE REVIEW

POSSUM. Version 2.5. Melbourne: Murdoch Institute, 1990. CP Export Pty Ltd.

POSSUM, full title, Pictures of Standard Syndromes and Undiagnosed Malformations, is the rather contrived title for a computerised pictorial dysmorphology database from Australia (get it?). For Latin scholars it also translates as 'I am able'. The program requires a microcomputer, a video disc player, and a video monitor so the initial outlay on hardware

is expensive. There is an instruction manual which is reasonably easy to follow, but I found the introductory video tutorial on how to use the system more helpful. Unfortunately, this video section is being dropped from the new edition, due out soon, to make more use of space on the disc.

There are several main components to the system. The illustrated trait directory is a very useful part. Dysmorphology like all other specialties has its own language, and it is helpful for those new to the discipline to see what exactly is meant by a keel shaped forehead, triangular face, prominent ear crus, camptodactyly, etc. The two main components of the system are the search/diagnostic facility and the listing of features of known syndromes. To initiate a search one has to set up a list of traits observed in the patient. The instruction tutorial recommends that between five and 15 features are entered. Personally I would go for as few and as distinct or unusual anomalies as possible and avoid 'soft' common features. Traits can be made optional, major, specific, or excluded in a search and there is the useful facility to check with the pictorial trait directory to ensure the correct term is used for a particular anomaly. After a search, a list of possible diagnoses is presented. For each disorder it is possible to access a brief description, a list of traits, and a few selected references. For an essentially visual specialty by far the best provision is the photographic file of children with specific disorders at different ages. Most of the 'commoner' syndromes are well illustrated but some more rare or recently described syndromes have no illustrations.

The second main use of the system is to review known disorders in the instances where one suspects a diagnosis but needs to confirm this by comparing with similar cases. Each disorder, with a few exceptions, has illustrations of affected children at different ages, many from the files of the Melbourne Children's Hospital. The value of the study of syndrome natural history and the phenotypic changes which occur with age has been recently recognised but most textbooks don't contain such a wealth of photographs. There is one tiny video clip of a patient with Rett's syndrome; more videos of other syndromes especially those with movement abnormalities are promised on the next videodisc update. Like syndrome natural history the value of the study of behavioural phenotypes is being recognised and it is an excellent feature of this system that the importance of movement in syndrome diagnosis has been identified.

The section of this system that I would not imagine using much in clinical practice is that of unknown cases. I believe this section badly needs pruning, as many children are included with rather soft signs, and even if one had a similar case there is so little distinctive to make a confident match. An unknown database, at least for general viewing, should only contain details of children with very distinct phenotypes.

The system is very user friendly and it is easy to get from one section to another. It is comprehensive in the range of disorders covered, but in some instances there is insufficient depth in the description of a syndrome and in the references given. Clearly only if a diagnosis is being considered would there be a need to go to the literature, but I feel that in many instances more information is needed to exclude a diagnosis. I understand a number of inaccuracies are being corrected for the next update. This is needed especially in the

sections on chromosome abnormalities where the karyotype nomenclature is frankly wrong in many instances. The accuracy of individual diagnoses where a diagnosis relies on phenotype alone is good, but for some disorders rather questionable cases are included and for others the 'blackout' required to preserve anonymity also hides diagnostic features.

POSSUM is designed for use by specialist geneticists and paediatricians as a diagnostic aid and for tutorial assistance for specialists in dysmorphology. As a dysmorphologist my main usage would be in verifying diagnoses and in teaching. It is a marvellous teaching aid for those in training and those trained in other disciplines and its ease of use makes self teaching an option. There are a number of rival systems especially the London Dysmorphology Database which has the edge in comprehensiveness of description and references of syndromes, but it currently is not illustrated. I think POSSUM has the edge as a teaching aid.

DIAN DONNAI
Consultant clinical geneticist

BOOK REVIEWS

Fits and Faints. By J B P Stephenson. (Pp 202; £21 hardback.) MacKeith Press, 1990. ISBN 0-632-02811-4.

This book provides information about conditions leading to loss of consciousness in childhood with emphasis on various types of anoxic seizures. As a history of loss of consciousness should not always lead to a diagnosis of epilepsy Stephenson analyses various clinical conditions in childhood and describes a series of case histories drawn from personal practice. A wide range of conditions are considered including common breath holding attacks, loss of consciousness due to vasovagal syncope, unusual cardiac and circulatory conditions, syncope which is self induced by the Valsalva manoeuvre, profound asphyxia, and seizures brought about by anoxia, epilepsy, or psychic disturbances. Anoxic seizures are dealt with extensively with a discussion of various types and an attempt is made to bring together relevant research from other disciplines including circulatory physiology and neurophysiology. Our understanding of anoxic seizures and the methods used in diagnosis is assisted by a useful reference section which draws attention to the author's own contribution over the years.

The first five chapters guide us through the definition of seizures, the extent of the problem in childhood, the importance of the clinical history in the diagnosis of paroxysmal disorders, fits and faints, and the differential diagnosis of fainting episodes. Having established this framework the next seven chapters are about various types of anoxic seizures. There is a detailed account of clinical manifestations which relate to specific types of anoxic seizures, and electroencephalographic and electrocardiographic recordings are illustrated. In a selection of case histories described