LETTERS TO THE EDITOR

Prostacyclin concentrations in haemolytic uraemic syndrome after acute shigellosis

SIR,—We read with interest the report of Alam and colleagues implicating prostacyclin (PGI₂) in the pathogenesis of haemolytic uraemic syndrome after acute shigellosis.¹ However, we were confused by the methodology that they used to establish the PGI₂ concentrations in the plasma from their patients.

The authors state their intention to measure PGI₂ by firstly incubating rabbit aortic rings with their patients' plasma, and then subsequently assaying 6-keto-PGF_{1α} in the incubation fluid, a method described by Moncada *et al.*² We have used a similar technique to investigate the role of PGI₂ in the pathogenesis of meningococcal shock, in which patients' serum was incubated with human umbilical vein endothelial cells.³ Both of these methods have been established to assess endothelial PGI₂ production in vitro in response to exogenous stimuli, and not, as suggested in this paper, to measure the concentration of PGI₂ directly in serum or plasma.

The distinction between direct measurement of PGI₂ and the effect of these patient samples on PGI₂ production by aortic rings is important in determining the significance of the results presented by Alam *et al.* A depression of PGI₂ synthesis in vitro in response to plasmas from patients with haemolytic uraemic syndrome suggests that there is either a deficiency of a factor necessary for PGI₂ synthesis, or that there is a circulating inhibitor. We have recently suggested that the former mechanism is the most likely explanation for the observed decrease in PGI₂ production by endothelial cells when incubated with the sera from

children with meningococcal shock.³ It is interesting that the PGI₂ concentrations reported by Alam and colleagues took a long time to recover. We have observed a defect in the serum of patients with meningococcal shock persisting for days, and sometimes weeks before normal endothelial PGI₂ synthesis can be demonstrated (unpublished results).

The authors conclude that the diminished PGI₂ concentrations reported in their study may be occur as a result of endothelial damage by endotoxin. However, the majority of in vitro, animal and human studies, to date, show that endothelial PGI₂ synthesis is stimulated rather than diminished by endotoxin.^{4,5} This may occur as a direct response to endotoxin or following the elaboration of cytokines such as tumour necrosis factor and interleukin-1.^{5,6} It would therefore seem unlikely that the action of endotoxin alone could adequately explain the depression of PGI₂ concentrations observed in haemolytic uraemic syndrome associated with shigellosis.

ROBERT S HEYDERMAN
NIGEL J KLEIN

Department of Paediatrics,
Queen Elizabeth the Queen Mother Wing,
St Mary's Hospital,
South Wharf Road,
London W2 1NY

- 1 Alam AN, Abdal NM, Wahed MA, *et al.* Prostacyclin concentrations in haemolytic uraemic syndrome after acute shigellosis in children. *Arch Dis Child* 1991;66:1231-4.
- 2 Moncada S, Higgs EA, Vane JR. Human arterial and venous tissues generate prostacyclin (prostaglandin X), a potent inhibitor of platelet aggregation. *Lancet* 1977;i:18-20.
- 3 Heyderman RS, Klein NJ, Shennan GI, Levin M. Deficiency of prostacyclin production in meningococcal shock. *Arch Dis Child* 1991;66:1296-9.
- 4 Ditter H, Mollenhauer J, Matthias FR, Voss R. Prostacyclin formation of the rabbit vessel wall: effects of endotoxin in vivo and in vitro. *Prostaglandins in Clinical Research* 1987: 95-100.
- 5 van der Poll, van Deventer, Büller HR, Sturk A, ten Cate JW. Comparison of the early dynamics of systemic prostacyclin release after administration of tumor necrosis factor and endotoxin

to healthy humans. *J Infect Dis* 1991;164: 599-601.

- 6 Kawakami M, Ishibashi S, Ogawa H, Murase T, Takaku F, Shibata S. Cachectin/TNF as well as interleukin-1 induces prostacyclin synthesis in cultured vascular endothelial cells. *Biochem Biophys Res Commun* 1986;141:482-7.

Diazoxide—an orphan drug?

SIR,—In the article on surgical treatment of hyperinsulinaemic hypoglycaemia in infancy and childhood the authors state that diazoxide, which inhibits glucose stimulated insulin secretion, is still the mainstay of medical management.¹ However, they fail to mention that, in its oral form, this drug is no longer readily available as the manufacture of this therapeutically desirable but commercially non-viable product has ceased and, thus, satisfies the definition of an orphan drug.²

This problem applies to other essential drugs and compounds used in paediatric practice, particularly for the treatment of inborn errors of metabolism. There is a clear need for the drug industry to be supported (by government subsidies if necessary) to maintain production of such essential drugs. Patients need orphan drugs and this must be the overriding consideration. Consequently, paediatricians, through the British Paediatric Association and the Royal Colleges should encourage the Department of Health to provide such direction to the drug industry, perhaps through the establishment of an orphan drug regulatory body under the auspices of the Medicine Control Agency.

D C DAVIDSON
A J NUNN

Royal Liverpool Children's NHS Trust
Alder Hey,
Eaton Road,
Liverpool L12 2AP

- 1 Spitz L, Bhargava RK, Grant DB, Leonard JV. Surgical treatment of hyperinsulinaemic hypoglycaemia in infancy and childhood. *Arch Dis Child* 1992;67:201-5.
- 2 Leonard JL. Orphan diseases and orphan drugs: A report based on the panel discussion. *Proceedings of the RTMDC Conference*. University of Keele, 14 September, 1991:5-8.