- 2 Wheeler MC, Roe MH, Kaplan EL, Schlievert PM, Todd JK. Outbreak of group A streptococcus septicemia in children. Clinical, epidemiologic, and microbiologic correlates. *JAMA* 1991; **266**: 533-7.
- 3 Centers for Disease Control and Prevention. Nursing home outbreaks of invasive group A streptococcal infections - Illinois, Kansas, North Carolina, and Texas. *MMWR Morb Mortal Wkly Rep* 1990; **39**: 577-9.
- 4 Centers for Disease Control and Prevention. Group A beta-hemolytic streptococcal bacteremia - Colorado, 1989. *MMWR Morb Mortal Wkly Rep* 1990; **39**: 3-6, 11.
- 5 Stromberg A, Romanus V, Burman LG. Outbreak of group A streptococcal bacteremia in Sweden: an epidemiological and clinical study. *J Infect Dis* 1991; **164**: 595-8.
- 6 Stevens DL. Invasive group A streptococcus infections. *Clin Infect Dis* 1992; **14**: 2-13.
- 7 Benjamin EM, Gershman M, Goldberg BW. Community-acquired invasive group A beta-hemolytic streptococcal infections in Zuni Indians. *Arch Intern Med* 1992; **152**: 1881-4.
- 8 Katz AR, Morens DM. Severe streptococcal infections in historical perspective. *Clin Infect Dis* 1992; **12**: 298-307.
- 9 Bisno AL. Group A streptococcal infections and acute rheumatic fever. *N Engl J Med* 1991; **325**: 783-93.
- 10 Schwartz B, Facklam R, Breiman RF. Changing epidemiology of group A streptococcal infections in the USA. *Lancet* 1990; **336**: 1167-71.
- 11 Smith TD, Wilkinson V, Kaplan EL. Group A streptococcus-associated respiratory tract infections in a day-care center. *Pediatrics* 1989 **83**: 380-4.
- 12 Falck G, Kjellander J. Outbreak of group A streptococcal infection in a day-care center. *Pediatr Infect Dis J* 1992; **11**: 914-9.
- 13 American Academy of Pediatrics. In: Peter G, ed. *Report of the committee on infectious disease*. 22nd Ed. Chicago, Illinois: AAP, 1991.
- 14 National Commission on Children. *Beyond rhetoric: a new American agenda for children and families*. Washington DC: National Commission on Children, 1991.
- 15 Chaudhary S, Bilinsky SA, Hennessy JL, et al. Penicillin V and rifampin for treatment of group A streptococcal pharyngitis: a randomized trial of 10 days penicillin vs 10 days penicillin with rifampin during the final 4 days of therapy. *J Pediatr* 1985; **106**: 481-6.
- 16 Wilson E, Zimmerman RA, Moody MD. Value of T-agglutination typing of group A streptococci in epidemiologic investigations. *Health and Laboratory Science* 1968; **5**: 199-207.
- 17 Moody MD, Padula J, Lizana D, Hall CT. Epidemiologic characterization of group A streptococci by T-agglutination and M-precipitation tests in the public health laboratory. *Health and Laboratory Science* 1965; **2**: 149-62.
- 18 Seppala H, Vuopio-Varkila J, Osterblad M, Jauhola Rummukainen M, Holm SE, Huovinen P. Evaluation of methods for epidemiologic typing of group A streptococci. *J Infect Dis* 1994; **169**: 519-25.
- 19 Fleming DW, Leibenhaut MH, Albanes D, et al. Secondary haemophilus influenzae type b in day-care facilities. Risk factors and prevention. *JAMA* 1985; **254**: 509-14.
- 20 Gray GC, Escamilla J, Hyams KC, et al. Hyperendemic streptococcus pyogenes infections despite prophylaxis with penicillin G benzathine. *N Engl J Med* 1991; **325**: 92-7.
- 21 Li KI, Dashefsky B, Wald ER. Haemophilus influenzae type b colonization in household contacts for infected and colonized children enrolled in day care. *Paediatrics* 1986; **78**: 15-20.
- 22 Reed BD, Huck W, Zazove P. Treatment of beta-hemolytic streptococcal pharyngitis with cefaclor or penicillin. Efficacy and interaction with beta-lactamase-producing organisms in the pharynx. *J Fam Pract* 1991; **32**: 138-44.

Commentary

Streptococcus pyogenes, the group A β -haemolytic streptococcus, remains a troublesome micro-organism. Good hygiene, antibiotics, and other attributes of a high standard of living have lessened the frequency of severe infection, but the bacterium has kept its virulence factors. Given a chance to spread among close contacts and to invade through

broken skin, outbreaks with severe or fatal cases are possible. The paper by Engelgau and colleagues reminds us of this and shows how investigations should be done. Three aspects merit a comment: the organism, the route of infection, and the outbreak management.

The M type protein of *S pyogenes* is an important determinant of its pathogenesis and epidemiology. This is clearest in glomerulonephritis, which is the sequel of infections by a few M types (for example, M types 12, 49, 55, and 60). T type 1 is equivalent to M type 1 in this incident, but T typing is less specific for most other types. M type 1 is the type most often reported in overwhelming infection, but it is a common strain and typing is most likely to be done for the bacteraemic and other severe infections. The M type is associated with specific protective immunity, which should not be confused with the antibodies against exotoxins used in diagnostic tests, of which streptolysin O is the best known. This type specific immunity partly explains why streptococcal diseases fluctuate under the influence of herd immunity to common strains.

Chickenpox is one of several skin conditions that may become secondarily infected by *S pyogenes*. Burns, insect bites, and minor injuries are commoner portals of streptococcal skin infection, but seem to have less risk of bacteraemia with consequent shock, osteomyelitis, or arthritis. Children are the population in whom *S pyogenes* circulate most as they have least type specific immunity and closest contacts with other infected individuals.

Streptococcal infection is endemic, and virulent bacteria in susceptible children will cause the occasional death. Should we investigate the contacts of people who have severe streptococcal disease, when little is done in the numerous mild cases? Is this not shutting the stable door after the horse has bolted? The death of a child creates a feeling that 'something must be done'. The paper by Engelgau *et al* shows that it is difficult to test everyone in a large group, and the time taken for laboratory tests diminishes the benefit of prophylactic treatment. The finding of lower carriage rates in children who had had antibiotics recently, suggests that good access to primary care is one reason why streptococcal infections have become less threatening in modern times.

R T MAYON-WHITE
Public Health Department,
Manor House,
John Radcliffe Hospital,
Oxford OX3 9DZ