- 7 Abu-Arafeh IA, Russell G. Current controversies – abdominal migraine. *Cephalalgia* 1993; 13: 138–9.
- 8 Mortimer MJ, Good PA, Marsters JB, Addy DP. Visual evoked responses in children with migraine: a diagnostic test. *Lancet* 1990; 335: 75–7.
- 9 Mortimer MJ, Good PA. The VER as a diagnostic marker for childhood abdominal migraine. *Headache* 1990; 30: 642–5.
- 10 Symon DNK. Is there a place for 'abdominal migraine' as a separate entity in the IHS classification? Yes! *Cephalalgia* 1992; 12: 345–6.
- 11 Hockaday JM. Is there a place for 'abdominal migraine' as a separate entity in the IHS classification? No! *Cephalalgia* 1992; 12: 346–8.
- 12 Symon DNK. Pizotifen. In: Gallai V, Guidetti V, eds. *Juvenile headache. Etiopathogenesis, clinical diagnosis and therapy*. Amsterdam: Elsevier Science, 1991: 405–8.
- 13 Gillies D, Sills M, Forsythe I. Pizotifen (Sanomigran) in childhood migraine. *Eur Neurol* 1986; 25: 32–5.
- 14 Symon DNK, Russell G. The general paediatrician's view of migraine – a review of 250 cases. In: Lanzi G, Balottin U, Cernibori A, eds. *Headaches in children and adolescents*. Amsterdam: Elsevier Science, 1989: 61–6.
- 15 Symon DNK, Russell G. Continued benefit after stopping pizotifen therapy in childhood migraine. *Cephalalgia* 1989; 9 (suppl 10): 422–3.
- 16 Peet KMS, Gray JA. Pizotifen (Sanomigran) in the prophylaxis of migraine headache: a long-term follow-up study. *Cephalalgia* 1989; 9 (suppl 10): 420–1.

Commentary

I agree with the authors that this is an important study. It is always comforting to learn that a widely used treatment has a rational basis.

Until recently, the migrainous nature of recurrent paroxysmal abdominal pain with facial pallor and a family history of migrainous headache has been contentious. The recent demonstration of the same fast wave abnormality in the visually evoked response of

children with paroxysmal abdominal symptoms,¹ as in those with migrainous headache, has helped to settle the debate. The present observation that the serotonin receptor stimulator, pizotifen, a proved treatment for cranial migraine, is equally effective in abdominal migraine is consistent with such a link.

The study also has important therapeutic implications. It is important to remember that prophylaxis is not appropriate in those children with mild, brief, or infrequent episodes. Pizotifen is also expensive.

Despite several convincing demonstrations to the contrary, the belief that recurrent abdominal pain without an organic basis is almost invariably psychogenic, dies hard. Hopefully, this paper, with others implicating defects in gastrointestinal motility for example,² will be further nails in the coffin of that particular hypothesis.

I W BOOTH
Institute of Child Health,
Nuffield Building,
Francis Road,
Birmingham B16 8ET

- 1 Mortimer MJ, Good PA, Marsters JB, Addy DP. Visual evoked responses in children with migraine: a diagnostic test. *Lancet* 1990; 335: 75–7.
- 2 Pineiro-Carrero VM, Andres JM, Davis RH, Mathias JR. Abnormal gastroduodenal motility in children and adolescents with recurrent functional abdominal pain. *J Pediatr* 1988; 113: 820–5.