CURRENT TOPIC

Mineral accretion in growing bones – a framework for the future?

N J Shaw, N J Bishop

The increasing sophistication and versatility of the techniques available for measuring mineral mass, coupled with the perception that adult degenerative bone disease may have its origins in early life, has led to increasing interest in measuring bone mineral accretion from birth through to adolescence.

Assessment of bone mineral

The techniques that have been used to study the accretion of bone mineral during childhood include single and dual photon absorptiometry, quantitative computed tomography, and dual energy *x* ray absorptiometry (DXA).

Single photon absorptiometry, which has been used for over 20 years, uses an iodine-125 source to produce a beam of collimated photons that pass to a photomultiplier detector. The forearm is positioned perpendicularly to the scan path. It produces a measurement of bone mineral content (g/cm) and assumes that the site of measurement is a cylinder of constant width. This technique has been used most effectively in studies of preterm infants. It has an accuracy and precision of 2–5%, a radiation dose of 5 mRem, and a scan time of 10 to 15 minutes. Its limitation is that it can only examine bones in the peripheral skeleton. As these are mainly cortical bones in children, which has a low turnover, this technique is less sensitive than modern methods that can examine the predominantly trabecular, high turnover, bone of the spine and hips.

Dual photon absorptiometry by using two separate energy levels is able to examine these areas and produces a measurement of bone mineral density (g/cm²) – a reflection of the attenuation of the beam over the area scanned. A scan takes 15–20 minutes and has a radiation dose of 5 mRem. It has an accuracy of 4–10% and precision of 2–4% but suffers the disadvantage in that the radioisotope decays with time thus reducing its precision.

Quantitative computed tomography is the only technique that measures true bone density (mg/cm³). It has the ability to separately measure trabecular and cortical bone within the spine and is not influenced by vertebral size. A scan takes 10–20 minutes, has an accuracy of 10–15%, and a precision of 2–4%. Its disadvantage, however, is a significant radiation dose of 100 mRem that

limits its use in children and in longitudinal studies.

DXA is currently the most favoured technique. A beam of collimated *x* rays are transmitted from a source that provides alternating pulses of 70 and 140 kV through a rotating calibration disk composed of bone and soft tissue equivalent materials and are then measured by a detector located above the subject. It produces a measurement of bone mineral density (g/cm²) by correcting the bone mineral content for the projected area of bone. A scan takes eight minutes on the first generation machines and only 45 seconds on the new generation fan beam machines. It has an accuracy and precision of 1–2% and a radiation dose of 3–5 mRem. It also has the ability to provide estimates of total lean body and fat mass in all but small infants.

A promising additional technique is ultrasound of the calcaneum with which there have been some preliminary studies in children.¹

The estimates of bone mineral density by dual photon absorptiometry and DXA need careful interpretation. It is important to appreciate that in growing children the values obtained are influenced by bone size and therefore body size. This is because it produces a measurement of bone area and does not correct for differences in bone thickness. Thus bone mineral density will be overestimated in large bones and underestimated in small bones. In an attempt to correct for this, a model has been recently proposed that expresses bone mineral adjusted for bone volume,² known as the bone mineral apparent density (g/cm³). An alternative approach when comparisons are made with normative data, is to match individuals for body size rather than age. Similarly as puberty has such a major influence on bone mineral density it is important to match for pubertal status when studying older children. This is of particular relevance in studies of children with chronic disease who may be small, underweight, and have delayed puberty. It is also important to ensure that normative data is ethnic specific as black people have a higher bone mineral density than white people. The reference data provided by manufacturers of DXA machines is based on studies of American children and is inappropriate for studying British children because of differences in diet, body size, and maturation. It is important therefore to develop reference data

Department of Growth and Endocrinology, Institute of Child Health, The Children's Hospital, Birmingham B16 8ET
N J Shaw

Department of Paediatrics, University of Cambridge, Addenbrooke's Hospital, Cambridge
N J Bishop

Correspondence to:
Dr Shaw.

for British children that take account of body size and pubertal status.

Although there is a considerable body of evidence to support the assumption that a reduction in bone mineral density is associated with an increased risk of fractures, it is worth noting that a measurement of bone density may not always correlate with bone strength. As an example of this, children with osteopetrosis have a large amount of mineral in their bones but they fracture easily.

Bone mineral accretion in health and disease

Studies of bone mineral content and density in childhood cover three main groups. (1) Preterm infants receiving different diets during and after their initial period of hospitalisation. (2) Late childhood and adolescence when the focus is on 'peak bone mass' achieved in relation to diet, exercise, and pubertal stage. (3) Children with chronic diseases where there is increased risk of reduced bone mineralisation.

(1) PRETERM INFANTS

The term infant spends the last trimester of pregnancy accreting large stores of mineral, and receives sufficient mineral substrate after delivery for the needs of normal growth and development to be satisfied. By contrast, the infant born preterm is growing at a faster rate, has low stores of many nutrients, is frequently in a catabolic state after delivery, and receives inadequate amounts of bone mineral (and possibly matrix) substrate.³ The preterm infant is therefore at risk of metabolic bone disease because of its rate of growth outstripping the available supplies of substrate.

Bone disease is seen most commonly in the most immature infants who have either received prolonged periods of parenteral feeding or who have been fed solely on human milk from birth. There is widespread agreement that deficiency of mineral substrate is the principal aetiological factor. Typically human milk contains 15–18 mg/100 ml of phosphorus and 32–35 mg/100 ml of calcium. Standard preterm formula contains 35 mg/100 ml of phosphorus and 70 mg/100 ml of calcium; some preterm formulas available in the US contain even higher amounts.

A number of studies have compared bone mineral content in preterm infants receiving different diets or receiving the same base diet with variable levels of calcium and phosphorus supplementation. Although bone mineral content is generally higher in those receiving the mineral enriched formulations, it is usually still less than that of newly born infants of the same postconceptional age.^{4,5} There is evidence from several studies, however, that bone mineral content increases rapidly after discharge from hospital,^{6–8} with no difference between those receiving different early diets at two years.⁹ Longer term studies of the effects of early diet on later bone mineralisation are still awaited. Decreased bone mineral content has also been observed in infants of diabetic

mothers¹⁰ and infants with intrauterine growth retardation.¹¹

(2) PEAK BONE MASS

The importance of studies of bone mineral accretion in children relates to the awareness that one of the main determinants of fracture risk in adulthood is the peak bone mass.¹² Until recently it was felt that peak bone mass was not achieved until the third decade of life. However, new longitudinal studies of bone mass acquisition during adolescence indicate that the majority of peak bone mass is acquired at this time. Several cross sectional studies have shown a progressive rise in bone mineral density with age with the most important influences being body weight and pubertal status.^{13,14} A recent study of bone mineral content in healthy Cambridge schoolchildren confirmed the correlation with body size with prediction equations and nomograms for bone mineral content being generated on the basis of the children's height and weight.¹⁵ A longitudinal survey from Geneva highlighted the importance of puberty producing a rapid increase in bone mineral density with no increment occurring beyond the age of 16 years in girls and 18 years in boys.¹⁶ Furthermore the bone mineral density seen at age 18 was very similar to that seen in adults between the ages of 20 and 35 years.

Why does this matter? The relevance is that events in childhood may have important implications for adults. Chronic conditions acquired in childhood may affect bone mineralisation producing a reduction in peak bone mass and thus increasing the risk of osteoporotic fractures in adulthood. Conversely if it were possible to produce positive increments in peak bone mass by interventions undertaken in childhood, this could potentially reduce the likelihood of osteoporosis with considerable implications for health care services for adults. Although it is felt that 80% of peak bone mass is genetically determined this still leaves 20% determined by environmental influences and thus potentially open to manipulation. Other important influences other than body size and pubertal status are dietary calcium intake and weightbearing exercise.

(3) CHRONIC CHILDHOOD DISEASE

Several studies have demonstrated the presence of osteopenia in many of the chronic childhood conditions.^{17–24} Although a variety of factors are of importance, the effects of poor nutrition and delayed puberty appear to be the strongest determinants. The importance of puberty in influencing peak bone mass has again been highlighted by a recent study of adult men with a history of untreated delayed puberty.²⁵ Both radial and spinal bone mineral densities were significantly reduced compared with controls; two thirds had values more than 1 SD below the mean. A reduction of bone mineral density of this extent probably represents a 50 to 100% increase in the incidence of fracture.²⁶ Furthermore, the timing of puberty

appeared to be the most significant determinant of bone mineral density.

The implication of this study is that puberty must occur during a critical period of bone development in order to achieve peak bone mass. Delayed puberty is a feature in many children with chronic disease and although most will eventually progress through puberty they could have a reduced bone mass in adulthood with an increased risk of fractures.

What possible therapeutic or preventive options are available that may influence bone mineralisation in chronic childhood conditions? Obviously, aggressive nutritional intervention is necessary in those conditions where malnutrition is a major feature. Calcium supplements that have been shown to increase bone density in normal prepubertal children are a potential option.²⁷ The initiation of puberty at an age appropriate point, using oestrogens or androgens is another consideration, but carries the risk of advancing the bone age faster than height age, with implications for final height.

Newer corticosteroids with apparent bone sparing effects²⁸ are a possible alternative in children on maintenance treatment but there has been little research so far undertaken in this group. The availability of bisphosphonates, which act as specific inhibitors of bone resorption, offer a potential therapeutic option in children who are having fractures as a consequence of their osteopenia. They have produced encouraging results in postmenopausal osteoporosis²⁹ but there is very limited experience of their use in children and some concerns about their effect in a growing skeleton.

Future research

The major question that now needs to be addressed is whether events in childhood predispose to an increased fracture risk in adults. Evidence is accumulating of a genetic predisposition towards osteoporosis with allelic variation in the gene encoding for the vitamin D receptor being linked to differences in bone density.³⁰ The significance of this if confirmed is that early identification of the 'at risk' population by a genetic test would permit targeted interventions in the areas already known to positively influence bone mass (exercise and nutrition). In this context, further longitudinal studies of mineral accretion throughout childhood and into adult life are now urgently required.

- 1 Mughal MZ, Utretch G, Specker BL, Langton CM. Comparison between broadband ultrasound attenuation of calcaneum and bone mineral density measurement by dual energy X-ray absorptiometry. *Proceedings of the 66th annual meeting of the British Paediatric Association*. University of Warwick 12–15 April 1994. London: BPA, 1994: abst G146.
- 2 Carter DR, Bouxsein ML, Marcus R. New approaches for interpreting projected bone densitometry data. *J Bone Miner Res* 1992; 7: 137–45.
- 3 Bishop NJ. Bone disease in preterm infants. *Arch Dis Child* 1989; 64: 1403–9.

- 4 Chan GM, Roberts CC, Pollard D, Jackson R. Growth and bone mineralization of normal breast-fed infants and the effects of lactation on maternal bone mineral status. *Am J Clin Nutr* 1982; 36: 438–43.
- 5 Chan GM, Mileur L, Hansen JW. Effects of increased calcium and phosphorus formulas and human milk on bone mineralization in preterm infants. *J Pediatr Gastroenterol Nutr* 1986; 5: 444–9.
- 6 Congdon PJ, Horsman A, Ryan SW, Truscott JG, Durward H. Spontaneous resolution of bone mineral depletion in preterm infants. *Arch Dis Child* 1990; 65: 1038–42.
- 7 Bishop NJ, King FJ, Lucas A. Increased bone mineral content of preterm infants fed with a nutrient enriched formula after discharge from hospital. *Arch Dis Child* 1993; 68: 573–8.
- 8 Pittard WB III, Geddes KM, Sutherland SE, Miller MC, Hollis BW. Longitudinal changes in the bone mineral content of term and premature infants. *Am J Dis Child* 1990; 144: 36–40.
- 9 Schanler RJ, Burns PA, Abrams SA, Garza C. Bone mineralisation outcomes in human milk-fed preterm infants. *Pediatr Res* 1992; 31: 583–6.
- 10 Mimouni F, Steichen JJ, Tsang RC, Hertzberg V, Miodovnik M. Decreased bone mineral content in infants of diabetic mothers. *Am J Perinatol* 1988; 5: 334–8.
- 11 Minton SD, Steichen JJ, Tsang RC. Decreased bone mineral content in small for gestational age infants compared with appropriate for gestational age infants: normal serum 25-hydroxyvitamin D and decreasing parathyroid hormone. *Pediatrics* 1983; 71: 383–8.
- 12 Hansen MA, Overgaard K, Riis BJ, Christiansen C. Role of peak bone mass and bone loss in postmenopausal osteoporosis: 12 year study. *BMJ* 1991; 303: 961–4.
- 13 Gialstre C, Brailon P, David L, Cochat P, Meunier PJ, Delmas PD. Measurement of bone mineral content of the lumbar spine by dual energy X-ray absorptiometry in normal children: correlations with growth parameters. *J Clin Endocrinol Metab* 1990; 70: 1330–3.
- 14 Southard RN, Morris JD, Mahan JD, et al. Bone mass in healthy children: measurement with quantitative DXA. *Radiology* 1991; 179: 735–8.
- 15 DePriester JA, Cole TJ, Bishop NJ. Bone growth and mineralisation in children aged 4 to 10 years. *Bone Miner* 1991; 12: 57–65.
- 16 Thientz G, Buchs B, Rizzoli R, et al. Longitudinal monitoring of bone mass accumulation in healthy adolescents: evidence for a marked reduction after 16 years of age at the levels of lumbar spine and femoral neck in female subjects. *J Clin Endocrinol Metab* 1992; 75: 1060–5.
- 17 Gibbens DT, Gilsanz V, Boechat MI, Dufer D, Carlson ME, Wang CI. Osteoporosis in cystic fibrosis. *J Pediatr* 1988; 113: 295–300.
- 18 Saggese G, Bertelloni S, Baroncelli GI, Federico G, Calisti L, Fusaro C. Bone demineralisation and impaired mineral metabolism in insulin-dependant diabetes mellitus. *Helvetica Paediatrica Acta* 1988; 43: 405–14.
- 19 Shaw NJ, White CP, Fraser WD, Rosenbloom LR. Osteopenia in cerebral palsy. *Arch Dis Child* 1994; 71: 235–8.
- 20 Gilsanz V, Carlson ME, Roe TF, Ortega JA. Osteoporosis after cranial irradiation for acute lymphoblastic leukaemia. *J Pediatr* 1990; 117: 238–44.
- 21 Chesney RW, Mazess RB, Rose PG, Jax DK. Bone mineral status measured by direct photon absorptiometry in childhood renal disease. *Pediatrics* 1977; 60: 864–72.
- 22 Hyer SL, Rodin DA, Tobias JH, Leiper A, Nussey SS. Growth hormone deficiency during puberty reduces adult bone mineral density. *Arch Dis Child* 1992; 67: 1472–4.
- 23 Bachrach LK, Guido D, Katzman D, Litt IF, Marcus R. Decreased bone density in adolescent girls with anorexia nervosa. *Pediatrics* 1990; 86: 440–7.
- 24 Chesney RW, Mazess RB, Rose P, Jax DK. The effect of prednisone on growth and bone mineral content in childhood glomerular disease. *Am J Dis Child* 1978; 132: 768–72.
- 25 Finkelstein JS, Neer RM, Biller BMK, Crawford JD, Klibanski A. Osteopenia in men with a history of delayed puberty. *N Engl J Med* 1992; 326: 600–4.
- 26 Gardsell P, Johnell O, Nilsson BE. The predictive value of forearm bone mineral content measurements in men. *Bone* 1990; 11: 229–32.
- 27 Johnston CC, Miller JZ, Slemenda CW, et al. Calcium supplementation and increases in bone mineral density in children. *N Engl J Med* 1992; 327: 82–7.
- 28 Loftus J, Allen R, Hesp R, et al. Randomised, double-blind trial of deflazacort versus prednisone in juvenile chronic (or rheumatoid) arthritis: a relatively bone-sparing effect of deflazacort. *Pediatrics* 1991; 88: 428–36.
- 29 Storm T, Thamsborg G, Steiniche T, Genant HK, Sorenson OH. Effect of intermittent cyclical etidronate therapy on bone mass and fracture rate in women with postmenopausal osteoporosis. *N Engl J Med* 1990; 322: 1265–71.
- 30 Morrison NA, Cheng Qi J, Tokita A, et al. Prediction of bone density from vitamin D receptor alleles. *Nature* 1994; 367: 284–7.