report of cost-benefit analysis of inhaled steroids in the prophylaxis of moderate to severe childhood asthma.

The patients studied were from the higher social classes and they were able to afford private medical attention. The excellent compliance rate of the patients in this study probably reflects the higher levels of education and intelligence of this particular social class among the general population. The very high literacy rate of around 88% that is found in Sri Lanka is likely to be another factor that contributed to this level of compliance. This study was not originally designed to look at the undesirable effects of inhaled steroids and as such, no comment could be made on this aspect of treatment. However, the mean starting doses and the mean maintenance doses of both beclomethasone dipropionate and budesonide were well below the levels at which side effects are known to occur.^{22 23}

Conclusion

This study clearly demonstrates the clinically efficacy and cost effectiveness of inhaled steroid prophylaxis in moderate to severe childhood asthma. The findings would be of significance not only to developing countries with their own financial problems but also to the more developed nations that are showing a reawakening of interest in medical audit and cost-benefit ratios of different treatment methods.

The author wishes to thank with gratitude the assistance provided by Dr D R Weerasekera, senior lecturer, Department of Statistics and Computer Science, University of Colombo, Sri Lanka, in the statistical analysis of this paper. Detailed calculations are available from the author.

- Williams HB, McNicol KN. Prevalence, natural history and relationship of wheezy bronchitis and asthma in childhood. An epidemiological study. *BMJ* 1969; iv: 321-5.
- Nocon A. Social and emotional impact of childhood asthma. *Arch Dis Child* 1991; 66: 458-60.
- Fletcher HJ, Ibrahim SA, Speight N. Survey of asthma deaths in the Northern region, 1970-85. *Arch Dis Child* 1990; 65: 163-7.
- Shaw RA, Crane J, O'Donnell TV, Porteus LE, Coleman ED. Increasing asthma prevalence in a rural New Zealand adolescent population: 1975-89. *Arch Dis Child* 1990; 65: 1319-23.
- Barnes PJ. Mechanisms of asthma. *Medicine International* 1991; 89: 3694-702.
- Warner JO, Gotz M, Landau LI, et al. Management of asthma: a consensus statement. *Arch Dis Child* 1989; 64: 1065-79.
- Godfrey S, Balfour-Lynn L, Tooley M. A three to five year follow up of the use of the aerosol steroid beclomethasone dipropionate in childhood asthma. *J Allergy Clin Immunol* 1978; 62: 335-9.
- Katz RM, Rachelefsky GS, Siegel SC, Spector SL, Rohr AS. Twice-daily beclomethasone dipropionate in the treatment of childhood asthma. *J Asthma* 1986; 23: 1-7.
- Central Bank of Sri Lanka. *Sri Lanka socio-economic data 1993*. Colombo, 1993.
- World Bank. *World development report 1993 - investing in health*. Oxford: Oxford University Press, 1993.
- Russell G. Asthma and growth. *Arch Dis Child* 1993; 69: 695-8.
- Balfour-Lynn L, Tooley M, Godfrey S. Relationship of exercise induced asthma to clinical asthma in childhood. *Arch Dis Child* 1980; 56: 450-4.
- Anderson HR, Bailey PA, Cooper JS, Palmer JC, West S. Morbidity and school absence caused by asthma and wheezing illness. *Arch Dis Child* 1983; 58: 777-84.
- Seddon PC, Heaf DP. Long term outcome of ventilated asthmatics. *Arch Dis Child* 1990; 65: 1324-8.
- Barnes PJ. New concepts in the pathogenesis of bronchial hyperresponsiveness and asthma. *J Allergy Clin Immunol* 1989; 83: 1013-6.
- Godfrey S, König P. Treatment of childhood asthma for 13 months and longer with beclomethasone dipropionate aerosol. *Arch Dis Child* 1974; 49: 591-6.
- Kerrabijn KF. Beclomethasone dipropionate in long term treatment of asthma in children. *J Pediatr* 1976; 89: 821-6.

- Russell G, Jones SP. Selection of skin tests in childhood asthma. *British Journal of Diseases of the Chest* 1976; 70: 104-6.
- Hall J, Mooney G. What every doctor should know about economics - part 1. The benefits of costing. *Med J Aust* 1990; 152: 29-31.
- Huskinson EC. Assessment for clinical trials. *Clinics in Rheumatic Diseases* 1976; 2: 37-49.
- de Silva AH. Cost analysis of patient care at the Lady Ridgeway Hospital. *Proceedings of the first scientific sessions of the College of Medical Administrators of Sri Lanka*. Colombo, December 1993.
- Ninan TK, Russell G. Asthma, inhaled corticosteroid treatment and growth. *Arch Dis Child* 1992; 67: 703-5.
- Priftis K, Milner AD, Conway E, Honour JW. Adrenal function in asthma. *Arch Dis Child* 1990; 65: 838-40.
- Mantel N. Biased inferences in tests for average partial associations. *American Statistician* 1980; 34: 190-1.
- Schlesselman JJ. *Case control studies, design, conduct analysis*. Oxford: Oxford University Press, 1982: 219.

Commentary

In any society the connection between value and costs must be made with regard to all forms of health care. To determine whether the value of a health activity is worth its costs should be made not only by professionals but also by the people who will actually receive the value (experience the benefits and harms) and pay the costs.¹ Perera's paper examines the efficacy and cost effectiveness of inhaled steroids in asthma in Sri Lanka, a developing country with severe resource constraints. Relatively high cost treatment may be justified in this setting if the cost-utility value to the patient is reduced by the treatment. Perera has shown this effect in the population of patients he studied, but one cannot conclude that inhaled steroid treatment for prophylaxis of asthma is a cost effective treatment in all developing country settings without considering some of the limitations of the study.

The sample studied was small (n=86) even though the results showed strong cost-utility improvements. Of potentially greater importance is the unrepresentativeness of the patients studied. As noted at the end of the paper, the subjects were mostly from the upper income groups (being self selecting because of fee paying). By the same token, the application of the results to the public sector would apply disproportionately to members of lower income groups. The potential significance of differences in education/literacy is acknowledged. It seems possible, however, that the effectiveness of treatment and the extent of the benefits could be affected quite substantially by other income related factors such as nutrition, home environment, and morbidity from other causes. This may also limit the extent to which the results could be applied to other developing countries.

Similarly, the fact that the subjects were users of the private health sector may well mean that their previous treatment is unrepresentative of public sector patients. All but two of the subjects were on oral β agonists, all but 14 on oral theophyllines, and all but five had had repeated courses of intravenous and oral steroids. This suggests that at least 75% had all three forms of treatment during the control period. Essentially, the analysis is a comparison of the proposed treatment with this alternative regimen, which is almost certainly different from the pattern of treatment in the public sector in Sri Lanka or elsewhere in

the developing world. The treatment protocol used seems more intensive than might normally be expected in a public sector hospital in a developing country (two days initial hospitalisation, and constant supervision thereafter). This might again affect the application of the results to the public sector.

In any cost analysis it is important to specify which costs have been measured. At the very least the costs to the health sector should be stated separately from the costs to the patients – this is a critical issue in resource constrained public health services. It should also be made clear whether the costs used are the fees charged in the private sector, the cost to the private hospital of providing the service, or estimated unit costs in the public sector. It is also helpful if there is some breakdown of costs before and after treatment. The costs of treatment will decline substantially over time as training

occurs only at the beginning of the process, the need for supervision should decline over time, and some patients are able to stop the treatment. Not all these points have been considered in Perera's study in which the reported cost per patient seems remarkably low at about £6.

Overall this is a valuable study of relevance both to clinicians and to health planners in developing countries where cost effectiveness must be a crucial factor in any planning decision.²

ANTHONY COSTELLO
DAVID WOODWARD
*Centre for International Child Health,
Institute of Child Health,
30 Guilford Street,
London WC1N 1EH*

1 Eddy DM. Connecting value and costs. *JAMA* 1990; **264**: 1737–9.

2 World Bank. *World development report 1993 – investing in health*. Oxford: Oxford University Press, 1993.