CURRENT TOPIC

Positron emission tomography and the central nervous system

R O Robinson, C D Ferrie, M Capra, M N Maisey

Sixty five years ago artificial radioactivity was discovered.¹ Radioactive tracers such as ^{11}C , ^{15}O , and ^{18}F , which decay by positron emission, were used to derive an image of the distribution in organs during the the mid 1970s. This became clinically important by the 1980s. The physics of position emission tomography (PET) and the logistics of its application are complex. Imaging is based on coincidence detection of annihilation radiation. The unstable radionuclides are neutron deficient and decay by emission of a positively charged electron (positron). The positron travels a few millimetres in the tissue before combining with an electron. Mass is lost and energy is emitted in the form of two gamma rays of energy 511 KeV, which travel in almost opposite directions from the site of annihilation. Multiple detectors on the scanner surrounding the patient identify each event, determine the site of origin, and reconstruct a map of distribution of the radionuclide. Because the path that the positron travels is unpredictable, this sets an irreducible limit to the resolution of the method, which with the currently available systems is in the order of 4–5 mm.

An approximation of the distribution of glucose metabolic rates can be derived by using ^{18}F labelled flurodeoxyglucose (FDG)—a glucose analogue. After cell membrane transport, FDG is phosphorylated to FDG-6 phosphate (FDG-6P), which is not metabolised thereafter. Because the cell is impermeable to FDG-6P, and enzymes that reverse this reaction are low in concentration, FDG is trapped within cells sufficiently long enough for rates of glucose uptake to be estimated. FDG uptake takes place over a 45 minute period and reflects the metabolic activity of the brain during that time. Some degree of standardisation of normal brain activity can be imposed by ensuring that during the period of FDG the subject lies with eyes closed in a quiet room. Although brief limited movement can be tolerated, young children may require sedation or a light general anaesthetic. The scan itself takes about 30–45 minutes and is not uncomfortable, threatening, or noisy. Where PET scanning is undertaken in the investigation of epilepsy, electroencephalogram (EEG) monitoring during the period of FDG uptake is recommended because ictal activity increases FDG uptake.

The radioactive isotopes are created in a cyclotron and it is the expense of this that limits the availability of these facilities. The isotopes have to be used within approximately one half life of the decay of the radioactive element. For ^{11}C compounds, this is about 20 minutes, but for ^{18}F it is about two hours, so FDG studies—for example, do not necessarily have to be on the same site as the cyclotron.

Cerebral blood volume can be derived using ^{11}C carboxyhaemoglobin derived from ^{11}C carbon monoxide; rates of cerebral blood flow can be derived using ^{15}O labelled water.

Consideration of radiation exposure is important but comparable with other procedures. For an FDG examination of the brain, the radiation dose is 6 mSv (effective dose equivalents). For a ^{11}C methionine (MET) scan, a further 3.5 mSv will be delivered. This compares with about 8 mSv for a barium enema, 5 mSv for a computed tomography (CT) scan, 4 mSv for an intravenous urogram, 3.8 mSv for a barium meal, and 2.1 mSv for a lumbar spine film. Background irradiation is about 2.2 mSv/year.

For clinical purposes, visual inspection of the images is often sufficient (qualitative PET). Absolute values for tissue metabolic rates, blood flow, and blood volume can be calculated if the activity of arterial samples is determined during the period of uptake (quantitative PET). In general, this has not been thought necessary in children. Alternative methods that measure simultaneous cardiac activity have been proposed. A compromise is to compare the activity of the structure under consideration (the region of interest) either with the whole brain activity, the analogous contralateral structure or a “neutral” structure, such as the cerebellum (“semi quantitative” PET). Each method has merits and disadvantages, particularly for the study of epilepsy (see below), but whichever region of comparison is used, semiquantitative analysis may be more sensitive than visual analysis by experienced personnel when—for example, there are rather diffuse changes in metabolic rate.

The production of new radiotracers offers synthetic chemists exciting but not insurmountable challenges. Thus, mapping the distribution of dopamine, opioid, benzodiazepine, and cholinergic receptors has all been achieved, with more to follow. The increasing

Department of
Paediatric Neurology,
Guy's Hospital, St
Thomas Street,
London SE1 9RT, UK
R O Robinson

Department of
Paediatric Neurology,
The General Royal
Infirmary at Leeds,
Belmont Grove, Leeds
LS2 9NS, UK
C D Ferrie

Paediatric Oncology
Department,
University Hospital,
Queens's Medical
Centre, Nottingham
NG7 2UH, UK
M Capra

Division of
Radiological Sciences,
Clinical PET Centre,
Guy's and St Thomas's
Hospitals, London
SE1 9RT, UK
M N Maisey

Correspondence to:
Dr Robinson.

realisation of the clinical applications of PET technology is being followed by the establishment of more clinical PET centres. The first facility in the world to be installed in a medical institute was at the MRC cyclotron unit at the Royal Post Graduate Medical Centre, London. Subsequent units in the UK have been established at Guy's and St Thomas's, London; Aberdeen Hospital; the National Hospital for Nervous Diseases, London; and Addenbrooke's Hospital, Cambridge

The epilepsies

Most reports of PET in the investigation of patients with epilepsies have used FDG to study the pattern of cerebral glucose metabolism.² This technique depends upon the different metabolic needs (and therefore uptake of FDG) of the cerebral cortex, giving rise to seizures, from that of the surrounding cortex. During seizures, there are large increases in the blood flow to and demand for metabolic substrates in the involved cortical area. This causes increased accumulation of FDG in epileptic foci during seizures ("hypermetabolism").³ However, the temporal resolution of FDG is ill suited for ictal studies; uptake of FDG takes 30–45 minutes and because most seizures last only a few minutes, "ictal" scans are usually a combination of interictal, ictal, and postictal data,⁴ making interpretation difficult.⁵ Clinical studies are nearly always planned to take place in the interictal state, with the expectation of detecting reduced FDG uptake ("hypometabolism"). The pathophysiology of interictal hypometabolism is unclear. It might simply reflect an obvious structural abnormality, such as a vascular malformation or hamartoma with decreased numbers of neurones and other metabolically active cells. However, in mesial temporal sclerosis, the degree of hypometabolism correlates poorly with neuronal loss. Reduced synaptic activity in non-functioning inhibitory neurones is an alternative explanation for interictal hypometabolism.^{4 6 7}

ANALYSIS OF PET SCANS IN CHILDREN WITH EPILEPSY

The most commonly used method of analysis of PET scans in patients with seizure disorders is visual inspection. This is analogous to the reporting of CT and magnetic resonance imaging (MRI) scans and depends on the differential uptake of radiotracer in different cerebral structures. Perceived limitations of visual inspection are the poor resolution of PET, the difficulty in identifying cerebral structures, and doubts as to whether the human eye is capable of detecting subtle abnormalities in radiotracer distribution.^{8 9}

The uptake of radiotracer in different cerebral structures can be quantified using tracer kinetic models and compared with data from controls. The objectivity of quantified methods makes them intuitively attractive but they are unpopular in clinical practice because large differences in radiotracer uptake between homologous cortical regions are detectable visually. Quantification is important, however,

because pronounced asymmetries between homologous cortical regions may exist, despite absolute metabolic rates for each lying within the "normal range". Unfortunately, quantified techniques are invasive, usually requiring repeated blood sampling from arterial lines. In paediatric practice there is also a lack of appropriate controls; maturational changes during infancy and childhood^{10–12} make adult control data unsuitable. Obtaining control data from normal children is ethically unacceptable.

Semiquantitative analysis, in which ratios of radiotracer uptake either from homologous brain regions or against a common comparator such as the cerebellum are calculated, is a third technique that avoids many of the problems of fully quantified techniques. Combined visual inspection and semiquantified analysis may maximise the detection of cortical metabolic defects in children with epilepsy.¹³

Techniques for the co-registration of PET and MRI images have been developed.^{14–26} However, in patients with epilepsy, the metabolic abnormality revealed by FDG PET is usually large, making it unlikely that co-registration will improve the localisation of the epileptic focus.

FDG PET AND PARTIAL EPILEPSIES

Patients with partial seizures arising in the temporal lobes have been the subject of a number of studies with FDG PET. Most patients investigated have been adults with mesial temporal sclerosis. FDG PET detects focal areas of hypometabolism interictally in 70–80% of patients.^{27–29} The abnormality usually extends well beyond the mesial temporal lobe structures and the nadir of FDG uptake may be in the lateral temporal lobe. Ipsilateral thalamic hypometabolism may also be seen.^{4 30} It has been claimed that FDG PET is more sensitive than MRI in detecting abnormalities in patients with temporal lobe epilepsy.³¹ However, this is unlikely to apply to the advanced MRI techniques used today.³² The superior resolution and localising value of MRI over PET, the lack of exposure to radiation, and the additional anatomical information gained makes it the investigation of choice.³³

FDG PET is considerably less sensitive in detecting abnormalities in patients with partial seizures arising from the temporal neocortex and extratemporal sites. Fewer than 50% of patients show regions of interictal hypometabolism.^{33–36} Moreover, abnormalities are often regional or hemispheric and therefore of poor localising value. It is hoped that newer techniques will improve the sensitivity of FDG PET in these epilepsies.³⁷

INFANTILE SPASMS

Infantile spasms occurring as part of West syndrome are an age dependent reaction to a variety of cerebral insults, both generalised and focal. After conventional investigations a cause is found in 89–90% of cases.³⁸ Identification of patients whose spasms are caused by focal brain pathology is important because resective surgery might offer the best hope of seizure

control and maximise the developmental potential of the child.³⁹ The occurrence of partial seizures before or concurrent with spasms, asymmetrical spasms, and atypical forms of hypsarrhythmia (especially if asymmetrical or with a fixed focus) increase the likelihood of focal cerebral pathology.^{38 40-43} Full evaluation requires detailed EEG (including video EEG) and high resolution MRI studies.

A number of groups have studied children with infantile spasms using FDG PET.⁴⁴⁻⁴⁹ Chugani and colleagues have reported about 30 children with either active spasms or a past history of spasms in whom FDG PET revealed focal cortical metabolic abnormalities and who subsequently underwent focal resections.⁴⁴⁻⁴⁶ Some of these children might reasonably have been operated upon based on clinical, EEG, and structural neuroimaging criteria, but PET made the likelihood of focal cortical pathology more certain in others, enabling surgery to proceed without the need for invasive EEG studies. Histological examination of resected cortex demonstrated circumscribed areas of cortical dysgenesis, often subtle and probably undetectable even with high quality MRI.^{50 51} It may even be worthwhile studying those children with spasms who lack electroclinical evidence of focal cortical pathology.⁴⁷ However, some caution is required, because some metabolic abnormalities in children with spasms can be transient, possibly dependent on the timing of the PET scans in relation to the duration of the seizure diathesis.⁴⁹

LENNOX-GASTAUT SYNDROME

A number of different patterns of cerebral glucose metabolism have been reported using FDG PET in the Lennox-Gastaut syndrome. This might reflect differences in how this syndrome is defined. If the definition is extended to include children with drug resistant seizures of multiple type (both generalised, partial, and of undetermined type) and learning difficulties, many children with partial epilepsies will be included and FDG PET may reveal focal, multifocal, or hemispheric metabolic defects.⁵²⁻⁵⁵ These abnormalities are usually congruent with clinical, surface EEG, and structural neuroimaging findings. However, except for case studies,⁵² there are no reports on children who have subsequently undergone surgical treatment based on FDG PET findings. If the term Lennox-Gastaut syndrome is confined to patients with multiple generalised seizures (tonic, atonic, and atypical absences, but not frequent myoclonic seizures), with generalised slow spike and wave EEG discharges (and generalised fast spikes in sleep), focal cortical metabolic abnormalities are not shown by FDG PET.^{56 57} The most common pattern seen in such patients is one of diffuse interictal hypometabolism.

OTHER CATASTROPHIC EPILEPSIES OF EARLY CHILDHOOD

There is little experience of FDG PET in other severe childhood epilepsies. FDG PET findings have been reported in a few children with severe myoclonic epilepsy in infancy and

epilepsy with myoclonic astatic seizures.⁴⁷ No consistent pattern of abnormalities was revealed, although there was a high incidence of diffuse or multifocal metabolic defects. FDG PET abnormalities have also been described in children with Sturge-Weber syndrome, tuberous sclerosis, and hemimegalencephaly.⁵⁸⁻⁶¹ These conditions are all better diagnosed with structural imaging methods, although there are anecdotal claims that FDG PET can give useful additional information regarding disease progression in the Sturge-Weber syndrome⁵⁸ and abnormalities of the "non-involved" hemisphere in hemimegalencephaly.⁶¹ Reports of FDG PET findings in Raussmussen's encephalitis suggests that either focal hypermetabolism or widespread hypometabolism may be seen, presumably reflecting the timing of the scan in relation both to seizures and to the chronicity of the process.⁶² In the Landau-Kleffner syndrome, FDG PET findings appear to be variable. Maquet *et al* found focal or regional hypermetabolism during both sleep and wakefulness in five of six children scanned during the active phase of the disease.⁶³ In the other patient, decreased metabolism was apparent during wakefulness, with a variable pattern during sleep. In the recovery phase of the disease, cortical metabolism was either normal or showed various patterns of focal or regional hypometabolism.

PET AND SPECT

The relative merits of PET and single photon emission computed tomography (SPECT) in the investigation of children being considered for epilepsy surgery have been debated. SPECT tracers for the measurement of cerebral blood flow are commercially available, making the technique widely available for centres with gamma cameras. Interictal SPECT is substantially less sensitive than interictal PET in detecting cortical abnormalities in patients with temporal lobe seizures.⁶⁴ However, the temporal characteristics of the main radiotracers used in SPECT while studying cerebral blood flow make ictal scans feasible. This latter facility is particularly attractive for studying patients with seizure disorders. The sensitivities of ictal SPECT and interictal PET are similar, at least for the detection of temporal lobe foci.⁶⁵ The resolution of SPECT is substantially inferior to that of PET, and the areas of abnormality detected are larger than those typically seen with PET. Claims that the two techniques (ictal SPECT and interictal PET) might be complimentary need further evaluation, bearing in mind the radiation exposure. The FDG technique has now been adapted for use with SPECT, but there are as yet no reports in which this has been compared with interictal FDG PET.

OTHER PET LIGANDS AND OTHER INSIGHTS PROVIDED BY PET

Experience with ligands other than FDG is limited. Flumazenil is an antagonist at benzodiazepine receptors and shows reduced interictal binding in the epileptic focus of patients with partial seizures. In such patients, the extent of

abnormality revealed with flumazenil appears to be smaller than that shown by FDG, giving improved localisation.⁶⁶ Flumazenil might also be more sensitive than FDG in the detection of extratemporal foci,⁶⁷ and might be useful in detecting areas of cortical dysgenesis.⁶⁸ The total radiation exposure is considerably less with flumazenil than with FDG.

PET offers unique opportunities to study the pathogenesis of seizure disorders and the action of antiepileptic drugs. The role of various neurotransmitter systems in seizure disorders can be studied using specific ligands.⁴ FDG PET has revealed differential effects of antiepileptic drugs on cerebral glucose metabolism, which might help to explain their different profiles of adverse effects.⁶⁹⁻⁷³ In the future, the distribution and binding of specific drugs will be studied by means of drug labelled PET ligands.

FDG PET studies have revealed evidence of subcortical, particularly thalamic, involvement in partial and idiopathic generalised epilepsies and in the epileptic encephalopathies of childhood, including the Lennox-Gastaut syndrome.⁵⁷ In West's syndrome, there is evidence of lentiform and brain stem activation by "antiepileptic drugs",⁷⁴ leading to new hypotheses concerning the aetiology of these seizure disorders. FDG PET studies have suggested a relation between frontal lobe hypometabolism and the mental dysfunction commonly seen in the childhood epileptic encephalopathies.⁷⁵ A further area of interest is the future use of PET activation studies for the non-invasive localisation of eloquent areas of cortex⁷⁶—an ability already possible with functional MRI, with which PET will need to be compared.

GUIDELINES

Investigating children with seizure disorders using functional neuroimaging should only be undertaken as part of a planned programme in patients being considered for possible epilepsy surgery. The choice between PET and SPECT is likely to be determined mainly by local availability. All children with partial epilepsies that are not within the spectrum of the idiopathic (age related) epilepsies of childhood, who are resistant to appropriate antiepileptic medication, should be considered as potential surgical candidates. Clinical evaluation, surface EEG (including video telemetry recordings) and high resolution MRI are the key investigations. If all point clearly to a mesial temporal lobe focus and there are no conflicting data or data suggesting bilateral foci, functional neuroimaging is probably redundant.⁷⁷ Where two of the modalities indicate a likely mesial temporal lobe focus, but the third is non-informative, or functional neuroimaging findings are equivocal, resective surgery should proceed without invasive EEG studies. When there is conflict or discrepancy between clinical, electrical, and MRI data, functional neuroimaging may help in the planning of invasive EEG studies.

Localisation of seizures arising in sites other than the mesial temporal lobes is more

difficult. Unless a lesion is apparent on MRI, nearly all such patients will require invasive EEG studies. Functional neuroimaging has a potential role in such patients in helping to plan invasive EEG studies. However, the reduced sensitivity of PET and SPECT in detecting abnormalities in extratemporal sites limits their usefulness.

Children with cryptogenic infantile spasms who do not respond rapidly to medical treatment should be investigated carefully by a tertiary epilepsy centre for the possibility of a cortical focus. PET might have a useful role to play in such cases. PET has not been shown to be sufficiently useful in other childhood epilepsies to justify its use outside of research studies.

Brain tumours

To date, most work with PET has been in adults with malignant gliomas. The resulting findings, extensive and promising as they are, unfortunately cannot be extrapolated directly to childhood brain tumours, because the spectrum of histology differs, and where the tumour type is identical, its clinical behaviour may vary with age.⁷⁸ In addition, the pattern of metabolic activity in the infant brain is different from the adult brain; children's brains attain adult characteristics by the age of 4, with only minor changes occurring through the rest of the first 2 decades.⁷⁹⁻⁸⁰

PET has the potential to play an important role in the prediction of tumour grade and therefore prognosis, assessing the response to treatment, differentiating between iatrogenic lesions and residual and recurrent tumour, evaluating the neurotoxicity of treatment, and finally studying the action of chemotherapeutic agents.

The rationale behind PET in brain tumours is that the rate of glycolysis of rapidly growing malignant tumours is greater than that of normal tissue.⁸¹ In 1982, Di Chiro *et al* demonstrated, *in vivo*, not only that FDG uptake is increased in malignant cells, but also that the increase is proportional to the histological grade of cerebral gliomas.⁸² Subsequently, this has been shown to hold true for posterior fossa tumours in children.⁸³⁻⁸⁴ The rate of FDG uptake correlates well with patient survival time,⁸⁵⁻⁸⁸ with some authors claiming superiority of PET over histological grade in predicting long term outcome.⁸⁹ Thus, in the future, the need for taking tumour biopsies associated with a high risk of morbidity may be reduced. PET might provide an additional tool to deal with two of the challenging characteristics (among many others) of brain tumours: the potential for heterogeneous histology within a tumour and the potential to transform from a benign to a malignant tumour. Remembering that the histological grade of a heterogeneous tumour is determined by the cell(s) with the greatest malignant transformation, PET can be useful in guiding a stereotactic biopsy needle to the area of greatest FDG uptake.⁹⁰⁻⁹³ Malignant change in low grade gliomas may be identified by PET before clinical manifestations of the disease or MRI changes become apparent.⁹⁴⁻⁹⁵

Because FDG uptake is an indicator of the overall rate of glycolysis and glucose cell membrane transport, a specific marker for protein synthesis, such as an amino acid, might also be expected to reflect tumour activity. This is indeed true, with the radiotracer MET having several advantages over other amino acids in the study of cerebral protein synthesis.⁹⁶ The uptake of methionine by tumours is related to cell membrane transport of the amino acid and the rate of protein synthesis,⁹⁷ not to changes in the blood-brain barrier.⁹⁸⁻⁹⁹ The advantages of MET over FDG is that it has a higher tumour to normal brain uptake ratio, thereby delineating the extent of the tumour more accurately, particularly for lower grade tumours.⁹⁸⁻¹⁰⁰ Although the degree of MET uptake correlates broadly with histologically assessed tumour grade, it is not as sensitive as FDG in this respect.⁹⁸⁻¹⁰¹ There is evidence to suggest that amino acid transport into tumour cells, rather than uptake and protein synthesis, is more closely associated with tumour grade.¹⁰²

FDG and MET uptake occurs not only in tumours but also in inflammatory or infectious foci.¹⁰³⁻¹⁰⁵ For this reason, patients with AIDS pose a particular problem. To date, no ligand has been identified to differentiate the above two pathologies.

Newer developments include the study of DNA synthesis with ¹¹C thymidine or iodine labelled pyrimidine analogues, such as iododeoxyuridine, which could provide a more specific measure of tumour cell division.¹⁰⁶⁻¹⁰⁷ This information has not yet been applied clinically on any scale. Benzodiazepine receptors have been shown to be increased 20-fold in brain tumours compared with normal brain.¹⁰⁸ As a result, ¹¹C flumazenil, a benzodiazepine antagonist, can now be used to delineate the outer margins of brain tumours.¹⁰⁹⁻¹¹⁰

PET has the potential to monitor treatment. A large reduction in the rate of glucose utilisation within a tumour reflects effective treatment.¹¹¹⁻¹¹³ Chemotherapy and radiotherapy can cause an initial rise in FDG uptake, which falls over the subsequent seven to 30 days. The amount of rise and fall are inversely and directly proportional, respectively, to patient survival.¹¹⁴⁻¹¹⁶ In a small study, FDG was useful in assessing the response to treatment of medulloblastoma in children.⁸⁴ In our experience, this has been useful, especially after radiotherapy, when the FDG uptake of a tumour may be reduced relative to the uptake of the surrounding brain. In a more recent study, MET was found to be superior to FDG in monitoring the effects of treatment in low grade gliomas.¹¹⁷ Combining information obtained from iododeoxyuridine and FDG uptake was a sensitive indicator of response in glial tumours.¹¹⁸ The dose of radiation involved in PET limits the frequency of serial studies.

Routine MRI (with the exception of diffusion weighted MRI)¹¹⁹⁻¹²⁰ and CT cannot reliably distinguish residual or recurrent tumours from postoperative inflammatory changes or radionecrosis.¹¹⁷⁻¹¹⁸⁻¹²¹⁻¹²³ The distinction is crucial in planning future management. PET can be invaluable in this regard

because postoperative change in itself does not cause increased FDG uptake at the site of surgery.¹²⁴⁻¹²⁵ Radiation necrosis causes decreased FDG uptake¹²⁶⁻¹²⁸ allowing ready discrimination from tumour. Low grade tumours may cause a problem because FDG uptake in these tumours may be less than is found in normal cortex. In this situation, combining FDG with MET PET is particularly helpful.¹²⁹ In our experience, this has been useful, especially during the post-radiotherapy reduction in FDG uptake. Additional sensitivity may be gained by prior phenylalanine loading. Phenylalanine competitively inhibits MET uptake in brain tumours but not in non-neoplastic lesions.⁹⁸⁻⁹⁹

Resolution of PET (at best ~ 4 mm) remains poor when compared with morphological imaging. As a result, anatomical localisation using PET alone may be difficult. However, co-registration of PET with CT or MRI is now possible, enabling accurate matching of cellular activity with anatomy.

PET can be used to monitor the unwanted effects of treatment. Long term survivors of acute lymphoblastic leukaemia, who received cranial irradiation and intrathecal chemotherapy, had a reduction in FDG uptake proportional to the reduction in their overall IQ.¹³⁰

PET can be used to study the delivery and the pharmacokinetics of chemotherapeutic agents, the most readily studied being 1,3-bis(2-chloroethyl)nitrosourea (BCNU).¹³¹⁻¹³² PET has also been used to monitor the kinetics of radiolabelled monoclonal antibodies—a technique with potential for identifying, as well as treating, tumours.¹³²⁻¹³⁵ Both of these applications of PET remain research based.

Conclusion

The amount of information to be gained from applying the increasing number of radionuclides in a variety of pathological situations is potentially very great. However, expense and scarcity of resources, as well as the radiation, discomfort, and anxiety to the individual child has, as with all diagnostic procedures, to be weighed against the potential benefit to be derived from the information gained from the investigation. Although the radiation involved is trivial in relation to radiotherapy doses, and is similar to many x ray and SPECT procedures, we feel that PET examination is only justifiable when the information is likely to affect management decisions, such as whether to offer surgery to a child with intractable epilepsy or re-operate in a child with recurrent tumour.

The technique is still in its infancy. Most of the references in this review were written within the past five years. There can be no doubt that the development of further ligands will find applications in a wide variety of conditions in paediatrics.

1 Joliot F, Curie I. Artificial production of a new kind of radio-element. *Nature* 1934;133:201-2.

2 Editorial. SPECT and PET in epilepsy. *Lancet* 1989;i: 135-6.

- 3 Phelps ME. Positron emission tomography (PET). In: Mazziotta JC, Gilman S, eds. *Clinical brain imaging: principles and applications*. Philadelphia: FA Davis, 1992:71-107.
- 4 Theodore WH. Epilepsy. In: Mazziotta JC, Gilman S, eds. *Clinical brain imaging: principles and applications*. Philadelphia: FA Davis, 1992:136-65.
- 5 Chugani HT, Rintahaka PJ, Shewmon DA. Ictal patterns of cerebral glucose utilization in children with epilepsy. *Epilepsia* 1994;35:813-22.
- 6 Engel J, Brown WJ, Kuhl DE, Phelps ME, Mazziotta JC, Crandall PH. Pathological findings underlying focal temporal lobe hypometabolism in partial epilepsy. *Ann Neurol* 1982;12:518-29.
- 7 Theodore WH, Katz D, Kufta C, Sato S, et al. Pathology of temporal lobe foci: correlation with CT, MRI and PET. *Neurology* 1990;40:797-803.
- 8 Sadzot B, Debets P, Comar C, Franck G. PET studies of patients with partial epilepsy: visual interpretation vs. semi-quantification/quantification. *Acta Neurol Scand* 1994;152(suppl):175-8.
- 9 Friberg L, Lomberg-Jensen H. Visual inspection versus quantified analysis of functional brain images. *Acta Neurol Scand Suppl* 1994;152:169-74.
- 10 Chugani HT, Phelps ME. Maturation changes in cerebral function in infants determined by ¹⁸FDG positron emission tomography. *Science* 1986;231:840-3.
- 11 Chugani HT, Phelps ME, Mazziotta JC. Positron emission tomography study of human brain functional development. *Ann Neurol* 1987;22:487-97.
- 12 Kinnala A, Suhonen-Polvi H, Kero P, et al. Cerebral metabolic rate for glucose during the first six months of life: an FDG positron emission tomography study. *Arch Dis Child* 1996;74:F153-7.
- 13 Ferrie CD, Marsden PK, Maisey MN, Robinson RO. Visual and semiquantitative analysis of cortical FDG PET scans in childhood epileptic encephalopathies. *J Nucl Med* 1997;38:1891-4.
- 14 Meltzer CC, Bryan RN, Holcomb HH, et al. Anatomical localization for PET using MR imaging. *J Comput Assist Tomogr* 1990;14:418-26.
- 15 Muller-Gartner HW, Links JM, Prince JL, et al. Measurement of radiotracer concentration in brain grey matter using positron emission tomography: MRI-based correction for partial volume effects. *J Cereb Blood Flow Metab* 1992;12:571-83.
- 16 Alpert NM, Bradshaw JF, Kennedy D, Correia JA. The principal axes transformation—a method for image registration. *J Nucl Med* 1990;31:1717-22.
- 17 Turkington TG, Jaszcak RJ, Pelizzari CA, et al. Accuracy of registration of PET, SPECT and MR images of a brain phantom. *J Nucl Med* 1993;34:1587-94.
- 18 Rusinek H, Tsui W, Levy AV, Noz ME, de Leon MJ. Principal axes and surface fitting methods for three-dimensional image registration. *J Nucl Med* 1993;34:2019-24.
- 19 Woods RP, Mazziotta JC, Cherry SR. MRI-PET registration with automated algorithm. *J Comput Assist Tomogr* 1993;17:536-46.
- 20 Kapouleas I, Alavi A, Alves WM, Gur RE. Registration of three-dimensional MR and PET images of the human brain without markers. *Radiology* 1991;181:731-9.
- 21 Steinmetz H, Huang Y, Seitz RJ, et al. Individual integration of positron emission tomography and high-resolution magnetic resonance imaging. *J Cereb Blood Flow Metab* 1992;12:919-26.
- 22 Meltzer CC, Leal JP, Mayberg HS, Wagner HN, Frost JJ. Correction of PET data for partial volume effects in human cerebral cortex by MR imaging. *J Comput Assist Tomogr* 1990;14:561-70.
- 23 Cooper MD, Chen CT, Pelizzari CA, Levin DN, Chen GTY. Analysis of PET images with the use of anatomic information from CT and MR images. *Acta Radiol* 1991;376(suppl):85-6.
- 24 Mangin J, Frouin V, Bloch I, Bendriem B, Lopez-Krahe J. Fast nonsupervised 3D registration of PET and MR images of the brain. *J Cereb Blood Flow Metab* 1994;14:749-62.
- 25 Ge Y, Fitzpatrick JM, Votaw JR, et al. Retrospective registration of PET and MR brain images: an algorithm and its stereotactic validation. *J Comput Assist Tomogr* 1994;18:800-10.
- 26 Rehm K, Strother SC, Anderson JR, Schaper KA, Rottenberg DA. Display of merged multimodality brain images using interleaved pixels with independent color scales. *J Nucl Med* 1994;35:1815-21.
- 27 Abou-Khalil BW, Siegel GJ, Sackellar JC, Gilman S, Hichwa R, Marshall R. Positron emission tomography studies of cerebral glucose metabolism in chronic partial epilepsy. *Ann Neurol* 1987;22:480-6.
- 28 Engel J, Kuhl DE, Phelps ME, Mazziotta JC. Interictal cerebral glucose metabolism in partial epilepsy and its relation to EEG changes. *Ann Neurol* 1982;12:510-17.
- 29 Theodore WH, Newmark ME, Sato S, et al. 18-F-fluorodeoxyglucose positron emission tomography in refractory complex partial seizures. *Ann Neurol* 1983;14:429-37.
- 30 Henry TR, Arbor A, Engel J, Phelps ME. Subcortical hypometabolism on interictal PET in complex partial seizures. *Neurology* 1989;39(suppl 1):299.
- 31 Chugani HT. The use of positron emission tomography in the clinical assessment of epilepsy. *Semin Nucl Med* 1992;XXII:247-53.
- 32 Jackson GD. New techniques in magnetic resonance and epilepsy. *Epilepsia* 1994;35(suppl 6):2-13.
- 33 Theodore WH. Neuroimaging in the evaluation of patients for focal resection. In: Wyllie E, ed. *The treatment of epilepsy: principles and practice*. Philadelphia: Lea and Febiger, 1993:1039-50.
- 34 Swartz BE, Halgren E, Delgado-Escueta AV, et al. Neuroimaging in patients with seizures of probable frontal lobe origin. *Epilepsia* 1989;30:547-58.
- 35 Henry TR, Sutherland WW, Engel J, et al. Interictal cerebral metabolism in partial epilepsies of neocortical origin. In: Luders H, ed. *Epilepsy surgery*. New York: Raven Press, 1991:243-50.
- 36 Henry JR, Mazziotta JC, Engel J. The functional anatomy of frontal lobe epilepsy studied with PET. In: Chauvel P, Delgado-Escueta AV, Halgren E, Bancaud J, eds. *Frontal lobe seizures and epilepsies*. New York: Raven Press, 1992:449-63.
- 37 Swartz BE, Khonsari A, Brown C, Mandelkern M, Simkins F, Kriskakumtorn T. Improved sensitivity of ¹⁸FDG-positron emission tomography scans in frontal and "frontal plus" epilepsy. *Epilepsia* 1995;36:388-95.
- 38 Aicardi J. *Epilepsy in children*. New York: Raven Press, 1994.
- 39 Wyllie E. Surgery for catastrophic localization-related epilepsy in infants. *Epilepsia* 1996;37(suppl 1):22-5.
- 40 Carrazana EJ, Lombroso CT, Mikati M, Helmers S, Holmes GL. Facilitation of infantile spasms by partial seizures. *Epilepsia* 1993;34:97-109.
- 41 Ploin P, Dulac O, Jalin C, Chiron C. Twenty-four hour ambulatory EEG monitoring in infantile spasms. *Epilepsia* 1993;34:689-91.
- 42 Olson DM, Chugani HT, Shewmon A, Phelps ME, Peacock WJ. Electroencephalographic confirmation of focal positron emission tomographic abnormalities in children with intractable epilepsy. *Epilepsia* 1990;31:731-9.
- 43 Gaily EK, Shewmon DA, Chugani DC, Curran JG. Asymmetric and asynchronous infantile spasms. *Epilepsia* 1995;36:872-82.
- 44 Chugani HT, Shields WD, Shewmon DA, Olson DM, Phelps ME, Peacock WJ. Infantile spasms. I. PET identifies focal cortical dysgenesis in cryptogenic cases for surgical treatment. *Ann Neurol* 1990;27:406-13.
- 45 Chugani HT, Shewmon DA, Shields WD, et al. Surgery for intractable infantile spasms: neuroimaging perspectives. *Epilepsia* 1993;34:764-71.
- 46 Chugani HT, Da Silva E, Chugani DC. Infantile spasms: III. Prognostic implications of bitemporal hypometabolism on positron emission tomography. *Ann Neurol* 1996;39:643-9.
- 47 Ferrie CD, Maisey M, Cox T, et al. Focal abnormalities detected by ¹⁸FDG PET in epileptic encephalopathies. *Arch Dis Child* 1996;75:1-6.
- 48 Wyllie E, Comair Y, Ruggieri P, Raja S, Prayson R. Epilepsy surgery in the setting of periventricular leukomalacia and focal cortical dysplasia. *Neurology* 1996;46:839-41.
- 49 Maeda N, Watanabe K, Negoro T, et al. Evolutional changes of cortical hypometabolism in West's syndrome. *Lancet* 1994;343:1620-3.
- 50 Vinters HV, Fisher RS, Cornford ME, et al. Morphological substrates of infantile spasms: studies based on surgically resected cerebral tissue. *Childs Nerv Syst* 1992;8:8-17.
- 51 Vinters HV, De Rosa MJ, Farrell MA. Neuropathologic study of resected cerebral tissue from patients with infantile spasms. *Epilepsia* 1993;34:772-9.
- 52 Gur RC, Sussman NM, Alavi A, et al. Positron emission tomography in two cases of childhood epileptic encephalopathy (Lennox-Gastaut syndrome). *Neurology* 1982;32:1191-4.
- 53 Chugani HT, Mazziotta JC, Engel J, Phelps ME. The Lennox-Gastaut syndrome: metabolic subtypes determined by 2-deoxy-2-fluoro-D-glucose positron emission tomography. *Ann Neurol* 1987;21:4-13.
- 54 Yanai K, Iinuma K, Matsuzawa T, et al. Cerebral glucose utilization in pediatric neurological disorders determined by positron emission tomography. *Eur J Nucl Med* 1987;13:292-6.
- 55 Miyauchi T, Nomura Y, Ohno S, Kishimoto H, Matsushita M. Positron emission tomography in three cases of Lennox-Gastaut syndrome. *Jpn J Psychiatry Neurol* 1988;42:795-804.
- 56 Theodore WH, Rose D, Patronas N, et al. Cerebral glucose metabolism in the Lennox-Gastaut syndrome. *Ann Neurol* 1987;21:14-21.
- 57 Ferrie CD, Marsden PK, Maisey MN, Robinson RO. Cortical and subcortical glucose metabolism in childhood epileptic encephalopathies. *J Neurol Neurosurg Psychiatry* 1997;63:181-7.
- 58 Chugani HT, Mazziotta JC, Phelps ME. Sturge-Weber syndrome: a study of cerebral glucose utilization with positron emission tomography. *J Pediatr* 1989;114:244-53.
- 59 Chugani HT. Imaging: anatomic and functional. In: Wallace SJ, ed. *Epilepsy in children*. London: Chapman and Hall, 1996:483-506.
- 60 Széles B, Herholz K, Heiss WD, et al. Hypometabolic cortical lesions in tuberous sclerosis with epilepsy: demonstration by positron emission tomography. *J Comput Assist Tomogr* 1983;7:946-53.
- 61 Rintahaka PJ, Chugani HT, Messa C, Phelps ME. Hemimegalencephaly: evaluation with positron emission tomography. *Pediatr Neurol* 1993;9:21-8.
- 62 Tien RD, Ashdown BC, Lewis DV, Atkins MR, Burger PC. Rasmussen's encephalitis: neuroimaging findings in four patients. *AJR Am J Roentgenol* 1992;158:1329-32.
- 63 Maquet P, Hirsch E, Metz-Lutz MN, et al. Regional cerebral glucose metabolism in children with deterioration of one or more cognitive functions and continuous spike-and-wave discharges during sleep. *Brain* 1995;118:1497-520.

- 64 Nagata T, Tanaka F, Yonekura Y, *et al.* Limited value of interictal brain perfusion SPECT for detection of epileptic foci: high resolution SPECT studies in comparison with FDG-PET. *Ann Nucl Med* 1995;9:59-63.
- 65 Ho SS, Berkovic SF, Berlangieri SU, *et al.* Comparison of ictal SPECT and interictal PET in the pre-surgical evaluation of temporal lobe epilepsy. *Ann Neurol* 1995;37:738-45.
- 66 Savic J, Ingvar M, Stone-Elinder S. Comparison of [^{11}C]flumazenil and [^{18}F]FDG as PET markers of epileptic foci. *J Neurol Neurosurg Psychiatry* 1993;56:615-21.
- 67 Savic J, Thorell JO, Roland P. [^{11}C]flumazenil positron emission tomography visualizes frontal epileptogenic regions. *Epilepsia* 1995;36:1225-32.
- 68 Richardson MP, Koepp MJ, Brooks DJ, Fish DR, Duncan JS. Benzodiazepine receptors in focal epilepsy with cortical dysgenesis: an [^{11}C]flumazenil PET study. *Ann Neurol* 1996;40:188-98.
- 69 Theodore WH, DiChiro G, Margolin R, Fishbein D, Porter RJ, Brooks RA. Barbiturates reduce human cerebral glucose metabolism. *Neurology* 1986;36:60-4.
- 70 Leiderman DB, Balish M, Bromfield EB, Theodore WH. Effect of valproate on human cerebral glucose metabolism. *Epilepsia* 1991;32:417-22.
- 71 Foster NL, VanDerSpek AFL, Aldrich MS, *et al.* The effect of diazepam sedation on cerebral glucose metabolism in Alzheimer's disease as measured using positron emission tomography. *J Cereb Blood Flow Metab* 1987;7:415-20.
- 72 Theodore WH. Antiepileptic drugs and cerebral glucose metabolism. *Epilepsia* 1988;29(suppl 2):48-55.
- 73 Gaillard WD, Zeffiro T, Fazilat S, DeCarli C, Theodore WH. Effect of valproate on cerebral metabolism and blood flow: an [^{18}F]2-deoxyglucose and [^{15}O] water positron emission tomography study. *Epilepsia* 1996;37:515-21.
- 74 Chugani HT, Shewmon DA, Sankar R, Chen BC, Phelps ME. Infantile spasms: II. Lenticular nuclei and brain stem activation on positron emission tomography. *Ann Neurol* 1992;31:212-19.
- 75 Ferrie CD, Madigan C, Tilling R, Maisey MN, Marsden PK, Robinson RO. Adaptive and maladaptive behaviour in children with epileptic encephalopathies: correlation with cerebral glucose metabolism. *Dev Med Child Neurol* 1997;39:588-95.
- 76 Perrine K. Future directions for functional mapping. *Epilepsia* 1994;35(suppl 6):90-102.
- 77 Polkey CE. Surgical treatment of epilepsy in children. In: Wallace S, ed. *Epilepsy in children*. London: Chapman and Hall, 1996:561-600.
- 78 Laws WE, Taylor WP, Clifton MB, Okazaki K. Neurosurgical management of low-grade astrocytoma of the cerebral hemispheres. *J Neurosurg* 1984;61:665-73.
- 79 O'Tuama LA, Phillips PC, Smith QR, *et al.* L-methionine uptake by human cerebral cortex: maturation from infancy to old age [see comments]. *J Nucl Med* 1991;32:16-22.
- 80 Chugani HT. Functional brain imaging in pediatrics. *Pediatr Clin North Am* 1992;39:777-99.
- 81 Warburg O. On the origin of cancer cells. *Science* 1956;123:309-14.
- 82 Di Chiro G, DeLaz RL, Brooks RA, *et al.* Glucose utilization of cerebral gliomas measured by [^{18}F] fluorodeoxyglucose and positron emission tomography. *Neurology* 1982;32:1323-9.
- 83 Hoffman JM, Hanson MW, Friedman HS, *et al.* FDG-PET in pediatric posterior fossa brain tumors. *J Comput Assist Tomogr* 1992;16:62-8.
- 84 Holthoff VA, Herholz K, Berthold F, *et al.* In vivo metabolism of childhood posterior fossa tumors and primitive neuroectodermal tumors before and after treatment. *Cancer* 1993;72:1394-403.
- 85 Patronas NJ, Di Chiro G, Kufta C, *et al.* Prediction of survival in glioma patients by means of positron emission tomography. *J Neurosurg* 1985;62:816-22.
- 86 Schifter T, Hoffman JM, Hanson MW, *et al.* Serial FDG-PET studies in the prediction of survival in patients with primary brain tumors. *J Comput Assist Tomogr* 1993;17:509-61.
- 87 Alavi JB, Alavi A, Chawluk J, *et al.* Positron emission tomography in patients with glioma. A predictor of prognosis. *Cancer* 1988;62:1074-8.
- 88 Mineura K, Sasajima T, Kowada M, *et al.* Perfusion and metabolism in predicting the survival of patients with cerebral gliomas. *Cancer* 1994;73:2386-94.
- 89 Di Chiro G. Positron emission tomography using [^{18}F] fluorodeoxyglucose in brain tumors. A powerful diagnostic and prognostic tool. *Invest Radiol* 1987;22:360-71.
- 90 Levivier M, Goldman S, Pirotte B, *et al.* Diagnostic yield of stereotactic brain biopsy guided by positron emission tomography with [^{18}F]fluorodeoxyglucose. *J Neurosurg* 1995;82:445-52.
- 91 Hanson MW, Glantz MJ, Hoffman JM, *et al.* FDG-PET in the selection of brain lesions for biopsy. *J Comput Assist Tomogr* 1991;15:796-801.
- 92 Levivier M, Goldman S, Bidaut LM, *et al.* Positron emission tomography-guided stereotactic brain biopsy. *Neurosurgery* 1992;31:792-7.
- 93 Pirotte B, Goldman S, Brucher JM, *et al.* PET in stereotactic conditions increases the diagnostic yield of brain biopsy. *Stereotact Funct Neurosurg* 1994;63:144-9.
- 94 Rubenstein LJ. The correlation of neoplastic vulnerability with central neuroepithelial cytogeny and glioma differentiation. *J Neurooncol* 1987;5:11-27.
- 95 Francavilla TL, Miletich RS, Di Chiro G, Patronas NJ, Rizzoli HV, Wright DC. Positron emission tomography in the detection of malignant degeneration of low-grade gliomas. *Neurosurgery* 1989;24:1-5.
- 96 Phillips PC. Positron emission tomographic studies of transport and metabolism in brain tumors. In: Packer R, Bleyer WA, Pochedly C, eds. *Pediatric neurooncology*. Philadelphia: Harwood Academic Publishers, 1992:91-110.
- 97 Bustany P, Chatel M, Derlon JM, *et al.* Brain tumor protein synthesis and histological grades: a study by positron emission tomography (PET) with C11-L-methionine. *J Neurooncol* 1986;3:397-404.
- 98 O'Tuama LA, Phillips PC, Strauss LC, *et al.* Two-phase [^{11}C]L-methionine PET in childhood brain tumors. *Pediatr Neurol* 1990;6:163-70.
- 99 Bergstrom M, Ericson K, Hagenfeldt L, *et al.* PET study of methionine accumulation in glioma and normal brain tissue: competition with branched chain amino acids. *J Comput Assist Tomogr* 1987;11:208-13.
- 100 Lilja A, Bergstrom K, Hartvig P, *et al.* Dynamic study of supratentorial gliomas with L-methyl- ^{11}C -methionine and positron emission tomography. *AJNR Am J Neuroradiol* 1985;6:505-14.
- 101 Ogawa T, Shishido F, Kanno I, *et al.* Cerebral glioma: evaluation with methionine PET [see comments]. *Radiology* 1993;186:45-53.
- 102 Meyer GJ, Harre R, Orth F. In vivo protein synthesis in human brain tumors measured with ^{11}C -L-methionine. *J Nucl Med* 1989;30:910-11.
- 103 Ishii K, Ogawa T, Hatazawa J, *et al.* High L-methyl- ^{11}C methionine uptake in brain abscess: a PET study. *J Comput Assist Tomogr* 1993;17:660-61.
- 104 Conti PS. Introduction to imaging brain tumor metabolism with positron emission tomography (PET) [review]. *Cancer Invest* 1995;13:244-59.
- 105 Villringer K, Jager H, Dichgans M, *et al.* Differential diagnosis of CNS lesions in AIDS patients by FDG-PET. *J Comput Assist Tomogr* 1995;19:532-6.
- 106 Blasberg RG. Prediction of brain tumor therapy response by PET [review]. *J Neurooncol* 1994;22:281-6.
- 107 Tjuvajev J, Muraki A, Ginos J, *et al.* Iododeoxyuridine uptake and retention as a measure of tumor growth [published erratum appears in *J Nucl Med* 1993;34:1645]. *J Nucl Med* 1993;34:1152-62.
- 108 Black KL, Mazziotta JC, Becker DP. Brain tumors [review]. *West J Med* 1991;154:186-97.
- 109 Pappata S, Cornu P, Samson Y, *et al.* PET study of carbon-11-PK 11195 binding to peripheral type benzodiazepine sites in glioblastoma: a case report. *J Nucl Med* 1991;32:1608-10.
- 110 Black KL, Ikezaki K, Santori E, Becker DP, Vinters HV. Specific high-affinity binding of peripheral benzodiazepine receptor ligands to brain tumors in rat and man. *Cancer* 1990;65:93-7.
- 111 Mineura K, Yasuda T, Kowada M, Ogawa T, Shishido F, Uemura K. Positron emission tomographic evaluation of radiochemotherapeutic effect on regional cerebral hemocirculation and metabolism in patients with gliomas. *J Neurooncol* 1987;5:277-85.
- 112 Rozental JM, Levine RL, Nickles RJ, Dobkin JA. Glucose uptake by gliomas after treatment. A positron emission tomographic study [see comments]. *Arch Neurol* 1989;46:1302-7.
- 113 Ogawa T, Uemura K, Shishido F, *et al.* Changes of cerebral blood flow, and oxygen and glucose metabolism following radiochemotherapy of gliomas: a PET study. *J Comput Assist Tomogr* 1988;12:290-7.
- 114 Rozental JM, Cohen JD, Mehta MP, Levine RL, Hanson JM, Nickles RJ. Acute changes in glucose uptake after treatment: the effects of carmustine (BCNU) on human glioblastoma multiforme. *J Neurooncol* 1993;15:57-66.
- 115 Rozental JM, Levine RL, Nickles RJ. Changes in glucose uptake by malignant gliomas: preliminary study of prognostic significance. *J Neurooncol* 1991;10:75-83.
- 116 Rozental JM, Levine RL, Mehta MP, *et al.* Early changes in tumor metabolism after treatment: the effects of stereotactic radiotherapy. *Int J Radiat Oncol Biol Phys* 1991;20:1053-60.
- 117 Wurker M, Herholz K, Voges J, *et al.* Glucose consumption and methionine uptake in low-grade gliomas after iodine-125 brachytherapy. *Eur J Nucl Med* 1996;23:583-6.
- 118 Takeda N, Diksic M, Yamamoto YL. The sequential changes in DNA synthesis, glucose utilization, protein synthesis, and peripheral benzodiazepine receptor density in C6 brain tumors after chemotherapy to predict the response of tumors to chemotherapy. *Cancer* 1996;77:1167-79.
- 119 Le Bihan D, Breton E, Lattmand D, Rubin ML, Vignand J, Laval-Jeantet M. Separation of diffusion and perfusion in intravoxel incoherent motion imaging. *Radiology* 1988;168:497-505.
- 120 Merholdt KD, Bruhn K, Frahm J, Gyngell ML, Hanicke W, Deimling M. MRI of "diffusion" in human brain: new results using a modified CE-FAST sequence. *Magn Reson Med* 1989;9:423-9.
- 121 Cairncross JG, Pexham JHW, Rathbone MP, DelMaestro RF. Postoperative contrast enhancement in patients with brain tumour. *Ann Neurol* 1985;17:570-2.
- 122 Dooms GC, Hecht S, Brant-Zawadzki M. Brain radiation lesions: MR imaging. *Radiology* 1986;158:149-55.
- 123 Patronas NJ, Di Chiro G, Brooks RA, *et al.* Work in progress: [^{18}F] fluorodeoxyglucose and positron emission tomography in the evaluation of radiation necrosis of the brain. *Radiology* 1982;144:885-9.
- 124 Coleman RE, Hoffman JM, Hanson MW, Sostman HD, Schold SC. Clinical application of PET for the evaluation of brain tumors [review]. *J Nucl Med* 1991;32:616-22.

- 125 Glantz MJ, Hoffman JM, Coleman RE, *et al.* Identification of early recurrence of primary central nervous system tumors by [18F]fluorodeoxyglucose positron emission tomography [see comments]. *Ann Neurol* 1991;29:347–55.
- 126 Di Chiro G, Oldfield E, Wright DC, *et al.* Cerebral necrosis after radiotherapy and/or intra-arterial chemotherapy for brain tumors: PET and neuropathologic studies. *AJR Am J Roentgenol* 1988;150:189–97.
- 127 Valk PE, Budinger TF, Levin VA, Silver P, Gutin PH, Doyle WK. PET of malignant cerebral tumors after interstitial brachytherapy. Demonstration of metabolic activity and correlation with clinical outcome. *J Neurosurg* 1988;69:830–8.
- 128 Doyle WK, Budinger TF, Valk PE, Levin VA, Gutin PH. Differentiation of cerebral radiation necrosis from tumor recurrence by [18F]FDG and 82Rb positron emission tomography. *J Comput Assist Tomogr* 1987;11:563–70.
- 129 Ogawa T, Kanno I, Shishido F, *et al.* Clinical value of PET with 18F-fluorodeoxyglucose and L-methyl-¹¹C-methionine for diagnosis of recurrent brain tumor and radiation injury. *Acta Radiol* 1991;32:197–202.
- 130 Phillips PC, Moeller JR, Sidis JJ, *et al.* Abnormal cerebral glucose metabolism in long-term survivors of childhood acute lymphocytic leukemia. *Ann Neurol* 1991;29:263–71.
- 131 Mitsuki S, Diksic M, Conway T, Yamamoto YL, Villemure JG, Feindel W. Pharmacokinetics of ¹¹C-labelled BCNU and SarCNU in gliomas studied by PET. *J Neurooncol* 1991;10:47–55.
- 132 Tyler JL, Yamamoto YL, Diksic M, *et al.* Pharmacokinetics of superselective intra-arterial and intravenous [11C]BCNU evaluated by PET. *J Nucl Med* 1986;27:775–80.
- 133 Daghighian F, Pentlow KS, Larson SM, *et al.* Development of a method to measure kinetics of radiolabelled monoclonal antibody in human tumour with applications to microdosimetry: positron emission tomography studies of iodine-124 labelled 3F8 monoclonal antibody in glioma. *Eur J Nucl Med* 1993;20:402–9.
- 134 Barnes PD, OTuama L, Tzika A. Investigating the pediatric central nervous system. *Curr Opin Pediatr* 1993;5:643–52.
- 135 Bergh J, Nilsson S, Liljedahl C, *et al.* In vivo imaging and treatment of human brain tumours utilizing the radiolabelled monoclonal antibody MUC 2–63. *Anticancer Res* 1990;10:655–60.