ARCHIVES OF DISEASE IN CHILDHOOD

The Journal of the Royal College of Paediatrics and Child Health

This issue live and online
www.archdischild.com

Annotations

Prognosis for vesicoureteric reflux

The prevalence of vesicoureteric reflux (VUR) has been estimated to be 2% of the child population.¹ In children with VUR demonstrated on micturating cystourethrography there is a tendency for the grade of VUR to improve or for VUR to disappear with time and with increasing age.²⁻³ VUR has been identified as a risk factor for the development of urinary tract infections (UTI) and is present in a third of young children presenting with this problem. In addition, it is a risk factor for renal scarring, otherwise called reflux nephropathy.⁴⁻⁵ VUR is also associated with renal dysplasia and other developmental abnormalities of the urinary tract.⁶ There is now abundant evidence for inheritance by an autosomal dominant mechanism.⁷

Pathogenesis of reflux nephropathy

Studies have suggested that reflux nephropathy develops following UTI in very early childhood or infancy.⁸ New scars have been observed relatively infrequently; however, there are sufficient case reports of new scar formation both on intravenous urography and using ^{99m}Tc DMSA scans to accept that at least a proportion of renal scars are acquired following UTI.⁹⁻¹⁰ The probability that most scars develop in this way cannot be proved because relatively few children have serial imaging studies; in particular, few children have had imaging investigations before the first UTI. The link between UTI, VUR, and renal scarring has been confirmed by several independent groups.⁴⁻¹¹ Smellie *et al* have also demonstrated a link between delay in diagnosis and treatment of UTI and the development of new renal scars.⁹⁻¹²

Symptoms and signs of VUR and reflux nephropathy

VUR and reflux nephropathy are silent conditions that do not usually give rise to symptoms or signs except when complications such as UTI develop. They can only be detected by invasive tests that are not routinely carried out in healthy children and are not usually indicated during the acute phase of treatment. Thus knowledge of pathogenesis and natural history has been gleaned from observational studies and additional imaging investigations carried out on the advice of paediatricians in the belief that they are important for management.¹³ The evolution of reflux nephropathy is a slow process. Most renal scarring develops very early in childhood, but progressive deterioration of damaged kidneys may continue slowly throughout life. The relative contributions of congenital renal dysplasia, acquired reflux nephropathy, and the final common pathway of progressive glomerulosclerosis¹⁴⁻¹⁵ are difficult

to disentangle in this group of patients. The development of proteinuria is indicative of progressive glomerulosclerosis and is a bad prognostic feature particularly when the patient also has hypertension.

Historical perspective

A review of literature in the preantibiotic era suggests that chronic pyelonephritis was a very serious condition in children and adults. Weiss and Parker described a series of postmortem cases¹⁶; antecedent clinical features included recurrent fevers, presumably due to persistent untreated infection, anaemia, hypertension, growth failure, and pregnancy complications. There is evidence for a falling prevalence of this condition, which is probably due to a true reduction of reflux nephropathy because of modern medical care, particularly the treatment of acute pyelonephritis with antibiotics; alternatively the decline may represent changing fashions in disease classification. The historical aspects are discussed in detail by MacGregor¹⁷ who considered that VUR was crucial to the development of reflux nephropathy.

Long term outcome of VUR

In her elegant long term study Smellie describes the natural history of VUR in 226 children, 85 of whom had renal scarring modified by close medical supervision, including long term, low dose prophylaxis, advice on double micturition, treatment of intercurrent UTI, and management of hypertension.¹⁸ In addition 33 patients had surgical procedures (nephroureterectomy or reimplantation of the ureter, or both). Twenty (9%) had hypertension and six (4%) had chronic renal failure, two of whom reached end stage. New scars visible on intravenous urography developed in only a small proportion of the cohort and this was attributed to the benefits of careful medical supervision. The incidence of new scars is lower than in other studies,¹⁹⁻²¹ but the diagnosis of new scars is imprecise and subjective so that comparisons between studies are difficult. New scars have only rarely been seen to develop after the age of 4 to 5 years.²² If scars are acquired following UTI the prompt treatment of symptomatic episodes particularly in early childhood may be equally or even more important than long term preventative measures. The natural history of asymptomatic bacteriuria is not significantly different in terms of incidence of new scars, although many girls have prolonged periods of exposure to infection.¹⁹⁻²³ The long gestation period for scar formation visible on intravenous urography makes it difficult to link new scars to specific episodes of infection.

Surgery for prevention of UTI and reflux nephropathy

Following the original observations by Hodson and Edwards²⁴ of the strong association between UTI, VUR, and reflux nephropathy, surgeons developed operations to correct VUR in the belief that this would reduce the risk of recurrent UTI and prevent the development of reflux nephropathy.²⁵ Although reimplantation of the ureter was quite successful for elimination of VUR, there was a significant complication rate²⁶ and the overall risk of UTI was altered little by successful surgery.^{20 21 27} Similarly, successful surgery did not benefit glomerular filtration rate or prevent new scar formation. More recently a procedure involving the suburothelial injection of Teflon or collagen has been used to correct VUR endoscopically.²⁸ The success rate of this procedure in eliminating VUR is less than for surgery but the procedure is simpler and involves a shorter hospital stay. Long term benefits from this procedure on infection rates and new scar formation have not been evaluated. Teflon has now been abandoned because of concerns about embolisation of particles causing granulomas at distant sites; collagen has been found to be safe but benefits are not permanent and VUR may recur after a period of months or years.

Long term, low dose prophylaxis

An alternative approach to correction of VUR is the use of long term, low dose prophylaxis initially with cotrimoxazole or nitrofurantoin²⁹ and later using trimethoprim. It was postulated that UTI usually develop as a result of ascending infection and that from time to time bacteria ascend the urethra and establish infection in the bladder or kidneys. A nightly dose of a broad spectrum antibiotic can sterilise the urine on a daily basis if the bacteria are sensitive to the drug chosen, reducing the chance that UTI can become established. Breakthrough infections can occur if the child's gut and perineum are colonised with resistant organisms. Breakthrough with sensitive organisms suggests that prophylaxis has been omitted. There was evidence of effectiveness of this treatment in reducing the rate of reinfection in children with normal urinary tracts,³⁰ but there are no controlled studies demonstrating that long term, low dose prophylaxis is superior to prompt treatment of UTI for the prevention of renal scarring. Thus the basis for widespread use of long term low, dose prophylaxis for the prevention of scarring is based on a hypothesis that has never been proved. Increasing resistance of organisms to trimethoprim raise doubts about effectiveness as a prophylactic agent in 1999. Nitrofurantoin is bitter tasting and less well tolerated than trimethoprim but the development of resistance is uncommon.

Guidelines

Because of the concern about development of reflux nephropathy, the Royal College of Physicians (RCP) published guidelines in 1991 on the diagnosis of UTI in childhood and recommended that imaging investigations should be carried out in every child following the first UTI,³¹ reflecting the view stated in the *Lancet* 20 years previously.¹³ The working group advised that all children should have an ultrasound examination and that a ^{99m}Tc DMSA scan should be carried out on all those younger than 7 years. Micturating cystourethrography was considered mandatory in children younger than 1 year. The indication for these tests was the high risk of VUR, around 30%, and the presumed risk of renal damage in these age groups. However, in the absence of good evidence for benefit from treatments and evidence of poor compliance with long term, low dose prophylaxis,³² the benefits of imaging and long term treatment may not be as great as originally expected.

The imaging investigations recommended in the RCP guidelines are in effect screening tests as they are carried out in a high risk population after the acute illness has been treated successfully, in the hope that better knowledge of the underlying anatomy will improve future management and prognosis. Unfortunately, there is no evidence that the prognosis is altered by these tests or by the widespread use of long term, low dose prophylaxis or successful surgery. However, the use of published guidelines has provided a very useful step in the evaluation of current practice, as the guidelines are followed widely throughout the UK and it is now possible to audit and assess the outcome of this standard practice.³³ Published studies suggest that many children do not have the recommended management.³⁴

Stark, in 1997, challenged the view that these imaging tests are worthwhile and suggested that they are excessively costly and invasive without giving any benefit to most children.³⁵ Tertiary specialists were heavily represented in the working group of the research unit of the RCP; their views and experiences represented the most severe end of the clinical spectrum of UTI and reflux nephropathy. In contrast, general practitioners and general paediatricians who see most children at the time of their first UTI were in the minority. Although there was no formal attempt to inform general practitioners of the guidelines some general practices are now referring large numbers of children for imaging investigations and a paediatric opinion following simple, non-febrile UTI. This has generated a massive workload for radiology departments and exposed large numbers of children to significant radiation without much evidence of benefit. The proportion of children seen with evidence of renal damage is lower than in earlier studies. This may be because UTI are diagnosed quickly in infants when they are referred to hospital,³⁶ or it may reflect the fact that many more straightforward cases are being referred for further investigation. Unfortunately, there is continuing evidence of delay in diagnosis of UTI in infants and toddlers in primary care where urine collection is perceived as difficult,³⁷ and some children have several consultations before the diagnosis of UTI is considered and further delays before it is confirmed.³⁸

Practice in Sweden

In Sweden, children over 2 years old with simple UTI are not referred for imaging procedures unless there is an additional risk factor such as recurrent UTI or evidence of upper tract involvement.³⁹ In spite of this practice, the prevalence of renal scarring is extremely low and there were no cases of chronic renal failure due to reflux nephropathy in a survey published in 1980,^{4 40} in contrast to the high incidence of VUR and reflux nephropathy in children in the UK.⁴¹ This has been attributed to the excellent facilities for the diagnosis of UTI in infants who are screened for UTI whenever they are febrile.⁴² The mean age for diagnosis of UTI in Sweden is during the 1st year of life in contrast to the UK where the mean age for diagnosis is around 4 years.

Need for controlled studies

Smellie and colleagues¹⁸ state that it would be unethical to carry out a randomised controlled study, and this may have been the view of most paediatricians at the time this study was started. However, now that the results of the two large prospective studies have failed to show superiority of either medical or surgical management, and there is mounting evidence that most scars are acquired very early in childhood, the climate of opinion may have changed. There is increasing pressure to establish the effectiveness of treatments and screening procedures both in the interest of the individual patient and in the interest of the National Health Service as a whole. Mounting costs of new and

sophisticated treatments have focused the attention of clinicians, managers, and politicians on the need to ensure that money is well spent. The public is becoming aware of the adverse effect of widespread use of antibiotics on bacterial resistance and of the small but not negligible risk of side effects of drugs used long term. The newly established National Institute of Clinical Excellence will examine new treatments, but it is unlikely that it will be able to examine established practice in retrospect. The use of systematic reviews to provide busy clinicians with a summary of previous studies has also aided critical evaluation of current practice.⁴³ The outcome of the systematic review of management of VUR is awaited with interest.

KATE VERRIER JONES

Laura Ashley Reader in Paediatric Nephrology,
KRUF Children's Kidney Centre for Wales,
Department of Child Health, University of Wales College of Medicine,
Heath Park, Cardiff CF14 4XN, UK
email: verrier-jones@cf.ac.uk

- Report of a meeting of physicians at the Hospital for Sick Children GOS London, Vesicoureteric reflux: all in the genes? *Lancet* 1996;**348**:725–8.
- Edwards D, Normand ICS, Prescod N, Smellie J. Disappearance of vesicoureteric reflux during long-term prophylaxis of urinary tract infection in children. *BMJ* 1977;ii:285–8.
- Smellie JM. The disappearance of reflux in children with urinary tract infection during prophylactic chemotherapy. In: Alwall N, Berglund F, Josephson B, eds. Proceedings of the 4th International Congress of Nephrology, Stockholm 1969, Vol 3. *Clinical nephrology, immunology*. Basel: Karger, 1969:357.
- Winberg J, Bollgren I, Kallenius G, Mollby R, Svenson SB. Clinical pyelonephritis and focal renal scarring. A selected review of pathogenesis, prevention and prognosis. *Paediatr Clin North Am* 1982;**29**:801–14.
- Hodson CJ. The kidneys in urinary tract infection. *Proc R Soc Med* 1966;**59**:416.
- Risdon RA. The small scarred kidney in childhood. *Pediatr Nephrol* 1993;**7**:361–4.
- Kenda RB, Fettich JJ. Vesicoureteric reflux and renal scars in asymptomatic siblings of children with reflux. *Arch Dis Child* 1992;**67**:506–8.
- Berg UB, Johansson SB. Age as a main determinant of renal functional damage in urinary tract infection. *Arch Dis Child* 1983;**58**:963–69.
- Smellie JM, Ransley PG, Normand ICS, Prescod N, Edwards D. Development of new renal scars; a collaborative study. *BMJ* 1985;**290**:1957–60.
- Vernon SJ, Coulthard MG, Lambert HJ, Keir MJ, Matthews JNS. New renal scarring in children who at age 3 and 4 years had had normal scans with dimercaptosuccinic acid: follow up study. *BMJ* 1997;**315**:905–8.
- McKerrow W, Davidson-Lamb N, Jones PF. Urinary tract infection in children. *BMJ* 1984;**289**:299–304.
- Smellie JM, Poulton A, Prescod NP. Retrospective study of children with renal scarring associated with reflux and urinary infection. *BMJ* 1994;**308**:1193–6.
- Anon. Vesico-ureteric reflux [editorial]. *Lancet* 1968;ii:1072–74.
- Becker GJ, Kincaid-Smith P. Reflux nephropathy: the glomerular lesion and progression of renal failure. *Pediatr Nephrol* 1993;**7**:365–9.
- Risdon RA. The small scarred kidney of childhood. A congenital or an acquired lesion? *Pediatr Nephrol* 1987;**1**:632.
- Weiss S, Parker F. Pyelonephritis: it's relation to vascular lesions and to arterial hypertension. *Medicine* 1939;**18**:221–315.
- MacGregor M. Pyelonephritis lenta. Consideration of childhood urinary infection as the forerunner of renal insufficiency in later life. *Arch Dis Child* 1970;**45**:159–72.
- Smellie JM, Prescod NP, Shaw PJ, Risdon RA, Bryant TN. Childhood reflux and urinary infection: a follow-up of 10–41 years in 226 adults. *Pediatr Nephrol* 1998;**12**:727–36.
- Aggarwal VK, Verrier Jones K, Asscher AW, Evans C, Williams LA. Covert bacteriuria: long term follow-up. *Arch Dis Child* 1991;**66**:1284–6.
- Birmingham Reflux Study Group. Prospective trial of operative versus non-operative treatment of severe vesicoureteric reflux in children: five years' observation. *BMJ* 1987;**295**:237–41.
- International Reflux Study Committee. Five-year study of medical or surgical treatment in children with severe reflux: radiological renal findings. *Pediatr Nephrol* 1992;**6**:223–30.
- Coulthard MG, Lambert HJ, Keir MJ. Occurrence of renal scars in children after their first referral for urinary tract infection. *BMJ* 1997;**315**:918–19.
- Cardiff-Oxford Bacteriuria Study Group. Sequelae of covert bacteriuria in schoolgirls. *Lancet* 1978;ii:889–93.
- Hodson CJ, Edwards D. Chronic pyelonephritis and vesicoureteric reflux. *Clin Radiol* 1960;**2**:219–31.
- Politano VA, Leadbetter WF. An operative technique for the correction of V.U.R. *J Urol* 1958;**79**:932.
- O'Donnell B. Controversies in therapeutics. Management of urinary tract infection and vesicoureteric reflux in children. The case for surgery. *BMJ* 1990;**300**:1393–4.
- Winberg J. Management of primary vesico-ureteric reflux in children—operation ineffective in preventing progressive renal damage. *Infection* 1994;**22**(suppl 1):S4–7.
- O'Donnell B, Puri P. Endoscopic correction of primary vesicoureteric reflux: results in 94 ureters. *BMJ* 1986;ii:1404–6.
- Normand ICS, Smellie JM. Prolonged maintenance chemotherapy in the management of urinary infection in childhood. *BMJ* 1965;ii:1023–6.
- Smellie JM, Katz G, Gruneberg RN. Controlled trial of prophylactic treatment in childhood urinary tract infection. *Lancet* 1978;ii:175–8.
- Report of a Working Group of the Research Unit Royal College of Physicians. Guidelines for the management of acute urinary tract infection in childhood. *J R Coll Physicians Lond* 1991;**25**:36–42.
- Smyth AR, Judd BA. Compliance with antibiotic prophylaxis in urinary tract infection. *Arch Dis Child* 1993;**68**:235–6.
- Jadresic L, Cartwright K, Cowie N, Witcombe B, Stevens D. Investigation of urinary tract infection in childhood. *BMJ* 1993;**307**:761–4.
- South Bedfordshire Practitioners' Group. Development of renal scars in children: missed opportunities in management. *BMJ* 1990;**301**:1082–7.
- Stark H. Urinary tract infections in girls: the cost-effectiveness of currently recommended investigative routines. *Pediatr Nephrol* 1997;**11**:174–7.
- Verrier Jones K, Hockley B, Pollock J, Scrivener R. Diagnosis of urinary tract infections in children under two years: a national audit [abstract]. *Pediatr Nephrol* 1998;**12**:C128.
- van der Voort J, Edwards A, Roberts R, Verrier Jones K. The struggle to diagnose UTI in children under two in primary care. *Family Practice* 1997;**14**:44–8.
- van der Voort J, Verrier Jones K, Edwards A. Can renal scarring be prevented by earlier diagnosis of urinary tract infection in infancy? [abstract] *Kidney Int* 1997;**52**:1130.
- Jodal U, Winberg J. Management of children with unobstructed urinary tract infection. *Pediatr Nephrol* 1987;**1**:647–56.
- Helin I, Winberg J. Chronic renal failure in Swedish children. *Acta Paediatr Scand* 1980;**69**:607–611.
- United Kingdom Transplant Support Service Authority. *Renal transplant audit 1984–1993*. Bristol: UKTSSA, 1995.
- Winberg J. Clinical aspects of urinary tract infection. In: Barratt TM, Holliday MA, Vernier RL, eds. *Paediatric nephrology*. 2nd ed. Baltimore: Williams & Wilkins, 1987:626–46.
- Chalmers I, Altman DG, eds. *Systematic reviews*. London: BMJ Publishing Group, 1995.

Cardiac causes for syncope or sudden death in childhood

Sudden death of a child is every parent's worst nightmare. It is a rare occurrence, although individual cases or causes may receive disproportionate attention in the press. Syncope, on the other hand, is a common symptom in childhood and is almost always benign. In a very few cases syncope may be a clue to the presence of an underlying cardiovascular problem and may predict a risk of sudden death. Retrospective assessment of victims of sudden death sometimes reveals a preceding history of syncope to suggest that earlier recognition of the problem might have led to a different outcome. This review considers the causes of sudden death, the causes of syncope, and which children with syncope need further investigation.

Cardiac causes of sudden death in infancy

Sudden death in infancy is usually either sudden infant death syndrome or caused by infection. A few neonatal or

infant deaths are caused by congenital cardiovascular malformations, particularly duct dependent abnormalities or obstructive left heart malformations.¹ Sustained tachycardia usually presents with heart failure but may cause collapse or even death. Atrioventricular re-entry tachycardia, the most common type of supraventricular tachycardia, maybe difficult to recognise in the neonate. The history is often non-specific and there are few physical signs. Ventricular arrhythmias in infancy are rare but some types are dangerous. Complete atrioventricular block is usually recognised in utero or soon after birth but may cause death if unrecognised or untreated.

Sudden infant death syndrome implies absence of any identifiable cause of death. Extensive investigation has failed to demonstrate any cardiovascular contribution to sudden infant death syndrome² and, despite recent reports, there is no convincing evidence that QT prolongation is implicated.³

Sudden death in children with known heart disease

In previous generations sudden cardiac death most often occurred in children with end stage pulmonary vascular disease associated with unoperated congenital heart disease or in children with unoperated aortic valve stenosis.⁴ Over the years surgical repair has been performed earlier and more effectively so that those most at risk now are children with repaired heart disease. The risk seems to be highest (5–7 per 1000 patient years' follow up) after atrial repair of transposition of the great arteries.^{5,6} Death is probably caused by an arrhythmia and there is circumstantial evidence to implicate atrial flutter⁶ although ventricular tachycardia or ventricular fibrillation are also candidates.⁵ The risk of sudden death after repair of tetralogy of Fallot is lower (around 1.5 per 1000 patient years) and is declining.⁵ Death is probably caused by a ventricular arrhythmia, perhaps in combination with a residual haemodynamic abnormality, but late atrioventricular block is also a possible cause. Late sudden death after repair of other common cardiovascular malformations is rare, with an incidence of around 0.1 per 1000 patient years.⁵

The risk of sudden death in the presence of known hypertrophic cardiomyopathy is impossible to establish because of the lack of a denominator. The impression given by published reports from specialised referral units is that the risk is high, whereas evidence from the few population based studies does not support this. "Risk factors" have been identified but are absent in most children. "Risk stratification" is encouraged and may identify a few individuals at increased risk. However, for the large majority the risk of sudden death is low, especially in the absence of symptoms.

Syncope in children with known heart disease

Syncope or presyncope in any child after cardiac surgery or with unoperated heart disease is an obvious indication for urgent specialist evaluation. It is to be hoped that all such children are under review by a paediatric cardiologist.

Sudden death in apparently normal children

The incidence of sudden death in apparently normal children is around 1–1.5 per 100 000 per year. Cardiovascular diagnoses first established after death include myocarditis, various types of cardiomyopathy, coronary artery abnormalities, and Marfan syndrome with aortic dissection. Sudden death associated with some types of structural heart disease, especially myocarditis or cardiomyopathy, may be related to an arrhythmia. In other children, death may remain unexplained after necropsy but the circumstances may lead to a presumptive diagnosis of a primary cardiac arrhythmia.

Sudden death from previously unsuspected hypertrophic cardiomyopathy often attracts publicity but is rare. The risk in apparently normal children or adolescents is less than 1 per 1 000 000 per year.⁷ Postmortem diagnosis should lead to screening of other family members but often no further cases are identified. Hypertrophic cardiomyopathy has an underlying genetic cause and it is likely that many cases presenting in this way are new mutations. Retrospective assessment will often reveal a prodromal history of syncope.^{8,9}

Unexpected sudden deaths that remain unexplained after necropsy are probably due to primary cardiac arrhythmias and are about 10 times as common as deaths from hypertrophic cardiomyopathy. Potentially fatal arrhythmias that would lead no trace after death include polymorphic ventricular tachycardia in congenital long QT syndrome,¹⁰ atrial fibrillation in Wolff-Parkinson-White syndrome,¹¹ primary ventricular arrhythmias such as those described by Brugada and colleagues¹² or Leenhardt *et al.*,¹³ and congenital complete atrioventricular block.¹⁴ Retrospective confirmation of an arrhythmia is not possible unless a familial

condition is recognised in a family member, but the circumstances surrounding death support a presumptive diagnosis in many cases. This group of arrhythmias also comprise most of the diagnoses made after resuscitation from out-of-hospital cardiac arrest.¹⁵ It is possible that some deaths attributed to "epilepsy" are in fact caused by a primary arrhythmia. Differentiation between syncope from a cardiac arrhythmia and epilepsy can be difficult.¹⁶

Syncope in apparently normal children

Syncope is a frequent problem in childhood. The common causes are benign whereas some rare causes are potentially dangerous. A simple faint or vasovagal syncope is said to occur in up to 15% of the normal population at some time during childhood. Other vascular but benign diagnoses include neurocardiogenic syncope in older children,¹⁷ and pallid syncope or reflex anoxic seizures in younger children.¹⁸ The mechanisms of these may be similar with inappropriate reflex bradycardia and hypotension. In the former syncope is often postural, and in the latter a frequent trigger is surprise, minor hurt or frustration.

Hyperventilation related to anxiety or behavioural disorders may produce syncope resulting from cerebral hypoperfusion, and the diagnosis will usually be apparent from the history.¹⁹ Psychogenic or hysterical syncope is rare—it usually occurs with an audience and does not result in injury.¹⁹

Studies that examine children presenting with syncope rarely identify any diagnoses with a significant risk of death because they are rare.²⁰ The spectrum of underlying abnormalities in children resuscitated from cardiac arrest is, not surprisingly, very different and more malignant. Diagnoses include various types of cardiomyopathy, particularly dilated cardiomyopathy or arrhythmogenic right ventricular cardiomyopathy, and substrates for primary arrhythmias including QT prolongation and ventricular pre-excitation.¹⁵

INVESTIGATION OF SYNCOPES

The history is of fundamental importance in the assessment of syncope. Common but benign causes such as vasovagal syncope or hyperventilation are usually easily recognised. A history of syncope on exertion, syncope preceded by palpitations, or syncope in the presence of a family history of sudden death, congenital long QT syndrome, or hypertrophic cardiomyopathy should lead to specialist evaluation. Detection of a cardiac murmur (or other abnormal signs on examination) should also lead to specialist referral as it may be a clue to the presence of aortic valve stenosis, coarctation of the aorta, or hypertrophic cardiomyopathy.

Electrocardiography is a valuable screening test in children with syncope, and may identify QT prolongation, ventricular pre-excitation, or atrioventricular block or it may be a clue to the presence of underlying ventricular arrhythmia or cardiomyopathy. Other investigations might include echocardiography, ambulatory or exercise electrocardiography, tilt testing,²¹ or invasive electrophysiology, but the investigative strategy depends fundamentally on the history.⁹

Tilt testing may be valuable in assessment of neurally mediated syncope with reasonable sensitivity and specificity.²¹ Treatment strategies include β blockade, vasoconstrictors, fludrocortisone, and even pacemaker implantation, but their efficacy can be difficult to evaluate and support from convincing controlled trials is lacking.¹⁷

Sudden death in young athletes

A small but significant proportion of sudden deaths occur during exercise. Deaths during sports attract publicity and focus attention on the possibility of an underlying cardiac abnormality. The common structural abnormalities

identified at necropsy are hypertrophic cardiomyopathy, coronary artery abnormalities, right ventricular cardiomyopathy, and aortic valve stenosis.^{9–22} Unexplained cases may result from ventricular arrhythmia or atrial fibrillation in Wolff-Parkinson-White syndrome. Health screening programmes are in place for some young athletes but their efficacy has not been evaluated.

Death during sports is a rare event: estimates from the USA suggest that 5 per 100 000 young athletes (including children, teenagers, and young adults) have a predisposing condition, and that only 10–25 such deaths per year occur among the total USA population of 260 000 000.⁹

Population screening for unsuspected heart disease

The infrequency of prodromal symptoms and the absence of physical signs in children who die suddenly and are found to have a cardiac problem may lead to calls for screening of the general population. There is no evidence that screening for hypertrophic cardiomyopathy is either feasible or appropriate, and no evidence that either early detection or treatment in the absence of symptoms has any effect on outcome.^{23–24} To my knowledge the only prospective screening programme of normal children has been in place in Tokyo, Japan: the prevalence of hypertrophic cardiomyopathy is around 1 in 15 000,²⁵ and children identified seemed to be at particularly low risk.²⁶ The programme does also detect children with long QT syndrome, atrioventricular block, and Wolff-Parkinson-White syndrome by routine electrocardiography but these problems are rare. So far it has not been possible to show that screening prevents sudden death.²⁷

Conclusions

Sudden death in children is rare. About 10% of all deaths are sudden but most occur in children with previously recognised disease. Syncope is very common but is only rarely a clue to the presence of a significant cardiovascular abnormality. A detailed history is of paramount importance and the ECG is a useful screening test. There is no evidence at present to support screening of all children for unsuspected cardiac problems.

CHRISTOPHER WREN

Consultant Paediatric Cardiologist,
Freeman Hospital, Newcastle upon Tyne NE7 7DN, UK

Covert video surveillance: an important investigative tool or a breach of trust?

There has been controversy among professionals and the public surrounding the use of covert video surveillance (CVS) in cases of suspected child abuse. Opinion is divided as to whether it is necessary to make a “diagnosis” of child abuse and whether it is ethical and legal. CVS for documentation of apnoea by smothering was first used in 1983,¹ and since then more than 20 reports have appeared.^{2–7} Its primary use is to detect imposed upper airways obstruction, which normally presents in small babies under the age of 1 year with an apparent life threatening event. It is a well recognised entity requiring immediate and urgent assessment. It is a dangerous form of abuse with high mortality and morbidity.⁸ It can be difficult to differentiate a “normal” apparent life threatening event from an imposed upper airways obstruction purely on clinical grounds. It may require multichannel monitoring of physio-

- 1 Abu-Harb M, Hey E, Wren C. Death in infancy from unrecognised congenital heart disease. *Arch Dis Child* 1994;71:3–7.
- 2 Guntheroth WG. Theories of cardiovascular causes of sudden infant death syndrome. *J Am Coll Cardiol* 1989;14:443–7.
- 3 Schwartz PJ, Stramba-Dadiale M, Segantini A, et al. Prolongation of the QT interval and the sudden infant death syndrome. *N Engl J Med* 1998;338:1709–14.
- 4 Lambert EC, Menon VA, Wagner HR, Vlad P. Sudden unexpected death from cardiovascular disease in children. *Am J Cardiol* 1974;34:89–96.
- 5 Silka MJ, Hardy BG, Menashe VD, Morris CD. A population-based prospective evaluation of risk of sudden cardiac death after operation for common congenital heart defects. *J Am Coll Cardiol* 1998;32:245–51.
- 6 Wren C. Late postoperative arrhythmias. In: Wren C, Campbell RWFC, eds. *Paediatric cardiac arrhythmias*. Oxford: Oxford University Press, 1996: 238–59.
- 7 Arola A, Jokinen E, Ruuskanen O, et al. Epidemiology of idiopathic cardiomyopathies in children and adolescents. A nationwide study in Finland. *Am J Epidemiol* 1997;146:385–93.
- 8 Driscoll DJ, Edwards WD. Sudden unexpected death in children and adolescents. *J Am Coll Cardiol* 1985;5:118B–21B.
- 9 Liberton RR. Sudden death from cardiac causes in children and young adults. *N Engl J Med* 1996;334:1039–44.
- 10 Garson A Jr, Dick M, Fournier A, et al. The long QT syndrome children. An international study of 287 patients. *Circulation* 1993;87:1866–72.
- 11 Timmermans C, Smeets JL, Rodriguez LM, Vrochous G, van den Dool A, Wellens HJ. Aborted sudden death in the Wolff-Parkinson-White syndrome. *Am J Cardiol* 1995;115:492–4.
- 12 Brugada J, Brugada R, Brugada P. Right bundle-branch block and ST-segment elevation in leads V1 through V3. *Circulation* 1998;97:457–60.
- 13 Leenhardt A, Lucet V, Denjoy I, Grau F, Ngoc DD, Coumel P. Catecholaminergic polymorphic ventricular tachycardia in children. *Circulation* 1995;91:1512–19.
- 14 Ross BA. Congenital complete atrioventricular block. *Pediatr Clin North Am* 1990;37:67–78.
- 15 Silka MJ, Kron J, Walence CG, Cutler JE, McAnulty JH. Assessment and follow-up of pediatric survivors of sudden cardiac death. *Circulation* 1990; 82:341–9.
- 16 Linzer M, Grubb BP, Ho S, Ramakrishnan L, Bramfield E, Estes NA. Cardiovascular causes of loss of consciousness in patients with presumed epilepsy: a cause of the increased sudden death rate in people with epilepsy? *Am J Med* 1994;96:146–54.
- 17 Benditt DG. Neurally mediated syncopal syndromes: pathophysiological concepts and clinical evaluation. *PACE* 1997;20:572–84.
- 18 Brenningstall GN. Breath-holding spells. *Pediatr Neurol* 1996;14:91–7.
- 19 Feit LR. Syncope in the pediatric patient: diagnosis, pathophysiology, and treatment. *Adv Pediatr* 1996;43:469–94.
- 20 Driscoll DJ, Jacobsen SJ, Porter CJ, Wollan PC. Syncope in children and adolescents. *J Am Coll Cardiol* 1997;29:1039–45.
- 21 Benditt DG, Ferguson DW, Grubb BP, et al. Tilt table testing for assessing syncope. *J Am Coll Cardiol* 1996;28:263–75.
- 22 Maron BJ, Shirani J, Poliac LC, Mathenge R, Roberts WC, Mueller FO. Sudden death in young competitive athletes. *JAMA* 1996;276:199–204.
- 23 Clark AL, Coats AJS. Screening for hypertrophic cardiomyopathy. *BMJ* 1993;306:409–10.
- 24 Spirito P, Seidman CE, McKenna WJ, Maron BJ. The management of hypertrophic cardiomyopathy. *N Engl J Med* 1997;336:775–85.
- 25 Ino T, Yabuta K, Yamauchi K. Heart disease screening in Japanese children [letter]. *BMJ* 1993;306:1128.
- 26 Ino T, Okubo M, Nishimoto K, Akimoto K, Yabuta K, Okada R. Clinicopathologic characteristics of hypertrophic cardiomyopathy detected during mass screening for heart disease. *Pediatr Cardiol* 1996;17:295–300.
- 27 Corrado D, Basso C, Schiavon M, Thiene G. Screening for hypertrophic cardiomyopathy in young athletes. *N Engl J Med* 1998;339:364–9.

logical function with concurrent CVS in an environment where prompt action can be taken to protect the child.

What is covert video surveillance?

Covert video surveillance is the video and audio recording of interactions between a child and his or her carer, usually the mother. It is conducted without the knowledge of the carer under strictly controlled conditions within a hospital based paediatric department. It is normally undertaken in a designated cubicle where two cameras (smaller than an electric light switch) are installed to provide coverage of the area from opposing positions. Two cameras reduce the possibility of a single camera angle being inadvertently obscured and abusive behaviour being undetected or misinterpreted. A typical CVS monitoring of a potentially life threatening situation would involve a child under 2 years of

age connected to a multifunction polygraph for clinical monitoring in a cubicle. This helps to maintain a fairly fixed position relative to the camera while giving the child some freedom to roam. The child is monitored from a remote site by a nurse or police officer or both. If a dangerous abusive action by the carer is observed, a separate nurse on the ward will be alerted to intervene. One of the main dilemmas is that the abusive action has to be allowed to continue long enough to obtain conclusive evidence, yet intervention has to occur in sufficient time to prevent harm to the child.

Southall *et al* reported 39 children (2–44 months old) who underwent CVS, of whom 36 were referred for investigation of an apparent life threatening event.⁹ Abuse was revealed in 33 with attempted suffocation in 30 patients.

What are the issues in covert video surveillance?

In 1996 the specialist advisory committee in paediatrics of the [UK] Northern Region was asked to consider the issues surrounding the use of CVS. A multidisciplinary group including professionals from health, social services, and the law produced a report and implementation plan.^{10 11} This group reviewed commonly expressed reservations and objections voiced by both professionals and the public.

ETHICAL AND MORAL CONSIDERATIONS

CVS is not in the best interest of the child

It has been suggested that the Children Act 1989 provides sufficient ways to protect children from abuse without the need for CVS,¹² and that if the child is going to be taken into care anyway such surveillance is unnecessary. However, it is important to recognise that although there may be enough grounds for legal proceedings, courts may make a supervision order, or no order at all, unless an appreciable level of risk is shown, and the child may be returned to the parents to face the same risk. CVS in such situations can provide valuable evidence for both care and criminal proceedings. The rights of the child must be paramount and CVS can provide the vital evidence that will allow protection.

CVS is a betrayal of trust and breach of partnership between parents and paediatricians

Concern has been expressed that CVS involves deception and it is therefore an infringement of the civil liberty of the parent. But this is no different from the current child protection procedures except that the duration of information gathering and arriving at a decision regarding further action may be prolonged, depending on the nature and seriousness of abuse in cases of induced illness syndrome. It has been criticised on the grounds that, “in a civilised society even child abusers have procedural rights”¹³ but the report of the Jasmine Beckford inquiry clearly stated that “parental rights cannot be insisted upon by a parent who has abused these rights”.¹⁴

CVS is a risk to the child

There are concerns that to obtain conclusive evidence for civil or criminal proceedings a decision to intervene (sending a nurse into the cubicle) may be delayed and the child may suffer unnecessary harm. Legally and ethically the principle of “double effect” applies in this situation—that is, an act definable as good in terms of its object can achieve a good effect only at the risk or expense of causing incidental but unavoidable harm.¹⁵ Premature intervention can miss the extent of the abuse and therefore leaves a child and possibly other children at the risk of being returned to the care of an abusing parent and of suffering continued abuse or death in childhood.

CVS contradicts the principle of openness and partnership with parents

The Children Act developed the concept of working in partnership with parents: “The development of working in partnership with parents is usually the most effective route to providing supplementary or substitute care for their children. Measures which antagonise, alienate, undermine or marginalise parents are counter-productive”.¹⁶ Clearly, CVS is in direct conflict with these principles. However in the types of abuse where perpetrators are devious, partnership with parents may need to be curtailed.¹⁷ The suggestion that overt video surveillance of carer and child is likely to succeed is inappropriate. If the carer knows the child is being videoed, he or she will almost certainly modify their behaviour.

CVS may cause possible harm to the parent

If a parent is a perpetrator and is caught on videotape, the diagnosis is certain and appropriate action can be taken to protect the child and to offer treatment to the parent. However, if the parent is not the perpetrator, then they are exonerated. CVS will help to prevent the separation of the child from an innocent parent. It is noteworthy that in the 14 cases published by Samuels *et al*,³ the 12 mothers concerned received non-custodial sentences and psychiatric care as did one grandmother, although the father involved received life imprisonment.

CVS may raise serious concerns among professionals

Professionals may be concerned about undertaking a criminal investigation that takes them away from traditional methods of intervention and treatment. The Royal College of Nursing had serious misgivings about nurses undertaking the role asked of them in the use of CVS,¹⁸ but it has given advice as to the safeguards necessary. CVS should only be implemented after full discussion with all staff who need to be involved, and they must have the right to opt out. The Staffordshire protocol¹⁹ on implementation of CVS was commended by the Department of Health as guidance all authorities should follow if it is to be attempted in any hospital.²⁰

LEGAL CONSIDERATIONS

One of the most important questions to address is whether there are any legal implications of undertaking CVS. It involves a major invasion of privacy and should be conducted within a justifiable legal framework. The Children Act 1989 upholds the principle that the needs of the child are paramount. The addendum to *Working together under the Children Act 1989* gives guidance to doctors stating that the welfare of the child must be of first importance and the overriding principle is to secure the best outcome for the child.²¹ The legal justification is the need to protect the welfare of the child and other children at risk. Health professionals must appreciate evidential issues and the fact that civil and criminal cases might fail due to inadequate evidence. However, it is never justifiable to use CVS simply to seek a criminal conviction. It is not easily dismissed by the court in child care matters, as the protection of the child is paramount. The criticism from courts has often been on the lack of clear guidelines and protocol for its use rather than the use of CVS itself. The usually unequivocal evidence obtained by CVS is admissible and has been accepted.

Two legal reports are noteworthy. In *Regina v Khan* 1996 House of Lords it is stated that “Under English law in general there is nothing unlawful about a breach of privacy”.²² As an occupier of hospital premises an NHS Trust is lawfully entitled to install surveillance equipment. Secondly, in *DH (A Minor) (Child Abuse) 1994* 1 Family

Law Reports 679, Wall J, clear guidance is given that "If a doctor considers that covert video surveillance is essential for the treatment of his patient, the doctor would be entitled to undertake this process without parental consent provided that he is satisfied that there is no risk that the patient will come to any serious harm".²³ In addition, the Home Office guidelines (1994) on the use of equipment in police surveillance²⁴ represent official approval of such covert means by the police in relation to discovery of crime. By analogy, lawful approval can be extended to cover protection of children from serious harm.

There are legal liabilities, particularly in today's climate of increasing numbers of claims against Trusts or health professionals. This needs careful consideration. Claims may be made by parents or carers for breach of confidentiality and trust. They may also potentially be made on behalf of the child on the basis of damage caused by delay of intervention if the child suffers additional harm by the continuation of CVS. Risk is minimised by the use of proper protocols based on professional advice, appropriate training, and experience. Conversely if a child dies or is injured because no CVS was used, and if this could have been proven to be the cause of the injury to the child, a claim could be pursued. In addition, claims based on stress and trauma of staff could be made against Trusts. Generally, protection against such claims will be afforded by approval of CVS at Trust Board level, by appropriate insurance cover by Trusts, and by operation of CVS under an agreed multiagency protocol.

PROFESSIONAL ISSUES

Professional issues concern all groups but are particularly relevant for nurses because of their direct involvement. The philosophy of paediatric nursing has changed in the past 20 to 25 years and has become more family focused and holistic. Consequently, for some, lack of openness with parents can cause considerable ethical dilemmas. The nursing role is crucial, and the principles must be understood and accepted by all involved in its operation. They must be adequately trained and fully supported. Assisting the police with the investigation of serious crime is an accepted exception to the professional duty of confidentiality. Consideration should be given to the concerns of health professionals that they may be disciplined if their activities are considered to be unprofessional or improper. This should be dealt with by appropriate clinical decision making in accordance with a protocol and by reassurance from the individual's professional body, and by operating within a proper multiagency protocol. It is important to note that Trusts have a vicarious liability for all employees acting within the course of their employment under agreed protocols.

There should be a clear agreement that unauthorised notification of CVS to parents and to others will be a disciplinary matter. It could also be a criminal issue if notification has been deemed to interfere with the detection of a crime. Care should be exercised in ensuring that those who volunteer to be involved in CVS are suitable and will be trained. As with any difficult ethical issue, it is wise to involve two independent and appropriate consultant paediatricians from the outset. The inappropriate and maverick use of CVS must be avoided by operating within a strict and rigorous multiagency protocol. There should be a unanimous view that all other alternatives have been addressed.

Conclusions

There has been much debate and controversy about the use of CVS in suspected cases of life threatening abuse over the past few years. Video surveillance of the public, overt or

covert, has been undertaken by police over a number of years in an attempt to prevent crime. The principle of video surveillance, therefore, has been well established. However, the debate around the use and interpretation of CVS continues.²⁵ We acknowledge and share many concerns expressed by various professionals. They conclude on the basis of the report produced by the Northern Region's paediatric subcommittee that CVS is both ethical and legal, and it is acceptable only when used with a rigorous multiagency protocol to safeguard all concerned: the child and his or her family as well as the professionals.²⁶

CVS is ethical if it is necessary to protect the interests of a child and if the child is at serious risk of abuse. It is justified if it is done in the best interest of the child to protect them from serious harm and death. It is the duty of health professionals to use all available approved methods to diagnose and to treat. CVS can therefore be regarded as a justifiable assessment tool to establish a firm diagnosis or to help to exclude deliberate harm to the child. If facilities are not available locally, then the child should be referred to another centre. However, it should never be undertaken without involvement of local child protection procedures, which will include medical, nursing, social work, and police staff.²⁷ Most paediatricians, given the nature and seriousness of the abuse, will agree in principle that CVS is an acceptable diagnostic tool in a small number of children presenting with an apparent life threatening event where child abuse is strongly suspected. Failure to use CVS can lead to a greater risk to life and wellbeing of the child or other children. It should be remembered that in some cases the use of CVS may also prevent the separation of children from innocent parents. A recent editorial²⁸ states that CVS needs to be available as a tool that can be used for some forms of factitious illness. It must also be taken in conjunction with a full child and family assessment, upon which interventions can be based. This is vital for preventing further harm.

Unless we can devise an alternative investigative tool, CVS appears to be necessary, safe, legal, and ethical when operated under a strict and rigorous multiagency protocol in cases of life threatening abuse.

The reports of the Working Group¹⁰⁻¹¹ (Specialist Advisory Committee in Paediatrics) are available from Dr Neela Shabde, Department of Community Child Health, Albion Road Clinic, Albion Road, North Shields NE29 0HG, UK.

NEELA SHABDE

Consultant Paediatrician, Community Child Health Department,
North Tyneside Health Care (NHS) Trust,
Albion Road, North Shields,
Tyne and Wear NE29 0HG, UK

ALAN W CRAFT

Professor and Head, Department of Child Health,
Sir James Spence Institute of Child Health,
Royal Victoria Infirmary,
Newcastle upon Tyne NE1 4LP, UK

- Rosen CL, Frost JD, Bricker T, et al. Two siblings with recurrent cardio-respiratory arrest: Munchausen syndrome by proxy or child abuse? *Pediatrics* 1983;71:715-20.
- Southall DP, Stebbens VA, Rees SV, Lang MH, Warner JO, Shinebourne EA. Apnoeic episodes induced by smothering: two cases identified by covert video surveillance. *BMJ* 1987;294:1637-41.
- Samuels MP, McLaughlin W, Jacobson RR, et al. Fourteen cases of imposed upper airway obstruction. *Arch Dis Child* 1992;67:162-70.
- Epstein MA, Markowitz RL, Gallo DM, Walker Holmes J, Gryboski JD. Munchausen syndrome by proxy: consideration in diagnosis and confirmation by video surveillance. *Pediatrics* 1987;80:220-4.
- Boros SJ, Ophoven JP, Anderson R, Brubaker LC. Munchausen syndrome by proxy: a profile of medical abuse. *Aust Fam Phys* 1995;24:768-73.
- Byard RW, Burnell RH. Covert video surveillance in Munchausen syndrome by proxy: ethics compromise or essential technique? *Med J Aust* 1994;160:352-4.
- Rosen CL, Frost JD, Glaze DG. Child abuse and recurrent infant apnoea. *J Pediatr* 1986;109:1065-7.
- British Paediatric Association Working Party. Report on the evaluation of suspected imposed upper airway obstruction. London: British Paediatric Association, 1994.

- 9 Southall DP, Plunkett MCB, Banks MW, Falkov AF, Samuels MP. Covert video recordings of life threatening child abuse; lessons for child protection. *Pediatrics* 1997;**100**:735–60.
- 10 Working Group of the Specialist Advisory Committee in Paediatrics (Northern). *The role of covert video surveillance in the management of induced illness syndrome*. Newcastle Upon Tyne, 1997.
- 11 Working Group of the Specialist Advisory committee in Paediatrics (Northern). *Covert video surveillance—a framework for developing a protocol*. Newcastle Upon Tyne, 1998.
- 12 Foreman DM, Farsides C. Ethical use of covert videoing techniques in detecting Munchausen syndrome by proxy. *BMJ* 1993;**307**:611–13.
- 13 Thomas T. Covert video surveillance. *New Law Journal* 1994;**114**:966–7.
- 14 Beal J, Blom-Cooper L, Brown B, Marshall P, Mason M. A child in trust. *The report of the panel of inquiry into the circumstances surrounding the death of Jasmine Beckford*. London Borough of Brent: Kingswood Press, 1985.
- 15 Shinebourne E. CVS and the principle of double effect: a response to criticism. *J Med Ethics* 1996;**22**:26–8.
- 16 Children Act 1989. London: HMSO, 1989.
- 17 Department of Health. *The challenge of partnership in child protection: practice guidelines*. London: HMSO, 1995.
- 18 Ethics and Nursing Sub-committee of the Royal College of Nursing. Covert video surveillance: the nursing view. *Bulletin of Medical Ethics* 1995;**106**:3–4.
- 19 North Staffordshire Hospital Trust, Staffordshire Social Services, Staffordshire Police. Guidelines for multi-agency management of patients suspected of or at risk of suffering from life threatening abuse resulting in cyanotic-apnoeic episodes. *J Med Ethics* 1996;**22**:16–21.
- 20 Department of Health. Protocol of videoing children in hospital. *Children Act News* 1994;**12**:5.
- 21 Department of Health. *Working together—under the Children Act 1989*. London: HMSO, 1995. [Addendum: Child Protection—medical responsibilities.]
- 22 Re: Regina v Khan (Sultan). Criminal evidence—admissibility—unlawful listening device. London and UK. *The Times Law Reports* 1994:313–14.
- 23 Re: DH, A minor (Child Abuse). London, UK: *Family Law Reports* 1994:679–716.
- 24 Home Office. Guidance on the use of equipment in police surveillance operations. London: HMSO, 1984.
- 25 Morley C. Concerns about using and interpreting covert video surveillance. *BMJ* 1998;**316**:1603–4.
- 26 Shabde N, Craft AW. Covert video surveillance is acceptable—but only with a rigorous protocol [commentary]. *BMJ* 1998;**316**:1604–5.
- 27 Williams C, Bevan VT. The secret observation of children in hospital. *Lancet* 1988;**i**:780–1.
- 28 Jones DPH, Lynch MA. Diagnosing and responding to serious child abuse [editorial]. *BMJ* 1998;**317**:484–5.