Disabilities and trends over time in a French county, 1980–91

C Cans, P Guillem, J Fauconnier, P Rambaud, P S Jouk

See end of article for authors' affiliations

Arch Dis Child 2003;88:114–117

Correspondence to:
Dr C Cans, Registre des
Handicaps de l'Enfant et
Observatoire Périnatal de
l'Isère, 23, Av Albert 1er
de Belgique, 38000
Grenoble, France;
christine.cans@imag.fr

Accepted 19 September
2002

Aim: To describe trends over time and types of disability for children born in a French county from 1980 to 1991.

Methods: Data were collected from medical records of a morbidity register; disabled children with at least one severe deficiency have been included. Prevalence rates are given per 1000 resident children, over four three-year periods.

Results: Overall, 7.73 per 1000 children (that is, 1360 children), had a severe childhood disability, and the prevalence rate had increased significantly since 1980. This increase was mainly owing to an increase in cerebral palsy and psychiatric disorder prevalence rates.

Conclusion: Future research aimed to explain these trends over time can be based on such data. The present knowledge is useful for planning purposes.

Considerable changes in obstetric and neonatal care over the past decades have resulted in a decrease in infant mortality. However, several authors have noted that this decrease in infant mortality has not been followed by a decrease in infant morbidity. In some places, the cerebral palsy prevalence rate has remained stable,¹ while in others it has increased^{2–4}; a decrease in some subgroups has recently been reported.⁵

The educational and medical care to be given to disabled children has been laid down by law in France since 1975.⁶ Since then, very few surveys have described the trends in childhood disability.^{7,8} However, knowledge about types of childhood disabilities and trends over time might prove to be very useful for those specialised schools and care institutions that deliver care to these children, and for those making health policy decisions. Thus, in this paper, we describe types of childhood disabilities and their prevalence trends over time for children born in one French county between 1980 and 1991.

POPULATION AND METHODS

Population

The morbidity register, "Registre des Handicaps de l'Enfant et Observatoire Périnatal (RHEOP)", was established in 1992 in France, with the aim of monitoring trends over time for severe childhood disabilities. Isère county has 1 000 000 inhabitants, and 14 000 live births per year. All disabled children living in Isère county, aged 7 years, with a clinical feature of a severe deficiency are registered.

Only children with a severe neurosensory deficiency were recorded. The criteria used to determine the severity of the deficiency were those usually used (table 1). Children with two or three mild or moderate deficiencies were not included, as it was considered that it would be too difficult to guarantee reproducibility over time, if such cases were included.

Registration of cases was done "actively" by a medical doctor going to the different sources. The main source was the County Commission for Specialised Care and Education of Disabled children (CDES), which represents the only way of obtaining financial support or of being admitted into a specialised school. Other sources were departments of paediatrics, psychiatric day care hospitals, and specialised institutions for disabled children. Data were collected from the

medical records at the CDES or at hospitals when the child was 7 years old. The registration form included: (a) administrative information; (b) medical information, using ICD9 codes for diagnosis of the main deficiency, aetiological factors, and associated disabilities, including congenital anomalies; and (c) when available, birth weight, gestational age, and neonatal care admission data. Parental authorisation was not obtained for 2% of the children (range 1–6%). The completeness of the data collection process was considered to be satisfactory.⁹

Methods

Prevalence rates for these childhood disabilities were calculated per 1000 children of the same age living in Isère county. With respect to population migration in and out of the county, it was decided to use results for prevalence rates per 1000 resident children rather than those per 1000 live births, as a previous survey showed that the magnitude of migration into this county was the same for healthy children and disabled children.¹⁰ Analysis of prevalence rates over time was carried out for three year periods, and the Armitage trend test¹¹ was used when checking for trends over time.

RESULTS

Types of disability

A total of 1360 children born between 1980 and 1991 (797 boys and 563 girls), satisfied the inclusion criteria for "at least one severe deficiency"—that is, around 120 children for each birth year cohort, with a significant overall male/female sex ratio of 1.42 ($p < 0.001$). Table 1 shows the different types of severe disabilities present in these 1360 children. As a child might present more than one disability, the total number of disabilities is greater than the number of disabled children. The mean number of ascertainment sources was 1.7 per child.

Trends over time

Table 2 shows the prevalence rates for the overall period, and for consecutive three year periods. The overall prevalence rate of 7.73 per 1000 has increased significantly since 1980 ($p < 0.001$). The prevalence rate of motor disability has increased slightly ($p = 0.17$); this increase mainly concerned the cerebral palsy prevalence rate ($p = 0.01$), while the prevalence rate of other motor disabilities has remained stable ($p = 0.31$). The prevalence rate for intellectual disability

Table 1 Severe disabilities at 7 years of age in Isere county 1980–91 (1360 children, 1643 disabilities)

	Birth years 1980–91 (n)	% within each type of disability	Proportion among all severely impaired children
<i>Motor disability (1)</i>	558		38.7
“IMC” among cerebral palsy	272	48.7	
Other cerebral palsies	88	15.8	
Nervous system congenital anomalies	37	6.6	
Other congenital anomalies	66	11.8	
Degenerative disease (hereditary or not)	52	9.3	
Other motor disability	43	7.8	
<i>Psychiatric disorders (2)</i>	353		24.5
Kanner autism	31	8.8	
Other autisms	58	16.4	
Psychosis with severe mental retardation	58	16.4	
Other psychoses	206	58.4	
<i>Intellectual disability (3)</i>	484		33.6
Down’s syndrome	167	34.5	
Other chromosomal anomalies	20	4.1	
Other congenital anomalies	63	13.0	
Other intellectual disability	234	48.4	
<i>Hearing disability (4)</i>	146		10.1
Without any other disability	85	58.2	
With at least one associated disability	61	41.8	
<i>Visual disability (5)</i>	102		7.1
Without any other disability	40	39.2	
With at least one associated disability	62	60.8	

Severity criteria

(1) None for Infirmitté Motrice Cérébrale (IMC) which is part of the whole group entitled Cerebral Palsy (CP), defined as a clinical association of posture and movement disorders resulting from a non-progressive lesion on an immature brain.¹²

Need for specific help and/or continuous physiotherapy care was the severity criterion for the other motor deficiencies.

(2) ICD codes F84.0 to F84.5.

(3) IQ <50 or mental retardation classified as profound, severe, or moderate.

(4) Loss of >70 db in each ear before correction.

(5) Visual acuity <3/10 in each eye after correction.

Table 2 Prevalence rates of severe disabilities in children born 1980–91

	Birth years				Total	95% CI
	80–82	83–85	86–88	89–91		
Children with at least one severe disability	314	308	351	387	1360	
Children living in Isere county at 7 years of age	45829	42950	43408	43732	175919	
Prevalence rate per 1000	6.85	7.17	8.09	8.85	7.73	7.33 to 8.15
<i>Motor disability</i>	134	132	144	148	558	
Prevalence rate per 1000	2.92	3.07	3.32	3.38	3.17	2.91 to 3.43
Of whom cerebral palsy	79	81	92	107	359	
Prevalence rate per 1000	1.72	1.89	2.12	2.45	2.04	1.83 to 2.25
<i>Psychiatric disorders</i>	65	65	105	118	353	
Prevalence rate per 1000	1.42	1.51	2.42	2.70	2.01	1.80 to 2.22
Of whom autism	14	12	35	28	89	
Prevalence rate per 1000	0.31	0.28	0.81	0.64	0.51	0.40 to 0.62
<i>Severe mental retardation</i>	128	122	104	130	484	
Prevalence rate per 1000	2.79	2.84	2.40	2.97	2.75	2.51 to 2.99
<i>Hearing or visual disability</i>	63	56	60	62	241	
Prevalence rate per 1000	1.37	1.30	1.38	1.42	1.37	1.20 to 1.54

(severe mental retardation) has remained stable over time ($p = 0.94$), even when looking at subgroups—that is, Down’s syndrome ($p = 0.68$), or other intellectual disabilities ($p = 0.70$). Overall, the prevalence rate has increased for psychiatric disorders ($p < 0.001$). However, for psychosis with severe mental retardation, the prevalence rate has remained stable ($p = 0.41$), while for autism and other psychoses, the rates have actually increased ($p = 0.001$ and $p < 0.001$, respectively). The prevalence rate for sensory disability has not changed since 1980 ($p = 0.80$).

Characteristics

The socioeconomic status, determined by the occupation of the father and/or mother of the disabled child, was compared with the distribution figures for parental occupation within the Isere population in 1990. Mothers of the disabled children were often without occupation, more often than other women of the same age in the whole population (55% versus 31%, $p < 0.001$). Parental occupation statistics were different, with more fathers of low economic status (working in factories) and fewer fathers of high economic status (businessmen,

professors, etc) among families with disabled children (46% versus 38%, and 8% versus 13% respectively, $p < 0.001$).

One hundred and nine disabled children (8.0%, range 6.6–9.4%) had a clear recognised postneonatal aetiology. Among the other severely disabled children with available information on pre/perinatal characteristics, 34.5% had a congenital anomaly, 6% had a very low birth weight (<1500 g), and 42% had been hospitalised in a neonatal care unit. Although there was a slight increase for the proportion of severely disabled children with a very low birth weight (from 3.9% to 8.1%, $p = 0.12$), the proportions with congenital anomaly or hospitalised in a neonatal care unit were remarkably stable during the studied period ($p = 0.87$ and $p = 0.93$ respectively).

One third of the severely disabled children (33.1%, range 30.1–36.1%), were born with a normal birth weight, without congenital anomaly, and had not been hospitalised during their first month of life. This proportion varied according to the type of deficiency: 25.1% for motor deficiencies, 22.0% for intellectual disability, 53.1% for psychiatric disorders, and 42.9% for sensory deficiencies.

DISCUSSION

According to our inclusion criteria, 7.73 per 1000 children born 1980–91 were registered with a severe disability at 7 years of age.

Data collection was conducted slightly differently before and after the 1984 birth cohort, with a retrospective data collection before 1984. This might account for the observed increase after 1984, as it is usually more difficult to find cases retrospectively than prospectively. However, we do not believe that this is the case, for two reasons:

- With the capture recapture method⁹ we were able to obtain a regular follow up of the degree of ascertainment, which remained stable over time, from 96% in the first period to 93% in the last period.
- We registered only severely disabled children, and it seems unlikely that children with such severe disability would have escaped retrospective data collection.

The increase in the overall prevalence rate for severely disabled children between 1980 and 1990 is mainly a result of the increase in the rates of cerebral palsy and severe psychiatric disorders. The cerebral palsy rate observed in our study is 2.04 per thousand, similar to the rates and trends observed elsewhere in Europe.^{5–13} The rate of autism observed over the whole period, 0.51 per 1000, is close to the rates published in other studies,^{14–16} and doubled during the study period from 0.31 to 0.64. The recent increase in the autism prevalence rate observed in different places has led to much discussion, without any clear and/or complete explanations being put forward. Gillberg *et al*, in Sweden, have suggested that the most probable reasons for this increase are classification bias and migratory movement into the area of study.¹⁷ Certainly it is necessary to obtain a more consistent definition of autism, and to have routine monitoring of autism trends, in order to avoid the problems of differences in definitions.¹⁸

The rate of severe mental retardation remained stable during the 1980–90 period, 2.75 per 1000, while elsewhere in France a decreasing trend was observed, Down's syndrome excluded, between 1976 and 1985.¹⁹ In England, the prevalence rate of severe mental retardation was higher in the 1970s, 3.0–4.1 per 1000 for children aged 5–9 years.²⁰ Recent studies in other countries have shown a rate of around 3.0 per 1000 children.^{21–22}

The rate of severe visual deficiency was 0.57 per 1000, which is lower than the rate of 0.70 per 1000 observed in the Northern Europe register,²³ but very close to the rate observed in Italy, 0.50 per 1000.²⁴ This cannot be explained by differences in definitions, as the threshold for defining severity was the same in these studies. In France, Arnaud *et al* observed great

variations between counties, according to the number of specialised care places in each area.²⁵

Social class differences in the prevalence rates for neurodevelopmental disabilities have been described,²⁶ and it has been suggested that the excess of disability in lower social classes could be the result of pregnancy complications, which are more frequent in deprived mothers.²⁷

In future studies about disabled children, the existence of a large group of children without any adverse pre/perinatal characteristics must be kept in mind. Knowledge of the high proportion of severely disabled children with congenital anomaly is useful for planning purposes, and also for early screening and care strategies in order to provide optimal support as soon as possible.

Authors' affiliations

C Cans, P Guillem, Registre des Handicaps de l'Enfant et Observatoire Périnatal de l'Isère, 23, Av Albert 1er de Belgique, 38000 Grenoble, France

J Fauconnier, Centre Hospitalier Universitaire de Grenoble, Département of Information and Biostatistics, BP 217, 38043 Grenoble cedex 09, France

P Rambaud, P S Jouk, Centre Hospitalier Universitaire de Grenoble, Département of Neonatology, BP 217, 38043 Grenoble cedex 09, France

We acknowledge the Conseil Général de l'Isère

REFERENCES

- 1 **Robertson C**, Hrynchysyn J, Etches P, *et al*. Population-based study of the incidence, complexity, and severity of neurologic disability among survivors weighing 500 through 1250 grams at birth: a comparison of two birth cohorts. *Pediatrics* 1992;**90**:750–5.
- 2 **Hagberg B**, Hagberg G. The changing panorama of infantile hydrocephalus and cerebral palsy over forty years—a Swedish survey. *Brain Dev* 1989;**11**:368–73.
- 3 **Pharoah PoD**, Cooke R, Rosenbloom L. Birthweight specific trends in cerebral palsy. *Arch Dis Child* 1990;**62**:379–84.
- 4 **Stanley F**, Blair E. Why have we failed to reduce the frequency of cerebral palsy? *Med J Aust* 1991;**154**:623–6.
- 5 **Hagberg B**, Hagberg G, Beckung E, *et al*. Changing panorama of cerebral palsy in Sweden. VIII. Prevalence and origin in the birth year period 1991–94. *Acta Paediatr* 2001;**90**:271–7.
- 6 **Loi d'orientation en faveur des personnes handicapées no. 75.534** du 30 Juin 1975 & Annexes XXIV, XXIVbis, XXIVter. In: Décret no. 89.878.
- 7 **Manciaux M**, Salbreux R, Tomkiewicz S. Mental retardation and multiple handicaps. *Arch Fr Pédiatr* 1981;**38**:627–34.
- 8 **Rumeau-Rouquette C**, du Mazaurun C, Verrier A, *et al*. *Prévalence des handicaps, Evolution dans trois générations d'enfants 1972, 1976, 1981*. Paris: INSERM, 1994.
- 9 **Guillem P**, Lavergne C, Cans C. How to calculate the completeness of ascertainment of a morbidity register? *Rev Epidemiol Sante Publique* 2000;**48**:41–51.
- 10 **Baillie MF**, Arnaud C, Grandjean H, *et al*. Prevalence calculation of childhood deficiencies: methodological points. *Archives of Public Health* 1996;**53**:13.
- 11 **Armitage P**, Berry G. *Statistical methods in medical research*, 2nd edn. Oxford: Blackwell Scientific Publications, 1987.
- 12 **Amiel-Tison C**. *L'infirmité motrice d'origine cérébrale*. Paris: Masson, 1997.
- 13 **SCPE Collaborative Group**. Surveillance of cerebral palsy in Europe: why a European collaboration of cerebral palsy surveys and registers? *Dev Med Child Neurol* 2000;**42**:816–24.
- 14 **Fombonne E**, du Mazaurun C, Cans C, *et al*. Autism and associated medical disorders in an epidemiological survey of French children. *J Am Acad Child Adolesc Psychiatry* 1997;**36**:1561–9.
- 15 **Cialdella P**, Mamelie N. An epidemiological study of infantile autism in a French department. *J Child Psychol Psychiatry* 1989;**30**:165–75.
- 16 **Burd L**, Fisher W, Kerbeshian J. A prevalence study of pervasive developmental disorders in North Dakota. *J Am Acad Child Adolesc Psychiatry* 1987;**26**:700–3.
- 17 **Gillberg C**, Steffenburg S, Schauman H. Is autism more common now than ten years ago? *Br J Psychiatry* 1991;**158**:403–9.
- 18 **Fombonne E**. Is there an epidemic of autism? *Pediatrics* 2001;**107**:411–13.
- 19 **Rumeau-Rouquette C**, Grandjean H, Cans C, *et al*. Prevalence and time trends of disabilities in school-age children. *Int J Epidemiol* 1997;**26**:137–45.
- 20 **Pharoah PoD**. The epidemiology of chronic disability in childhood. *Int Rehabil Med* 1985;**7**:11–17.
- 21 **Schaefer G**, Bodensteiner JB. Evaluation of the child with idiopathic mental retardation. *Pediatr Clin North Am* 1992;**39**:929–43.

- 22 **Murphy CC**, Yeargin-Allsop M, Decoufflé P, *et al*. The administrative prevalence of mental retardation in 10 year-old children in Metropolitan Atlanta 1985 through 1987. *Am J Public Health* 1995;**85**:319–23.
- 23 **Riise R**, Flage T, Hansen E, *et al*. Visual impairment in Nordic children. I. Nordic registers and prevalence data. *Acta Ophthalmol Scand* 1992;**70**:145–54.
- 24 **Nicolosi A**, Marighi P, Rizzardi P, *et al*. Prevalence and causes of visual impairment in Italy. *Int J Epidemiol* 1994;**23**:359–64.
- 25 **Arnaud C**, Baille MF, Grandjean H, *et al*. Visual impairment in children: prevalence, aetiology and care, 1976–85. *Paediatr Perinat Epidemiol* 1998;**12**:228–39.
- 26 **Dolk H**, Parkes K, Hill AE. Cerebral palsy prevalence in relation to socio-economic deprivation in Northern Ireland. *Paediatr Perinat Epidemiol* 1996;**10**:A4.
- 27 **Taylor DJ**, Davidson J, Howie PW, *et al*. Do pregnancy complications contribute to neurodevelopmental disability. *Lancet* 1985;**30**:713–16.

IMAGES IN PAEDIATRICS.....

Dumbbell tumour of the mediastinum

A 6 month old male infant presented with fever and cough of one month duration. The chest radiograph showed bilateral radio-opacities occupying the medial aspect of each hemithorax (fig 1). Computerised tomography (CT) revealed a dumbbell shaped tumour occupying both hemithoraces (fig 2). The tumour was continuous across the midline by an isthmus passing posterior to the descending aorta and displacing the oesophagus anteriorly. The tumour had fat and soft tissue attenuation and did not enhance significantly. It was completely excised by bilateral posterolateral thoracotomy. Postoperative recovery was uneventful. Lipoblastoma was diagnosed on histopathology.

Lipoblastoma is a rare tumour of embryonal fat that occurs almost exclusively in children under the age of 3 years.¹ The tumour presents as a localised, well circumscribed lesion known as lipoblastoma or a diffuse/multicentric form known as lipoblastomatosis.² Most lesions present as a subcutaneous soft tissue mass on the extremities or head and neck. Truncal lesions are less common.³ Focal tumours behave in a benign manner and diffuse lesions may recur if incompletely excised.¹ To our knowledge, only five patients with mediastinal lipoblastoma have been reported in the literature.² This is the only instance of a dumbbell shaped mediastinal tumour occupying both hemithoraces.

K L N Rao, B Eradi, R K Marwaha, A Trehan, A K Saxena, P Menon

Department of Paediatric Surgery, Paediatric Medicine and Radiodiagnosis, Postgraduate Institute of Medical Education and Research, Chandigarh, 160012, India; klnrao@hotmail.com

References

- 1 **Dilley AV**, Patel DL, Hicks MJ, *et al*. Lipoblastoma: pathophysiology and surgical management. *J Pediatr Surg* 2001;**36**:229–31.
- 2 **Frederici S**, Cuoghi D, Sciutti R. Benign mediastinal lipoblastoma in a 14-month-old infant. *Pediatr Radiol* 1992;**22**:150–1.
- 3 **Irgau I**, McNicholas KW. Mediastinal lipoblastoma involving the left innominate vein and the left phrenic nerve. *J Pediatr Surg* 1998;**33**:1540–2.

Figure 1 Anteroposterior chest radiograph showing bilateral hilar radio-opaque shadows.

Figure 2 CT scan showing a dumbbell shaped mediastinal tumour with fat and soft tissue attenuation. The tumour is compressing both lungs and displacing mediastinal structures anteriorly.