as they cope with the pain at home. This would include appropriate use of oral analgesia, home visits by sickle nurse specialists, alternative therapies, and help in the development of psychological coping mechanisms.

Authors' affiliations

S Chakravorty, K Newell, J Ramchandani, K Qureshi, R Ibrahim, B Datta, P T Telfer, Royal London Hospital, UK

Correspondence to: Dr P T Telfer, Department of Haematology, Royal London Hospital, Whitechapel Road, London E1 1BB, UK; ptelfer@cytanet.com.cy

Accepted 6 July 2003

REFERENCES

- 1 **Platt OS**, Thorington BD, Brambilla DJ, *et al*. Pain in sickle cell disease. Rates and risk factors. *N Engl J Med* 1991;**325**:11–16.
- 2 **Brozovic M**, Davies S. Management of sickle cell disease. *Postgrad Med J* 1987;**63**:605–9.
- 3 **Ware MA**, Hambleton I, Ochaya I, *et al*. Day-care management of sickle cell painful crisis in Jamaica: a model applicable elsewhere? *Br J Haematol* 1999;**104**:93–6.
- 4 **Thomas VJ**, Hambleton I, Serjeant GR. Psychological distress and coping in sickle cell disease: comparison of British and Jamaican attitudes. *Ethnicity and Health* 2001;**6**:129–36.
- 5 **Fuggle P**, Shand PA, Gill LJ, *et al*. Pain, quality of life, and coping in sickle cell disease. *Arch Dis Child* 1996;**75**:199–203.
- 6 **Shapiro BS**, Dinges DF, Orne EC, *et al*. Home management of sickle cell-related pain in children and adolescents: natural history and impact on school attendance. *Pain* 1995;**61**:139–44.

IMAGES IN PAEDIATRICS.....

Wheezing with a whistle

A 2 year old boy presented with history of sudden onset respiratory distress of three days duration. On examination he had right sided wheeze. Chest x ray examination revealed a radio-opaque foreign body in the right main bronchus. An emergency bronchoscopy was done and the foreign body (a whistle used in toys) was removed, after which his wheeze subsided.

A B Sivam, M Nath

Kanchi Kamakoti Child's Trust Hospital, Chennai, India;
adhisivam1975@yahoo.co.uk

doi: 10.1136/ad.2003.037358