ORIGINAL ARTICLE

Should parents accompany critically ill children during inter-hospital transport?

J Davies, S M Tibby, I A Murdoch

Arch Dis Child 2005;90:1270-1273. doi: 10.1136/adc.2005.074195

See end of article for
authors' affiliations

Correspondence to:
Dr S Tibby, Paediatric
Intensive Care Unit, Guy's
and Saint Thomas' NHS
Foundation, Guy's
Hospital, St Thomas Street,
London SE1 9RT, UK;
shane.tibby@gstt.sthames.
nhs.uk

Accepted 6 May 2005
Published Online First
12 May 2005

Background: Parental accompaniment during inter-hospital transportation (retrieval) of critically ill children is not commonplace in the United Kingdom.

Methods: A three month pilot of parental accompaniment was undertaken in 2002 (143 retrievals), after which time the policy was adopted as standard practice. A follow up audit was performed in 2004 (136 retrievals).

Results: Findings were remarkably consistent between the two periods. Staff perceived little or no added stress during the majority of transfers (96% in 2002, 98% in 2004), and felt able to perform medical interventions without hindrance (98% in 2002, 100% in 2004). There was good agreement between medical and nursing staff regarding perception of stress and ability to perform interventions (phi statistic 0.57 to 1.00). Adverse events occurred during 11 (3.9%) retrievals; six of these involved a parent exclusively. Stress tended to be associated with adverse events or parental behaviour rather than disease acuity. Staff vetoed the offer of accompaniment on 11 occasions, for a variety of reasons. The majority of parents found the experience safe, beneficial, and perceived a reduction in stress as a result. These data may inform other retrieval services who are considering adopting a similar policy.

The ethos that parents should have unimpeded access to their child when hospitalised is accepted widely, and is endorsed in the Children's National Service Framework.¹ It is debatable however, whether parents should be present throughout every phase of their child's treatment, particularly during episodes that may involve risk.²⁻⁵ Two situations often cited are the induction of anaesthesia,^{6,7} or when cardiopulmonary resuscitation is required.⁸ A third example, namely inter-hospital transportation of the critically ill child, has become increasingly relevant in the United Kingdom over the last decade. This has come about as a result of centralisation of paediatric intensive care services; with over 4000 episodes estimated to occur annually, the majority via specialised retrieval teams.⁹

It is unusual, however, for a parent to be offered the opportunity to accompany their child during transfer, a view acknowledged by two national documents detailing standards for inter-hospital transport.^{10,11} Reluctance to allow parental accompaniment may be for a variety of reasons, including lack of space in the ambulance, the potential for inducing stress in both staff and parents (particularly in an emergency), and even diversion of staff's attention from the patient.^{12,13}

The South Thames Retrieval Service is one of the largest paediatric retrieval services in the UK, undertaking approximately 600 retrievals per year.¹⁴ A recent postal survey of 233 parents revealed a high degree of satisfaction with the service; however, a recurrent theme that emerged was the level of parental distress as a consequence of separation from their child during the transfer period.¹⁵ In response, a three month pilot was undertaken in October 2002, whereby a parent or carer was offered the opportunity to travel in the ambulance with their child. Staff and parent experiences were audited over this period, and on the basis of positive findings from the audit, parental accompaniment was adopted as standard practice. A follow-on audit was carried out 18 months later. This report details the findings of the two audits, examining the impact that parental accompaniment has had on the delivery of the service.

METHODS

A preliminary, anonymised survey of 100 PICU staff was undertaken in April 2002, to gauge attitudes regarding parental accompaniment. Although 72% of respondents supported this concept, one third of this group specified that accompaniment should not be offered unconditionally. Several themes were identified, including: severity of the child's illness, mental state of the parents, a clear explanation of the benefits and risks being given to parents before embarkation, and the ability of the ambulance crew to look after parents if cardiopulmonary resuscitation was needed. Many staff expressed concerns that they would be placed under undue stress, and may be inhibited from carrying out medical procedures.

On the basis of the survey, a three month pilot was undertaken between October and December 2002 (period 1), whereby one parent or guardian was offered the opportunity to travel in the ambulance. Any of the retrieval team members (doctor, nurse, ambulance driver) had right of veto. Parents were given an information sheet about what to expect when travelling in the retrieval ambulance, and retrieval staff discussed aspects of safety with parents prior to departure. Before commencing the pilot, clarification was sought about cover provided by the ambulance service's insurance policy with respect to parental accompaniment.

Data were collected prospectively concerning patient demographics, retrieval team composition, medical and nursing interventions required during the journey (interventions were defined as those over and above routine physiological monitoring), and the occurrence of adverse events. The definition of an adverse event has been published previously.¹⁶ Briefly, an adverse event was defined as any unforeseen occurrence which could actually or potentially affect patient care or the safety of anyone in the ambulance. These included: (a) physiological deteriorations (for example, hypotension, hypoxia); (b) equipment/therapy failures (for example, loss of oxygen supply, mechanical ventilator malfunction); or (c) other (motor vehicle accident, aggressive/abusive behaviour). Nursing and medical staff were

Table 1 Patient demographics

	Period 1	Period 2	p value
Total number	143	136	
Median patient age (months)	7.6 (1.4–43)	18.3 (1.7–60)	0.14
Median mortality risk (%) [*]	10 (5–14)	5.3 (2.5–11)	0.02
Number (%) mechanically ventilated	109 (76%)	80 (59%)	0.003
Number (%) requiring inotropic support	19 (13%)	19 (14%)	0.87

^{*}Calculated using the Paediatric Index of Mortality score.
Numbers in parentheses represent the interquartile range.

asked independently, via a questionnaire, to rate their stress levels and perceived ability to intervene during the transport. Rating was via a five point Likert scale, ranging from “stress free” to “very stressful”, and from “intervene easily” to “felt inhibited”. Staff were also asked to state whether or not they thought it was beneficial to have a parent accompanying their child.

Parents were given a questionnaire after arriving in the PICU, asking if they had been given the opportunity to travel in the ambulance, and asking about perceived levels of safety and information given during the transfer.

Following the pilot, parental accompaniment was adopted as standard practice, and a follow up audit was undertaken 18 months later, between April and June 2004 (period 2).

Ethical approval was not sought for this audit; however, the audit was registered with our hospital Trust via the standard procedure.

Statistics

Data are presented as raw counts and percentages, while summary statistics are reported as median (interquartile range). Demographic data are compared using the Mann-Whitney test (continuous) and χ^2 test (categorical). For the staff questionnaire, inter-observer agreement was assessed using the phi statistic (Φ).¹⁷ Phi is more robust than kappa in situations where the majority of outcomes cluster at one extreme (for example, if the majority of retrievals were perceived as stressful). When this occurs, kappa derived agreement over and above chance is small, producing very low kappa values. Phi corrects for this phenomenon, producing a chance independent measure of inter-observer agreement.¹⁸ Like kappa, values for Φ range from -1.0 to $+1.0$, with level of agreement interpreted as follows: less than 0, poor agreement; 0 to 0.2, slight; 0.2 to 0.4 fair; 0.4 to 0.6 moderate; 0.6 to 0.8 substantial; and 0.8 to 1.0, excellent agreement.

RESULTS

Staff questionnaires were returned for 143/155 (92%) retrievals during period 1 and 136/150 (91%) during period

2, and form the basis of this report. The median (inter-quartile) journey time from referring hospital to PICU was 40 minutes (25–65). Case mix differed slightly between the epochs; patients in period 2 were slightly older with a lower incidence of mechanical ventilation, and hence disease severity (table 1).

In both periods, the retrieval team typically comprised a doctor, a nurse, and a dedicated ambulance crew. On 32% of occasions, the team included more than one member of the medical staff (typically a fellow and resident), and in 15% there was more than one nurse in attendance (usually for training purposes). The most senior medical staff member present was a consultant (2%), fellow (87%), and resident (11%). The grade of the most senior nurse in attendance was H (1%), G (14%), F (45%), and E (40%). Medical or nursing intervention was required in 39% (109/279) of retrievals (many of these required more than one type of intervention), with the commonest interventions being: administration of sedation and/or neuromuscular blockade (26%), fluid bolus (10%), commencement or adjustment of an inotrope infusion (7%), and endotracheal suction (5%). Adverse events occurred during 11 retrievals (3.9%), and are shown in table 2. Interestingly, the majority of these were not patient related, but rather involved staff or parents, or pertained to a logistic aspect of the retrieval.

Parental accompaniment was offered for the majority (250/279) of transfers, and was accepted in 71% (178/250) of these. The commonest reason for declining accompaniment was the desire for both parents to travel together. The team was unable to offer accompaniment due to a parent not being present at the referring hospital on 18/279 episodes (6.5%), and staff veto occurred on 11/279 occasions (3.9%). Reasons for retrieval team veto were varied, and are shown in table 3. There was no obvious relation between the occurrence of team veto and the seniority of the team members.

The majority of staff found parental accompaniment to be non or minimally stressful (96% period 1, 98% period 2), and found little or no difficulty in performing medical/nursing interventions (98% period 1, 100% period 2). There was generally high agreement between nursing and medical staff

Table 2 Adverse events occurring during retrieval

Period	Age (mth)	Mortality risk	Diagnosis	Type
1	180	54%	Hanging	Cardiopulmonary resuscitation
1	1.2	52%	Bronchiolitis/HLH	Mechanical ventilator failure
1	156	14%	Seizures	Unable to fit seatbelt on mother
1	11	0.9%	Upper airway obstruction	Delay leaving while waiting for mother's bag
1	7	51%	HLH	Mother travelsick
1	0.1	37%	TGA	Nurse travelsick
2	172	4.7%	Status epilepticus	Low battery on infusion pumps
2	2	9.8%	Anaphylaxis	Mother phobic of blue lights (didn't inform staff)
2	166	8.9%	Head injury	Monitor failure
2	6	4.4%	Pneumonia	Mother travelsick
2	1.2	6.3%	Pneumonia	Mother verbally aggressive as father not able to also accompany

HLH, hypoplastic left heart syndrome; TGA, transposition of the great arteries.

What is already known on this topic

- Parental accompaniment during inter-hospital road transportation of critically ill children is uncommon in the UK
- Experience from the United States suggests that this may place extra stress on staff

Although rare, parent related adverse events during the retrieval did have an impact on our staff, producing a degree of stress that was higher among nurses. Adverse events causing greatest stress were those that involved a parent becoming unwell, or exhibiting aggressive behaviour. Although staff did not perceive difficulty in performing interventions on these occasions, we are unable to assess whether stress compromised performance in other ways (such as decision making and technical competence); thus this factor must be born in mind by other retrieval services that are considering parental accompaniment. Conversely, many staff members reported that having a parent present produced a calming effect on non-ventilated patients, similar to the United States experience.¹³ Other benefits reported by staff included opportunities to develop a rapport with the parent, explain aspects of the child's illness, clarify what to expect while in the intensive care unit, and to take a full medical history.

These themes were mirrored by the parents' survey, where the perception of benefit was an unambiguous and recurrent qualitative finding. It was remarkable how similar many of the comments were to those reported from an audit in the United States.²⁰ The latter report highlighted that parental requests for accompaniment were more likely for children who were older, non-ventilated, and for those who have been previously well (as opposed to suffering from chronic illness); conversely disease acuity did not appear to influence the desire to travel in the ambulance.²⁰ We also found that parents who did not accompany often reported additional stress associated with negotiating transport to the referral centre.

A service development may initially be followed by a honeymoon phase, after which time enthusiasm for the development wanes, or deficiencies become apparent. Thus we were reassured to see consistency in results, both for staff and parents, across the two audit periods spanning 20 months. As a result, we have continued to provide the service in its current format, including preservation of the right of staff veto. It is hoped that our results may inform other services that are considering adopting a similar policy.

Authors' affiliations

J Davies, S M Tibby, I A Murdoch, Department of Paediatric Intensive Care, Guy's and Saint Thomas' NHS Foundation Trust, London, UK

Competing interests: none

What this study adds

- Parental accompaniment per se does not appear to increase staff stress nor compromise the performance of medical interventions in the majority of occasions. When present, staff stress tends to be associated with parental behaviour and the occurrence of adverse events rather than the child's disease acuity
- Although adverse events were rare, a significant proportion involved a parent directly

REFERENCES

- 1 **Department of Health.** *Getting the right start: National Service Framework for Children, Young People and Maternity Services. Part 1: Standard for hospital services.* Department of Health, 2003.
- 2 **Bauchner H,** Waring C, Vinci R. Parental presence during procedures in an emergency room: results from 50 observations. *Pediatrics* 1991;**87**:544–8.
- 3 **Merritt KA,** Sargent JR, Osborn LM. Attitudes regarding parental presence during medical procedures. *Am J Dis Child* 1990;**144**:270–1.
- 4 **Fein JA,** Ganesh J, Alpern ER. Medical staff attitudes toward family presence during pediatric procedures. *Pediatr Emerg Care* 2004;**20**:224–7.
- 5 Beckman AW, Sloan BK, Moore GP, et al. P50 (Parental Presence during Painful Pediatric Procedures) Research Group. Should parents be present during emergency department procedures on children, and who should make that decision? A survey of emergency physician and nurse attitudes. *Acad Emerg Med* 2002;**9**:154–8.
- 6 **Kain ZN,** Caldwell-Andrews AA, Krivtza DM, et al. Trends in the practice of parental presence during induction of anesthesia and the use of preoperative sedative premedication in the United States, 1995–2002: results of a follow-up national survey. *Anesth Analg* 2004;**98**:1252–9.
- 7 **Kain ZN,** Mayes LC, Wang SM, et al. Parental presence and a sedative premedication for children undergoing surgery: a hierarchical study. *Anesthesiology* 2000;**92**:939–46.
- 8 **Boudreaux ED,** Francis JL, Loyacono T. Family presence during invasive procedures and resuscitations in the emergency department: a critical review and suggestions for future research. *Ann Emerg Med* 2002;**40**:193–205.
- 9 **Universities of Leeds, Leicester and Sheffield.** Paediatric Intensive Care Audit Network. Annual report Mar 2003–Feb, 2004:p49.
- 10 **Department of Health.** *Paediatric intensive care. "A framework for the future". Report from the National Coordinating Group on Paediatric Intensive Care to the Chief Executive of the NHS executive.* Department of Health, 1997.
- 11 **Paediatric Intensive Care Society.** *Standards for retrieval of critically ill children.* Paediatric Intensive Care Society, UK, 2001.
- 12 **Stack CG,** Bennett NR, Goddard JM. Should paediatric intensive care be centralised. *Lancet* 1997;**350**:65–6.
- 13 **Woodward GA,** Fleegler EW. Should parents accompany pediatric interfacility ground ambulance transports? Results of a national survey of pediatric transport team managers. *Pediatr Emerg Care* 2001;**17**:22–7.
- 14 **Doyle YG,** Orr FE. Interhospital transport to paediatric intensive care by specialised staff: experience of the South Thames combined transport service, 1998–2000. *Arch Dis Child* 2002;**87**:245–7.
- 15 **Colville G,** Orr F, Gracey D. 'The worst journey of our lives': parents' experiences of a specialised paediatric retrieval service. *Intensive Crit Care Nurs* 2003;**19**:103–8.
- 16 **Edge WE,** Kanter RK, Weigle CG, et al. Reduction of morbidity in interhospital transport by specialized pediatric staff. *Crit Care Med* 1994;**22**:1186–91.
- 17 **Cook RJ,** Farewell VT. Conditional inference for subject-specific and marginal agreement: two families of agreement measures. *Can J Stat* 1995;**23**:333–44.
- 18 **Meade MO,** Cook RJ, Guyatt GH, et al. Interobserver variation in interpreting chest radiographs for the diagnosis of acute respiratory distress syndrome. *Am J Respir Crit Care Med* 2000;**161**:85–90.
- 19 **Bauchner H,** Vinci R, Bak S, et al. Parents and procedures: a randomized controlled trial. *Pediatrics* 1996;**98**:861–7.
- 20 **Woodward GA,** Fleegler EW. Should parents accompany pediatric interfacility ground ambulance transports? The parent's perspective. *Pediatr Emerg Care* 2000;**16**:383–90.