respectively.³ Hospital admissions during the same time period totalled approximately 350 and 530 for varicella and zoster, respectively; hospitalisation was most common for those aged <4 years old.³

Of the mothers interviewed, just over 30% (61/200) were aware of the availability of an effective vaccine. Women of childbearing age are an at-risk population and should be a target group for immunisation. Also, the vaccine is available in Ireland upon request and parents may choose to administer to their child. This information needs to be disseminated to parents principally via public health clinics and general practitioner practices.

Mothers interviewed were from a cross-section of Irish society (table 1). Education level varied, the majority agreeing to partake had completed at least secondary level education, 70% had completed some form of third-level education. Willingness to comply with vaccination policies varied according to social grouping. It is generally accepted that women of a higher educational background are more questioning of vaccination policies while those of lower income tend to be more trusting of health-care providers.⁴ The majority of mothers (91%) would have agreed to have their child vaccinated were universal recommendations in place despite socioeconomic or educational grouping, with most willing to pay independently of insurance status.

The most useful information obtained from this study was that if the varicella vaccine were introduced, according to the figures obtained from this study, 91% of mothers interviewed would vaccinate their child, with the possibility of another 4% (table 3). Despite the small numbers interviewed, there was a cross-section of society sampled, and if given the option, the majority would have been happy for their child to be immunised with this safe and effective vaccine.

V O'Dowd, M O'Grady, E Moylett

Academic Department of Paediatrics, National University of Ireland, Galway, Ireland

Correspondence to: Edina Moylett, Department of Paediatrics, Clinical Science Institute, National University of Ireland, Galway, Ireland; edina.moylett@nuigalway.ie

Acknowledgements: We would like to thank the mothers on the postnatal ward for their willingness to participate, and the nursing staff on the postnatal ward of University Hospital Galway for their assistance.

Competing interests: None declared.

Accepted 6 February 2009

Arch Dis Child 2009;**94**:648–649.
doi:10.1136/adc.2009.159111

REFERENCES

1. **Marin M**, Meissner HC, Seward JF. Varicella prevention in the United States: a review of successes and challenges. *Pediatrics* 2008;**122**:e744–51.
2. **Royal College of Physicians of Ireland**. National Advisory Immunisation Committee. *Immunisation guidelines for Ireland*. 2008 edn. <http://www.ndsc.ie/hpsc/> (accessed 27 Apr 2009).
3. **Kavanagh P**, O'Flanagan D. Varicella vaccine: Policy options for Ireland. *Epi-Insight* June 2008;**9**:2–3. <http://www.ndsc.ie/hpsc/> (accessed 27 Apr 2009).
4. **Moseley KL**, Clark SJ, Gebremariam A, *et al*. Parents' trust in their child's physician: using an adapted Trust in Physician Scale. *Ambul Pediatr* 2006;**6**:58–61.

CORRECTION

doi:10.1136/adc.2007.127233corr2

Pijpers MAM, Tabbers M, Benninga MA, *et al*. Currently recommended treatments of childhood constipation are not evidence based. A systematic literature review on the effect of laxative treatment and dietary measures. *Arch Dis Child* 2009;**94**: 117–31.

Two errors were noticed by the authors in their systematic review of the literature on treatment of childhood constipation. These errors did not affect the study's conclusions. Here is the corrected data. Corrections are in *italics*.

Table 4b PEG compared to lactulose

Study (quality)	Intervention	Controlle intervention	Outcome measure	Results	Efficacy
Gremse <i>et al</i> ²³ , 2002 (LQ)	PEG 3350, 10 g/m ² /day	Lactulose, 1.3 g/kg/day	Mean defecation frequency/2 weeks	I: 14.8 ± 1.4 C: 13.5 ± 1.5 (p < 0.05)	More effective
			Global assessment of treatment succes	I: 31/37 (84%) C: 17/37 (45.9%) (p = 0.002)	More effective

Table 4i Laxatives investigated in one single study

Study (quality)	Intervention	Controlle intervention	Outcome measure	Results	Efficacy
Bongers <i>et al</i> ¹⁵ , 2007 (HQ)	New formula with high concentration of sn-2 palmitic acid, a mixture of prebiotic oligosaccharides and partially hydrolysed whey protein (Nutrilon Omneo)	Standard formula (Nutrilon 1)	Mean defecation frequency/week	I: 5.6 ± 2.8	Not significant
				C: 9 ± 2.5 (p = 0.36)	
			Improvement of hard to soft stools	I: 9/10 (90%) C: 5/10 (50%) (p = 0.14)	Not significant