Abstract 563 Figure 1 NMR

Results First treatment was initiated with 2 immunoglobulin's cycles and pulse therapy with methylprednisolone not getting satisfactory motor response. The patient developed with respiratory effort and hypoventilation (PCO₂ max - 90mmHg) becoming necessary to introduce a course of 5 days of non invasive ventilation. Subsequently, a 5-day plasmapheresis cycle was initiated, solving the clinical case with an increasing recovering of motor and ventilation function after 1 week of treatment.

Conclusion Enterovirus infections can cause several clinical manifestations such as transverse myelitis. In most of transverse myelitis cases the treatment is immunoglobulin and pulse therapy with methylprednisolone, yet this case was only solved after plasmapheresis cycle.

564 AUTOIMMUNE ENCEPHALOPATHY NINE YEAR OLD BOY WITH NO HISTORY OF INTEREST

doi:10.1136/archdischild-2012-302724.0564

C Perez Moreno, MDM Campillo. *Paediatric Area, Health Centre, Altea, Spain*

Background and Aims Nine year old boy admitted with meningeal syndrome. six days ago began with fever being treated with azithromycin. three days before entering a high fever, bilious vomiting, no diarrhea, the next day associated dizziness, headache, drowsiness, the day before admission diplopia and pain associated neck.

Strikes disease, no rashes or petechiae.

Methods Hemogram: 13.24 leukocytes, neutrophils 87.4%

Cerebrospinal fluid red blood cells 160/mm³ 30/mm³ polynuclear leukocytes 5%, lymphocytes 95%, Gram negative, PCR Herpes simplex I and II negative, negative enterovirus, varicella zoster negative. Negative blood cultures, Mantoux negative.

Abnormal EEG tracing during wakefulness slow waves of high amplitude delta, acute, which are located in anterior and temporal area.

RMN ill-defined hyperintense areas in pons, cerebellum, basal ganglia, right parietal subcortical white matter. suspected encephalitis retrovirus by areas of parenchymal signal alteration above and infratentorial level.

Three days later reduction in the number and extent of intraxial lesions above and infratentorial.

Results The clinical and resonancia were doing suspect herpes virus encephalopathy, so income at the start of treatment with intravenous acyclovir. When we receive negative results and the improvement of symptoms, treatment it was suspended treatment with acyclovir on the fifteenth day and start treating autoimmune encephalitis with five boluses of methylprednisolone one gram every 24 hours.

Conclusions The day of discharge was treated with 60 mg of prednisolone daily. the fever subsides completely within three days before discharge, and intention tremor persists discrete gait instability.

565 REFLECTION ON A CASE OF DOPAMINE-RESPONSIVE DYSTONIA

doi:10.1136/archdischild-2012-302724.0565

ML Neamtu, L Dobrota. *Pediatric Clinic Hospital Sibiu, Lucian Blaga University of Sibiu, Sibiu, Romania*

Background Characteristic symptoms of Dopamine-responsive dystonia (DRD) are increased muscle tone and Parkinsonian features. Children with DRD are often misdiagnosed. The disorder responds well to treatment with Levodopa.

Aim To reflect on a case of DRD.

Method Case report of a 13 years old girl misdiagnosed with tetany.

Results The girl was hospitalized for opisthotonus, positive Trousseau and Chvostek signs, diagnosed as tetany. The laboratory analysis have shown: normal serum of calcium (2.28 mmol/l), normal serum of magnesium (0.80 mmol/l), normal serum of phosphor (1.26 mmol/l), normal alkaline phosphatase (261 u/l) and normal PTH (27.9 pg/ml). Although the initial evolution was favorable (with intravenous calcium gluconate), the hypoparathyroidism diagnosis requiring reconsideration. The final diagnosis was DRD with long good evolution after Levodopa treatment. Referring to family history we learned that the patient have a cousin with the same symptoms.

Conclusions The misdiagnosis results from the following similarities: increased muscle tone with opisthotonus, writer's cramp with Trousseau sign, facial dystonia with Chvostek sign and difficult speech (due to facial dystonia) with patient illiteracy. All these similarities delayed the DRD diagnosis.

566 KLIPPEL TREUNANAY SYNDROME IN DIFFERENTIAL DIAGNOSIS OF CEREBRAL PALSY

doi:10.1136/archdischild-2012-302724.0566

¹P Karakaya, ¹Y Topçu, ¹E Bayram, ¹U Yiş, ²H Çakmakçı, ¹S Hız. *¹Pediatric Neurology; ²Radiology, Dokuz Eylül University, Izmir, Turkey*

In cerebral palsy (CP) atrophy of the paretic body half results in disturbed growth. Disturbed growth is also a feature of a rare disorder; Klippel Treunanay syndrome (KTS). Here we report a child with an initial diagnosis of CP because of limping and thinning of the extremities on the right side who had a final diagnosis of KTS. Five year old male was admitted to our department of Pediatric Neurology. He had been followed up with the diagnosis of CP since he started walking because of limping and thinning of the extremities on the right side of his body. His perinatal and natal period was uneventful. Developmental milestones were normal. On physical examination hypertrophy of the left upper and lower extremities

with widespread port wine stain on his face, lower lip, arms, legs and trunk were noted. Brain magnetic resonance imaging revealed vascular malformations located periventricularly and adjacent to the corpus callosum. On brain magnetic resonance angiography ectatic, varicose deep veins (venous malformations) were detected. Lower extremity MRI showed hypertrophy of bones and soft tissues on the left side. As the child has capillary malformations (port wine stain), soft tissue and bone hypertrophy and vascular malformations a diagnosis of KTS was made. KTS consists of two major features; congenital vascular malformations and disturbed growth. For diagnosis presence of either capillary or venous malformations with disturbed growth of the bone or soft tissues is required. Children with the diagnosis of CP should be carefully examined for any finding suggesting a genetic disease.

567 NEAR FATAL CASE OF AMLODIPINE POISONING IN AN INFANT

doi:10.1136/archdischild-2012-302724.0567

SM Nimbalkar, DV Patel. *Department of Pediatrics, Pramukhswami Medical College, Anand, India*

Background and Aims Infants contribute about 5.25% of total poisoning exposures in humans. We report first case of infant to have survived Amlodipine intoxication.

Methods 11 month old infant received 12.5 times the maximum therapeutic dose of amlodipine as a result of a medication error. He presented with vomiting, lethargy, breathlessness, muffled heart sounds and progressed to hypotensive shock within hours of admission. He received mechanical ventilation, fluid therapy with normal saline and inotropes. Peripheral pulses remained feeble and blood pressure was 70/40 mm of Hg. High dose insulin infusion at 0.5 u/kg/hr with intravenous 25% dextrose and intramuscular glucagon for maintaining euglycemia was given. Continuous infusion of calcium gluconate at 0.5 mEq/kg/hr was started simultaneously. To manage prerenal failure, oliguria and congestive cardiac failure induced pulmonary oedema peritoneal dialysis was initiated.

Result Infant improved rapidly after insulin and dextrose infusions (for 15 hours) along with glucagon and calcium gluconate infusion (for 72 hours) was initiated. Calcium channel blockers (CCB) are phenylalkylamines. CCB act by binding to the α unit of the L Type calcium channels. Insulin secretion being calcium dependent, blockage of the L type channel results in impaired insulin secretion causing hypoglycemia. CCB poisoning results in insulin resistance. A continuous increase in blood sugar levels may predict sudden decline in hemodynamic variables rather blood pressure and pulse measurements.

Conclusion Hyperinsulinaemia/euglycaemia therapy is beneficial in seriously intoxicated patients with CCB induced hypotension, hyperglycemia, and acidosis. There may be benefit in using them in combination with other standard therapies.

568 NEONATAL HEMOLYTIC UREMIC SYNDROME

doi:10.1136/archdischild-2012-302724.0568

L Tiwari, N Vashisht, P Kaur, N Gupta, B Saikia, J Puliye. *Paediatrics and Neonatology, St. Stephens Hospital, Delhi, India*

Haemolytic uremic syndrome (HUS) is characterised by the triad of non immune haemolytic anaemia, thrombocytopenia and renal failure. It is rare in newborns. Only 5 cases of neonatal HUS have been reported to-date. We describe a series of 2 cases of neonatal HUS where one had definite history of birth asphyxia and another was an abandoned baby without clinical details about perinatal period. The known modalities of treatment include transfusion of plasma and plasmapheresis. In first baby haematological and renal parameters improved after double volume exchange transfusion while second

baby responded to transfusion of fresh whole blood (packed red cells + plasma). Exchange transfusion essentially provides plasma exchange besides augmenting the haemoglobin levels. Arguably this is the first report of neonatal HUS treated with exchange transfusion.

569 THORACOSCOPIC SYMPATHECTOMY FOR LONG-QT SYNDROME IN A SICKLE CELL ANEMIA PATIENT

doi:10.1136/archdischild-2012-302724.0569

¹B Banieghbal, ²J Harrisberg. *¹Pediatric Surgery, Netcare Sunninghill Hospital; ²Pediatric Cardiology, Walter Sisulu Cardiac Centre for Africa, Netcare Sunninghill Hospital, Johannesburg, South Africa*

Background and Aim Long QT syndrome (LQTS) is a relatively common disease, however, it is rarely described in Black African patients. We would like to present a west African patient with sickle cell disease and LQTS who was treated with thoracoscopic left sympathectomy and 1/3 lower stellectomy.

Methods An 11-year old boy with sickle cell anemia (Hb=7.2g/dl) was also found to have LQTS. He has had >15 defibrillations in 5 years. He was on high dose beta-blockers but required a number of shocks despite it. Implantable defibrillator was rule out due to his home country lack of cardiac surgical expertise in case of complications. He was considered to be a suitable candidate for sympathectomy. This was done through a 3 trocar approach to the left upper chest. Sympathetic chain including stellate ganglion was exposed. 20mls of 0.5% lignocaine was sprayed over the ganglions. Prior to excision of sympathetic chain, all visible nerves traveling medially (to the heart) was excised with hook diathermy. Lower 1/3 of stellate ganglion as well as T2-T4 ganglia was resected en-bloc and submitted for histology.

Results There were no per- or post-operative complications and no evidence of Horner's syndrome post surgery. The patient was discharged 72 hours later. QTc interval before surgery was 532ms and reduced to 448ms prior to discharge.

Conclusion Sympathectomy including lower portion of stellate ganglion is a viable alternative for patient with LQTS.

570 PULMONARY HYPERTENSION AS A COMPLICATION OF VON RECKLINGHAUSEN DISEASE

doi:10.1136/archdischild-2012-302724.0570

LT Celik, K Papakostas, A Artmann, TP Le. *Pediatric Cardiology, Klinikum Links der Weser, Bremen, Germany*

Introduction Neurofibromatosis (NF) is occasionally complicated by systemic vasculopathy such as myocardial infarction, renovascular hypertension etc. The appearance of pulmonary embolism (PE) is not described until yet.

Case Report 15-year-old female patient with neurofibromatosis type I. Progressive dyspnoea, general fatigue and chest pain beginning 6 days before hospital admission. Anamnestic no clinical sign for deep vein thrombosis. No oral contraceptive, non smoker. Laboratory findings: Elevated troponin, plasma brain natriuretic peptide (BNP) and D-dimer. Normal ECG, echocardiogram showed right ventricular and atrial enlargement with moderate tricuspid regurgitation and estimated right ventricular systolic pressure at 45 mm Hg and dilated main pulmonary artery. Left ventricular function was normal.

Chest x ray showed right heart enlargement but normal lung structure. Lung CT revealed massive PE in left and right branches and a mediastinal tumor. Lysis with Alteplase was given for 4 days. Normalisation of laboratory and echocardiogram. Because of the tumor angiographic-MRI was done for follow-up and showed normal flow pattern in pulmonary vasculature. The tumor was classified as a ganglioneuroma. Genetic thrombophilia testing was negative,