an incidence around 1/100 000/year. We studied the characteristics and outcome in PICU patients with ICH.

Methods Children with ICH admitted to PICU during 2000–2010, were retrospectively studied. Clinical information was abstracted via chart review.

Results 21 consecutive cases, aged 5.6 ± 4.5 years, 12 girls, were analyzed. 70% of the children presented with vomiting, 55% with seizures, 38% with headache and 9.6% with focal neurological signs. Mean Glasgow Coma Scale (GCS) before intubation was 7.33 ± 2.45 . In 66.6% emergent evacuation of hematoma or hydrocephalus at admission was performed. 23.5% had an arteriovenous malformation (AVM) and 15% of these children underwent embolization for AVM obliteration. Mortality was found 19%. Patients who died had lower GCS (4.75 ± 2.06 vs. 7.9 ± 2.16), $P < 0.05$, higher PRISM III-12 (first 12 hours from admission), (21.5 ± 8.1 vs. 7.8 ± 7.7), $P < 0.005$, higher PRISM III-24 (next 12 hours), (17.2 ± 8.3 vs. 5.2 ± 5.1), $P < 0.001$, and longer α PTT, $P < 0.01$ than those who survived. Patients who needed inotropic support the 1st day of PICU stay had 12 times greater mortality than children who didn't need inotropic support. Among survivors 68.7% presented neurologic deficit at PICU discharge. Children with neurologic deficit had lower GCS (6.8 ± 1.6 vs. 10 ± 1.5), $P < 0.005$ than those without deficit.

Conclusions As outcome is dismal in children with ICH and critical illness at presentation, low GCS, high PRISM III and need for inotropic support, prompt diagnosis seems essential to improve prognosis in these children.

811 USEFULNESS OF THE PRISM III SCORE TO PREDICT OUTCOME IN CRITICALLY ILL CHILDREN WITH VAP

doi:10.1136/archdischild-2012-302724.0811

S Stabouli, E Volakli, A Violaki, A Tsolaki, M Dimitriadou, K Skoumis, M Sdouga. *PICU, Hippokraton Hospital, Thessaloniki, Greece*

Background and Aims Ventilator-associated pneumonia (VAP) is associated with increased length of stay and adverse outcomes in PICU patients. In a retrospective study, we examined if PRISM III score at admission or at the day of VAP development could better predict the outcome in patients with VAP.

Methods The medical records of PICU patients admitted to a 8-Bed PICU of a tertiary-care hospital from January–December 2011 were reviewed. Clinical data, PRISM III score at admission or at the day of VAP development were recorded. VAP was diagnosed according to CDC criteria.

Results 27 patients, mean age 4.40 ± 4.23 years, 59.3% boys, developed VAP. 4 patients presented 2 VAP episodes. Mean PRISM III score at admission was 10.19 ± 7.65 , at the day of first VAP episode 7.31 ± 6.94 , and at the day of 2nd VAP episode 4.75 ± 3.60 . The receiver operator characteristic curve (ROC) analysis showed that PRISM III at admission could better predict mortality in PICU patients with VAP than PRISM III at the day of VAP episode. (Figure 1).

Abstract 811 Figure 1

The area under the curve was found 0.85 (asymptotic 95%CI 0.59 to 1, $P < 0.01$) for PRISM III at admission and 0.72 (95%CI 0.48 to 0.95, $P = 0.081$) for PRISM III at VAP episode.

Conclusions PRISM III at admission could better predict mortality in PICU patients with VAP than PRISM III at the day of VAP episode suggesting that VAP may not independently affect mortality.

812 CARDIOPULMONARY ARREST IN PEDIATRIC EMERGENCY CARE AND INTENSIVE CARE: A MULTICENTER STUDY IN TURKEY

doi:10.1136/archdischild-2012-302724.0812

¹T Kendirli, ²N Erkek, ³T Köroğlu, ⁴B Bayrakçı, ⁵A güzel, ⁶A Çitak, ⁷D Demirkol, ⁸H Ağın, ⁹AE Arslanköylü, ¹⁰NO Kutlu, ¹¹Ş Paksu, ¹²AB Anıl, ¹³G Kalkan, ¹⁴D Yıldızdaş, ¹⁵M Duman, ¹⁶R Dündaröz, ¹⁷N Aşlıoğlu, ¹⁸A Yaman, ¹⁹S Kesici, ²⁰D Tekin, ²¹O Dursun, ²²Ç Ödek, ²³C Ateş, ²⁴HL Yılmaz, ²⁵E İnce, ²⁶M Karaböcöoğlu. ¹Pediatric ICU, Ankara University; ²Dr. Sami Ulus Hospital, Ankara; ³Dokuz Eylül University, İzmir; ⁴Pediatric ICU, Hacettepe University, Faculty of Medicine, Ankara; ⁵Pediatric Emergency Care, Ondokuz Mayıs University, Samsun; ⁶Pediatric ICU, Istanbul University Istanbul Medical Faculty; ⁷Pediatric ICU, Bezmialem University, Istanbul; ⁸Pediatric ICU, Behçet Uz Children's Hospital, İzmir; ⁹Pediatric ICU, Mersin University, Mersin; ¹⁰Pediatric ICU, Meram Faculty of Medicine, Konya; ¹¹Pediatric ICU, Ondokuz Mayıs University, Samsun; ¹²Pediatric ICU, Tepecik Children's Hospital, İzmir; ¹³Pediatric ICU, Gazi University Faculty of Medicine, Ankara; ¹⁴Pediatric ICU, Çukurova Faculty of Medicine, Adana; ¹⁵Pediatric emergency Care, Dokuz Eylül University, İzmir; ¹⁶Pediatric Emergency Care, Bezmialem University, Istanbul; ¹⁷Pediatric ICU; ¹⁸Pediatric Emergency Care, Ankara University, Faculty of Medicine, Ankara; ¹⁹Pediatric ICU, Akdeniz University, Antalya; ²⁰Biostatistics, Ankara University, Faculty of Medicine, Ankara; ²¹Pediatric Emergency Care, Çukurova University Faculty of Medicine, Adana, Turkey

Background and Aim The most cause of cardiopulmonary arrest (CPA) is respiratory system disorders. Usually the survive from CPA is 30% in hospital and under 10% in out of hospital. The aim of this study, the cause of CPA, applications and results of CPA in pediatric ICU and emergency care in Turkey.

Methods This study conducted between January 15 and July 15, 2011, multicenter, prospective, observational from Turkey.

Results We enrolled 239 children whose CPA developed. Fifty-four percent of all patients were boy and their mean age were 42.4 ± 58.1 months. The causes of CPA were respiratory failure in 49.8%, sepsis in 30.1%, cardiac disease in 21.3% and rhythm disorders in 8.8%. The place of CPA occurred were PICU in 68.6%, services in 18%, out of hospital in 10% and emergency care in 3.3% of patients whose CPA developed. Adrenalin was performed in 221, defibrillation in 16 and automatic external defibrillation in patients. Mean resuscitation time was 30.7 ± 23.6 minutes. Return percent after first resuscitation application was 44.8%. We check to mortality rate after first resuscitation 43.3% in PICU, 41.9% in services, 50% in Emergency Care, 41.7% at out of hospital ($p = 0.539$). The 83% of them were unconsciousness, renal replacement therapy was applied in 16 patients. After first resuscitation, 54.2 patient survived and neurologic sequelae was in 32% of them.

Conclusion Mortality and morbidity are higher either hospital and out of hospital CPA, therefore prevention to CPA and well resuscitation applications are very important.

813 PALLIETERBURGH: DEVELOPMENT OF A HIGH DEPENDENCE TRANSITIONAL CARE UNIT

doi:10.1136/archdischild-2012-302724.0813

¹M Smit-van den Berg, ²C Joosen, ³D Tibboel, ⁴E Ista, ⁵S Gischler. ¹PICU; ²PICU and Department of Pediatric Surgery, Erasmus MC - Sophia Children's Hospital, Rotterdam, The Netherlands

Background Ongoing advances in paediatric intensive care led to increased survival, with increased morbidity and long-lasting sequelae.