There was no significant correlation between the number of wheezing attacks ($p>0.05$).

No correlation between vitamin D levels and acute bronchiolitis was found but this may be the result of few numbers of patients that we studied with.

829 EVALUATION OF IL-6 AND HIGHLY SENSITIVE CRP VALUE IN CSF AND SERUM CHILDREN REFERRED TO PEDIATRIC EMERGENCY ROOM

doi:10.1136/archdischild-2012-302724.0829

¹A Hamed, ²H Ayatollahi. ¹Pediatrics; ²Mashhad University of Medical Sciences, Mashhad, Iran

Introduction Acute bacterial meningitis which is a pediatric emergency with high mortality and morbidity, must be diagnosed and treat promptly. Determination of some inflammatory mediators example IL-6 and hsCRP were useful in differential diagnosis of bacterial and viral meningitis.

Objectives Determination of hs CRP and IL6 in serum and CSF in children suspected meningitis.

Materials and Methods From of children that hospitalized in pediatric emergency ward and for they performed lumbar puncture, 1cc serum and csf of they were taken to laboratory ward and have measured IL-6 with Elisa method and hsCRP with Immunoturbidimetry method patients were followed up and Finally we compared levels of this two mediators.

Results From 81 cases, 27 cases (33/3%) were bacterial meningitis 27 cases (33/3%) were viral meningitis and 27 cases (33/3%) were Normal. IL-6 concentration in the csf and serum were significantly raised in cases of bacterial meningitis ($P=0/001$, $P=0/01$) but hsCRP concentration in the CSF and serum we not significantly raised in cases of bacterial meningitis ($p=0.46$, $p=0.29$). Mean IL-6 in serum in bacterial meningitis was (50.01) and in viral meningitis was (10.64) mean IL-6 concentration in the CSF in bacterial meningitis was (180.74) and in viral meningitis was (39.08) mean hsCRP in CSF in bacterial meningitis was (2.22) and viral meningitis was (1.29).

Conclusion The measurement of IL-6 in the CSF and serum is potentially a very useful diagnostic tool for differential diagnosis of bacterial and viral meningitis.

830 THE VALUE OF MEAN PLATELET VOLUME IN THE DETERMINATION OF COMMUNITY ACQUIRED PNEUMONIA IN CHILDREN

doi:10.1136/archdischild-2012-302724.0830

¹E Karadag Oncel, ¹Y Ozsurekci, ¹A Kara, ²S Karahan, ¹AB Cengiz, ¹M Ceyhan. ¹Pediatric Infectious Disease, Hacettepe University Ihsan Dogramaci Childrens' Hospital; ²Biostatistics, Hacettepe University Faculty of Medicine, Ankara, Turkey

Background The aim of this study was to evaluate whether mean platelet volume (MPV) could be used for the diagnostic tool of community acquired pneumonia (CAP) and for making the decision for hospitalization.

Methods The computerized records of children aged 1 to 18 years who were diagnosed with CAP based on WHO criteria were evaluated. A standard protocol was followed, and patients with severe CAP were hospitalized. CAP patients were divided into two groups based on disease severity. The control group consisted of age- and gender-matched healthy children who attended the "Well-child" clinic during the study period. Values for hemoglobin, white blood cell count (WBC), platelet count, MPV and C-reactive protein (CRP) obtained on first presentation were recorded for each patient.

Results A total of 139 patients were diagnosed with CAP during the study period, 82 (58.9%) of which had severe disease, which required

hospitalization, while the remaining 57 (41.4%) were followed-up as outpatients. The control group consisted of 61 healthy children. Patients with CAP had lower MPV values than their healthy counterparts (7.14 ± 0.70 vs. 8.35 ± 1.26 fL; $p<0.001$). Following ROC analysis, the cut-off value established for MPV was 8.05, which had a sensitivity of 91% and a specificity of 51% for making a diagnosis of CAP. Patients with severe CAP that required hospitalization had significantly higher CRP levels than those who were followed-up on an out-patient basis ($p<0.001$).

Conclusions Our findings suggest that MPV could be used for the diagnosis of CAP in children.

831 AN ANALYSIS OF THE PROGRESSIVE CLINICAL DETERIORATION IN THE CASES WITH SUBACUTE SCLEROSING PANENCEPHALITIS

doi:10.1136/archdischild-2012-302724.0831

S Saygili, C Yazar, G Arslan, E Aksuyek. *Pediatrics, Turkish Ministry of Health, Kanuni Sultan Suleyman Training and Research Hospital, Istanbul, Turkey*

Subacute sclerosing panencephalitis (SSPE) is devastating progressive neurodegenerative disorder of the central nervous system by an aberrant measles virus. SSPE is still important health problem in countries where measles immunization is incomplete and in world regions where genetic polymorphism to this particular infection is present. We aimed to make a retrospective analysis of the clinical and laboratory findings and prognostic factors of children with SSPE.

Method A total 55 consecutive cases (22 female, 33 male), aged 5–19 years, followed-up for SSPE at the Pediatric Neurology Department were included. The risk factors for progression time of SSPE from stage I–II to IV according to the criteria of Jabbour examined were gender, age at diagnosis of SSPE, presence of neuroimaging abnormalities, age of patient during measles infection and the antibody index for measles. SSPE risk factors were obtained through a questionnaire via personal interviews and the medical records and were assessed using Mann-Whitney U, Student t and Pearson correlation tests.

Results Presence of neuroimaging abnormalities, gender, age at diagnosis of SSPE, presence of neuroimaging abnormalities, age of patient during measles infection and the antibody index for measles were not associated with a progressive clinical deterioration ($p>0.05$). 66.7 percent of cases were vaccinated with measles vaccine, 43.5 percent of cases have had measles infection before age of 9 months.

Conclusion Viral, host and environmental factors are probably involved in the development and progression of SSPE. Successful measles vaccination programmes which accordance with the requirements of that country can protect the population against SSPE.

832 ACUTE TRANSVERSE MYELITIS IN A PEDIATRIC CASE OF LYME DISEASE

doi:10.1136/archdischild-2012-302724.0832

¹I Erol, ²B Kiliçaslan, ³Ş Demir, ⁴H Gülcan, ¹F Alehan. ¹Department of Pediatrics, Division of Child Neurology; ²Department of Pediatrics; ³Department of Radiology; ⁴Department of Pediatrics, Division of Neonatology, Baskent University Hospital, Adana, Turkey

Background and Aim A broad range of neurologic disorders have been described in children with Lyme disease. In contrast, there are few reports of transverse myelitis in pediatric Lyme disease patients.

Case Report A 12-old girl admitted with the complaints of weakness in legs, sensory loss in lower limb, and backache for 10 days. Mental status, cranial nerve, and upper-extremity motor and sensory examinations produced normal results. The strength in his iliopsoas and hamstring muscles was scored 2/5. Deep tendon reflexes

were absent in both lower extremities. Babinski's signs were not noted. The Romberg sign was present. Magnetic resonance imaging (MRI) of the spinal cord revealed enhancing T₂ hyperintensity within the spinal cord, from T₁₀-L₁. (Figure 1). Cerebrospinal fluid (CSF) showed no pleocytosis and normal protein and glucose concentrations. The CSF did not show oligoclonal banding and immunoglobulin (Ig)G index was normal. Serum serologic evaluation of *Borrelia burgdorferi* was positive for IgM but negative for IgG. CSF serologic evaluation was negative (both ELISA and Western blot). The patient was diagnosed as manifesting acute transverse myelitis. He was treated with intravenous methylprednisolone pulse therapy (1000 mg/day for 5 consecutive days), followed by oral prednisolone (1 mg/kg per day). After the start of steroid therapy, the patient showed gradual clinical improvement and was able to walk on the 30th day of illness. We also administered doxycycline 4 mg/kg per day for 14-days for borreliosis.

Conclusion This case serves as a reminder that acute transverse myelitis can be a rare clinical manifestation of Lyme disease.

Abstract 832 Figure 1

833 THE EFFECT OF PROBIOTICS ON COLONIZATION OF RESISTANT ORGANISMS IN PRETERM INFANTS

doi:10.1136/archdischild-2012-302724.0833

¹A Kurt, ¹A Ecevit, ¹D Anuk Ince, ¹A Abbasoglu, ²O Kurt Azap, ¹Z Ecevit, ³E Ogus, ¹A Tarcan. ¹Pediatrics; ²Infectious Diseases; ³Biostatistics, Baskent University Faculty of Medicine, Ankara, Turkey

Background and Aims To investigate the effect of probiotics on colonization of resistant organisms in preterm infants receiving antibiotics.

Methods This study comprised of preterm infants who were born < 36 weeks and received antibiotic treatment or prophylaxis. Pre-term infants were divided into two groups according to receiving probiotic (*Lactobacillus reuteri*). Stool culture and nasal swab culture were taken to determine colonization.

Results In the group receiving probiotics, mostly colonizing the stool cultures bacteria were *Klebsiella spp.*, *Escherichia coli*, *Enterococcus spp.*, *Enterobacteriaceae spp.*, *Staphylococcus spp.* respectively, and in the group not receiving probiotic mostly colonizing the stool cultures bacteria were *Klebsiella spp.*, *Enterococcus spp.*, *Staphylococcus spp.*, *Escherichia coli*, *Enterobacteriaceae spp.* respectively. When probiotic receiving group compared with not receiving group, proliferation rate of stool cultures was higher in probiotic group. In the groups receiving and not receiving probiotic, proliferation of the nose cultures were similar. Increase in the proliferation rates of weekly stool cultures in probiotic receiving group was statistically significant but there was no statistically difference in the proliferation rates of nose and other cultures that were taken weekly. There was no statistical difference in both groups in the development of resistant organisms.

Conclusions The use of probiotics in neonatal intensive care unit for premature infants who received treatment with antibiotics, did not prevent the colonization of pathogenic microorganisms.

834 THE SENSITIVITY OF BACTERIAL MENINGITIS SCORE IN CHILDREN WITH ACUTE MENINGITIS IN QATAR

doi:10.1136/archdischild-2012-302724.0834

¹H Abdllrahman, ¹A Alhamadi, ¹K Al-Saddi, ¹W Seleem, ¹H Badran, ¹N Sharaf, ²A Farooq. ¹Hamad Medical Center; ²Asptar Medical Center, Doha, Qatar

Background The empirical use of antibiotics in children with suspected meningitis is a common clinical practice worldwide that often leads to drug resistance. It is difficult to clinically differentiate bacterial when compared to viral meningitis until a culture study of cerebrospinal fluid (CSF) Or CSF viral PCR study is performed. A 'wait and see' approach may lead to undesirable outcome. Bacterial Meningitis Score (BMS) is a tool that was developed to help physicians to differentiate between viral versus bacterial meningitis.

Aim To determine the usefulness if any of BMS for discriminating between bacterial or viral meningitis in young children.

Methodology We retrospectively reviewed the charts of all children (from birth till 14 years old) who were admitted with the diagnosis of meningitis to Hamad general hospital in last 2 years period.

Result A total 120 patients (68% boys) with confirmed meningitis were reviewed during the study period. The mean age was (6.3±2.7 year). The majority of patients 112 (93.3%) had viral type meningitis while the remaining had bacterial meningitis (*Strep Pneumonia*, *Neisseria meningitis* and *H. Influenza*). The sensitivity of BMS tool revealed a sensitivity of 100% (95% CI: 75.1 to 100.0) and a specificity of 60.9% (95% CI:50.1-69.7).

Conclusion Our study shows that BMS is a simple, easy and highly sensitive tool that can differentiate bacterial from viral meningitis and it is use may limit the use of unnecessary antibiotic s and hospitalizations.

835 THE VALUE OF A POSITIVE NITRATE TEST IN ROUTINE URINE ANALYSIS FOR THE DIAGNOSIS OF CHILDHOOD UTI

doi:10.1136/archdischild-2012-302724.0835

HA Bin-Nakhi, E Bu-Hamrah, S Sadeq, AR Moodambail. *Pediatrics, Adan Hospital, Kuwait, Kuwait*

Urinary tract infection (UTI) is one of the most common diseases in children. Its diagnosis depends mainly on urine culture. Urinalysis is used as a screening test to exclude UTI. A hospital based, prospective case-review study was carried out at the Pediatric Department, Al-Adan Hospital, State of Kuwait. The objectives of this study is:

1. To determine the proportion of positive urine cultures identified in patient less than 12 years of age, admitted to hospital and was found to have a positive nitrite test in the routine urinalysis.