¹F Abbasi, ²P Vahdani, ³S Korooni, ²A Naderi. ¹Bushehr University of Medical Sciences, Bushehr; ²Infectious Diseases and Tropical Medicine Research Center, Shaheed Beheshti Medical University, Tehran; ³Shiraz University of Medical Sciences, Shiraz, Iran

Background and Aims Hydatidosis, caused by *Echinococcus granulosus*, is an endemic parasitic disease. The most frequent anatomic locations are liver and lung. Dogs and other canines are the primary definitive hosts for this parasite. Hydatid cyst may develop after accidental ingestion of tapeworm eggs, excreted with the feces of these animals. Diagnosis is usually based on radiological and serological findings.

Patient We present a 6 Year-Old-Boy with abdominal pain since months ago. He had history of travelling to village and exposure to cattle and dog. Physical examination of head and neck, chest, extremities, and neurologic exam were normal. In abdominal Physical examination he had mild right upper quadrant tenderness without rebound tenderness. Liver was palpable 3 Cm below costal margin. Abdominal sonography and CT scan showed multiple huge cysts in liver that occupied almost all the liver space. Serum hydatid cyst antibody was positive. With diagnosis of hydatosis surgery was done and albendazole started for him. The patient improved after a while and he was discharged with good general condition.

Conclusion Hydatid cyst should be considered in every patient with liver cystic lesion in endemic area.

958 FATAL HUMAN BOCAVIRUS INFECTION IN A BOY WITH IPEX-LIKE SYNDROME AND VACCINE-ACQUIRED ROTAVIRUS ENTERITIS AWAITING STEM CELL TRANSPLANTATION

doi:10.1136/archdischild-2012-302724.0958

¹D Klinkenberg, ¹R Kobbe, ²R Schneppenheim, ¹I Müller, ¹M Blohm, ³M Malecki, ³V Schildgen, ³O Schildgen. ¹Clinic of Paediatrics; ²Clinic for Paediatric Haematology and Oncology, University Medical Center Hamburg-Eppendorf, Hamburg; ³Institute of Pathology, Kliniken der Stadt Köln, Köln, Germany

We report about a 21-month-old boy presenting with chronic diarrhea and obstructive lung disease since infancy. His older brother, the first male child of non-consanguine Egyptian parents, had died at the age of 20 months suffering from BCGitis and severe CMV infection, suggesting a severe primary immunodeficiency syndrome.

We found immunological dysregulation, endocrine dysfunction and enteropathy compatible with IPEX syndrome, therefore the child was listed for stem cell transplantation (SCT). *FOXP3* gene showed no IPEX associated mutations, but sequencing *IL2RG* we found an mutation in Exon 2 associated with x-linked IPEX-like SCID (c.252C>A, p.N84K).

The pulmonary condition of the boy deteriorated and he was admitted on the ICU, where he was mechanically ventilated ever since. After common respiratory infections were excluded, chronic human bocavirus (hBoV) was identified by multiplex-PCR as the primary causative pulmonary agent causing his respiratory failure. hBoV was found in decreasing quantity in respiratory material, blood and stool specimen.

Interestingly, chronic rotavirus shedding was notified in repetitive stool specimen. As the boy had been vaccinated against rotavirus, vaccine-acquired chronic infection was suspected and confirmed by typing of a vaccine virus specific gene variant.

The boy died before SCT could be performed.

Since its discovery, there is an ongoing discussion if hBoV can cause serious infections or represents only a harmless "bystander". This case report shows that hBoV can result in lethal respiratory infections in immunocompromised children.

959 MYCOPLASMA PNEUMONIAE ASSOCIATED WITH STEVENS JOHNSON SYNDROME IN TWO CHILDREN

doi:10.1136/archdischild-2012-302724.0959

L Sfaihi, K Baklouti, I Maaloul, A Hakim, H Aloulou, T Kamoun, M Hachicha. *CHU Hedi Chaker, Sfax, Tunisia*

Background Mycoplasma pneumonia (MP) infections are often asymptomatic but can involve multiple organ systems. Secondary skin reactions are common, although few patients infected develop Stevens Johnson syndrome (SJS).

Results We describe 2 cases of Mycoplasma pneumoniae chest infection associated with Stevens Johnson syndrome. The two patients had prodromal symptoms of an upper respiratory tract infection before the onset of the eruption. The patients had extensive epidermal bullous vesicles, oropharyngeal and genital ulceration, injected conjunctivae and sclera and swollen lips with flaccid bullae. The mycoplasma IgG and IgM titers returned positive, and blood cultures and other titers were negative. They were successfully treated with macrolides and glucocorticoids.

Conclusion Although the clinical course may be severe and prolonged, the prognosis is uniformly good with complete recovery.

960 SECONDARY PSEUDOHYPOALDOSTERONISM DUE TO PYELONEPHRITIS: TWO CASES REPORT

doi:10.1136/archdischild-2012-302724.0960

¹I Llana, ²MD Sanchez-Redondo, ¹M Benedit, ²J Cobas, ¹MT Garcia de Alvaro. ¹Hospital HM Torrelodones, Madrid; ²Hospital Virgen de la Salud, Toledo, Spain

Secondary pseudohypoaldosteronism occurs due to transient resistance in renal tubules causing renal sodium loss, hyponatremia and hyperkalemia. This may resemble congenital adrenal hyperplasia. This is to be considered specially when a urinary tract infection is present as treatment and prognosis varies.

We report two infants who developed pseudohypoaldosteronism secondary to acute tract urinary infection and urologic malformations.

Patient 1 27 days old infant, weight: 4056 gr. He suffers severe dehydration, hyponatremia 119 mmol/L, K 7 mmol/L and metabolic acydosis, normal serum creatinine and normal 17 OH progesterone. Pathological urine analysis and high levels of serum rennin and aldosterone. Urine culture: E.Coli. We can also appreciate left Ultrasonography hidronefrosis IV/V and primary megaureter. He was treated with Ampicillin + gentamicin and ions stabilisation within the following 36 hours after hospital admission.

Patient 2 14 days old infant, 2510 gr. Admitted at emergencies with a severe dehydration. Hyponatremia and hyperpotasemia that needed urgent treatment and severe metabolic acydosis. Electrolytic disorders were amended, then a urine culture was done with positive results in E. Coli test. He was treated with ampicillin + gentamicin. The renal scan evinced a bilateral ureteral hydronephrosis. The infant showed a recovery thanks to antibiotics treatment.

Conclusions An endocrinological evaluation is necessary to a diagnosis but we mustn't forget the urinary cultures.

Most probable cause is the immature renal responsiveness to aldosterone in three first months of life when exist urinary tract anomalies and/or urinary tract infection.

This secondary pseudohypoaldosteronism is reversible with early treatment.

961 THE DIAGNOSTIC VALUE OF NEUROIMAGING IN RECOGNIZING AND THERAPY MONITORING OF ENCEPHALITIS AND MENINGITIS IN CHILDREN

doi:10.1136/archdischild-2012-302724.0961

¹J Surmik, ²E Emich-Widera, ¹J Wróblewska, ¹I Maruniak-Chudek, ³E Kluczevska. ¹Department of Neonatal Intensive and Special Care; ²Department of Neuropediatrics,