Conclusions EOS in term infants is associated with significantly lower maternal and neonatal 25-OHD levels. We also found that levels of 25-OHD in neonates were positively correlated with those in mothers. These data suggest that adequate vitamin D supplementation during pregnancy may be helpful to prevent EOS in term neonates.

PS-222 STAPHYLOCOCCUS CAPITIS IN NEONATAL LATE-ONSET SEPSIS: UNEXPECTED WORLDWIDE DISSEMINATION OF AN ENDEMIC MULTI-RESISTANT CLONE

¹M Butin, ²P Martins Simoes, ³H Lemriss, ³S Lemriss, ²F Vandenesch, ⁴O Claris, ⁴JC Picaud, ¹JP Rasigade, ¹F Laurent. ¹Clinical Microbiology, Hospices Civils de Lyon, Lyon, France; ²U1111 - CIRI, Inserm, Lyon, France; ³Laboratoire de Recherche Et d'Analyses Médicales, Fraternelle de La Gendarmerie Royale, Rabbat, Morocco; ⁴NICU, Hospices Civils de Lyon, Lyon, France

10.1136/archdischild-2014-307384.521

Background Multi-resistant *Staphylococcus capitis* NRCS-A is involved in late-onset sepsis (LOS) in French NICUs.

Aims To investigate the geographical distribution of NRCS-A, and to precise its susceptibility profile.

Methods Twelve *S. capitis* isolates from distant NICUs (Australia, Belgium, France, United Kingdom, n = 3 each) and 2 *S. capitis* isolates from adult patients were analysed using PFGE, SCCmec typing, dru-typing, a MLST-like analysis, and antimicrobial susceptibility testing. To explore impact of vancomycin selective pressure, after 15 daily subcultures with vancomycin, we determined vancomycin, daptomycin and linezolid minimal inhibitory concentrations (MICs).

Results All NICU *S. capitis* (i) shared >80% similarity of PFGE profile and were similar to NRCS-A profile, (ii) harboured a type V-related SCCmec element, (iii) exhibited the same dru-type, (iv) formed a monophyletic group, (v) harboured a same antimicrobial susceptibility profile including aminoglycosides and methicillin resistance, and vancomycin heteroresistance. These molecular and antimicrobial susceptibility profiles differed from those of adult isolates. An increase of vancomycin and daptomycin (but not linezolid) MICs was observed, significantly faster ($p < 0.05$) for NRCS-A isolates than other tested strains.

Conclusion Our analysis demonstrates an unexpected worldwide distribution of *S. capitis* NRCS-A, specifically in NICUs. Recently, we collected complementary NICU isolates belonging to NRCS-A from Norway, Denmark, the Netherlands, USA, Brazil, New Zealand and Canada, confirming the worrisome dissemination of NRCS-A. Its multi-resistant profile and its ability to rapidly adapt to vancomycin selective pressure, constitute a selective advantage to NRCS-A in NICUs, and raise the issue of potential therapeutic failure and the need for alternative antimicrobial regimens.

Neonatology Clinical

PS-223 IMPROVING STAFF COMPETENCE AND CONFIDENCE FOR NEONATAL RESUSCITATION, A QUALITY IMPROVEMENT INITIATIVE

S Hackett, M Cassidy, S Gormally. Neonatal Intensive Care Unit, Our Lady of Lourdes Hospital Drogheda Co. Louth, Drogheda, Ireland

10.1136/archdischild-2014-307384.522

Aims To introduce a Quality Improvement Initiative aimed at instilling confidence and competence with Neonatal Resuscitation in Labour Ward (LW) and Neonatal Unit (NICU) staff.

Methods A Neonatal Resuscitation Initiative Team (NRIT) consisted of LW and NICU midwives, an Advanced Nurse Practitioner (ANP) and a Paediatric Consultant. LW, NICU and paediatric staff were invited to attend a presentation to outline the aims and methodology of the initiative. Emphasis was placed on teamwork with effective leadership and communication. This was followed by a demonstration of 2 low fidelity simulated neonatal resuscitations performed by the NRIT.

Midwifery and NICU staff were then invited to partake in a Simulated Team Response (STR) to 2 simulated resuscitations.

A random selection of participants completed a closed questionnaire before and after the NRIT demonstration and before and after their STR. Levels of confidence in all aspects of neonatal resuscitation, including teamwork and leadership skills were evaluated using a Likert Scale.

Results One hundred and thirteen staff participated in the quality initiative from January 2013 to August 2013. Seventy four staff participated in the NRIT demonstrations and 39 in the STR. Forty three questionnaires were completed following the NRIT and 13 following the STR. Using a Wilcoxon test the pre-post gain in confidence score was statistically significant ($p = 0.0020$) for doctors and midwives.

Conclusion Simulation training is a valuable means of maintaining skills and enhancing confidence for neonatal resuscitation. This study quantifies the positive impact of a Quality Improvement Initiative aimed at improving performance at neonatal resuscitation.

PS-224 VIDEO EDUCATION TO IMPROVE BAG MASK VENTILATION DURING SIMULATED NEWBORN RESUSCITATION

¹P Deindl, ²J Schwindt, ²A Berger, ³G Schmoelzer. ¹Department of Neonatology and Pediatric Intensive Care Medicine, University Medical Center Hamburg-Eppendorf, Hamburg, Germany; ²Department of Pediatrics and Adolescent Medicine Division of Neonatology Pediatric Intensive Care and Neuropediatrics, Medical University of Vienna, Vienna, Austria; ³Department of Pediatrics and Adolescent Medicine Division of Neonatology Pediatric Intensive Care and Neuropediatrics, Royal Alexandra Hospital, Edmonton, Canada

10.1136/archdischild-2014-307384.523

Aim To evaluate if video based education could improve quality of positive pressure ventilation (PPV) performed by novice health care providers during neonatal resuscitation.

Methods Twenty-eight 4th year medical students were randomly paired and instructed to give PPV to a modified manikin as single-person resuscitators, then as two-person paired resuscitators using either an anatomical shaped neonatal face mask with an air cushion rim (IS) or a Laerdal round face mask (LM). After watching a video-tutorial they randomly repeated each mask ventilation performance. Airway pressure, gas flow, tidal volume, and mask leak were recorded. PPV performance quality was analysed using video recording.

Results Mask leak was lower during one-person ventilation when using IS ($56 \pm 16\%$) compared to LM ($71 \pm 19\%$). LM mask leak during one-person ventilation was significantly lower when using the two point top hold in contrast to the ok rim hold (before training: $63 \pm 22\%$ vs. $72 \pm 18\%$, after training: $57 \pm 17\%$ vs. $77 \pm 12\%$, respectively). Watching a video-tutorial improved correct head position (score: 1.4 ± 0.7 vs. $3.8 \pm$

Poster symposium

Abstract PS-224 Figure 1

Abstract PS-224 Figure 2

0.4) and chin lift manoeuvre (score: 2.3 ± 0.2 vs. 3.5 ± 0.2) during PPV, however mask leak was not significantly reduced.

Conclusion A self-instructional educational video on adequate bag mask ventilation significantly improves performance quality scores in novice health care providers.

PS-225 RISK FACTORS (RF) ASSOCIATED WITH ADVANCED NEONATAL RESUSCITATION IN ≥ 34 W GA NEWBORNS: A MULTICENTER, PROSPECTIVE, CASE-CONTROLLED STUDY. THE ADVANCED NEONATAL RESUSCITATION (ANR) STUDY

¹JP Berazategui, ²A Aguilar, ³M Escobedo, ⁴F Althabe, ⁵R Ginsburg, ⁶F De Almeida, ⁷G Albornoz, ⁸F Saker, ⁹MG Puig, ¹⁰D Amado, ¹¹M Valera, ¹²E Szyld. ¹Maternal and Infant, FUNDASAMIN / Hospital Universitario Austral, Pilar - Derqui, Argentina; ²Maternal and Infant, FUNDASAMIN (Fundacion Para La Salud Materno Infantil), Buenos Aires, Argentina; ³Maternal and Infant, University of Oklahoma, Oklahoma, USA; ⁴Maternal and Infant, IECS Instituto de Efectividad Clínica y Sanitaria, Buenos Aires, Argentina; ⁵Maternal and Infant, Escola P. de Medicina da Universidade F. de Sao Paulo, Sao Paulo, Brazil; ⁶Maternal and Infant, Clinica y Maternidad Suizo Argentina, Buenos Aires,

Abstract PS-225 Table 1

Variables	OR	P	95% (CI)
GA < 37wk (GA < 37)	3.0	0.001	(2.0–4.6)
Eclampsia (EC)	6.0	0.049	(1.0–36.6)
Maternal fever during labour (MFL)	7.8	0.001	(3.1–20.0)
Clinical chorioamnionitis (CC)	6.1	0.005	(1.7–22.0)
Fetal bradycardia (FB)	20.9	0.001	(11.3–38.6)
Abruptio placentae (AP)	21.7	0.001	(8.1–57.7)
MSAF	11.3	0.001	(7.3–17.6)
Emergency CS (ECS)	14.1	0.001	(8.6–23.1)
General anaesthesia (Ga)	17.2	0.001	(6.9–42.9)
PROM >18 h	2.10	0.003	(1.2–3.4)

Argentina; ⁷Maternal and Infant, Cleveland Clinic, Cleveland, USA; ⁸Maternal and Infant, Maternidad Martin, Rosario, Argentina; ⁹Maternal and Infant, Maternidad NS de la Merced, Tucuman, Argentina; ¹⁰Maternal and Infant, Hospital Fernandez, Buenos Aires, Argentina

10.1136/archdischild-2014-307384.524

Background Approximately 1% of newborns (NB) require advanced resuscitation (AR) [intubation (ET), and/or chest compression (CC) and/or medication (ME)] at birth. The NRP recommends checking risk factors maternal (MF), intrapartum (IF) and fetal (FF) before each birth (evidence level of expert recommendation) but the need for a team with advanced skills after risk factors have been identified remains undetermined. This imprecision leads to underprovision of expertise which is unsafe or costly overprovision of expertise.

Objective To evaluate the relationship of RF and the need for AR in NB ≥ 34 w gestational age (GA).

Design/methods Prospective, case-controlled study conducted in 16 sites (ARG, CHL, USA and BRA) during 18 months. DR management followed NRP guidelines. Eligible cases were NB ≥ 34 w GA receiving AR at birth and the 4 consecutive NB not requiring AR were selected as controls for the study. Exclusion criteria: prenatal diagnosis of major congenital malformations. Univariate analysis and multivariate logistic regression (MLR) were used to estimate OR and the associated 95% CI.

Results From 61,593 deliveries, 58,429 NB were ≥ 34 w GA (95%). Out of 219 NB receiving AR (0.37%), 23 were excluded, resulting in 196 cases and 784 controls. We found 21 RF

Abstract PS-225 Figure 1