A CONTRIBUTION TO THE STUDY OF THE CAUSATION OF FŒTAL DEATH*

BY

CLARA DUNBAR TINGLE, M.D.,

Demonstrator in Pathology, Sheffield University;

Assistant Pathologist, Sheffield Royal Hospital.

During the last few years, more attention has been paid to what has been, beforetime, a much neglected branch of pathology—the subject of fœtal and neo-natal death. Its importance is obvious. The yearly wastage of fœtal life assumes enormous proportions; nor is this all. The death rate during the first week of life is still very high, and it is to be presumed that some, at any rate, of the factors causing fœtal death may be instrumental in the production of this high death rate, and that a sounder knowledge of fœtal pathology and its treatment would not only reduce the number of still births, but also be associated with a lowered infantile mortality.

Classification of the causes of fœtal death in this series. It is very difficult to classify the causes of fœtal death; indeed the very term 'still birth' and even 'fœtus' are ill-defined in the literature. The word 'fœtus' should not be used indiscriminately, and is not interchangeable with the words 'child' and 'infant.' The term covers a definite period of existence extending from the end of embryonic life to the time when the tissues are aerated by means of its own respiratory efforts. Thus the fœtus has an extra-uterine existence. This is usually very short, but in some instances (*e.g.*, after Cæsarian section) there may be a prolonged interval before respiration is established. It is then obvious that the term 'still birth' must be confusing because it involves more than one entity. The fœtus may die in utero, and later be delivered macerated; it may be killed during labour, or it may be born alive with the heart beating and fœtal movements occurring, and yet its pulmonary respiration may not be established. To each of these cases the term 'still born' applies, yet, as regards their causation and pathology they are widely different. This subject was fully discussed in 1914 by Ballantyne who suggested a solution of the problem by the additional use of the word 'dead birth.' Thus the use of the term 'still birth' would be restricted to the fœtus born alive and dying before the establishment of pulmonary respiration, and the term 'dead birth' reserved for the fœtus born without signs of life.

*The material of this paper is derived from 160 autopsies performed at the Jessop Hospital for Women, Sheffield, on still-born viable fœtuses, or on children dying within seven days of birth, forming a consecutive series, from March, 1923, to March, 1925. My thanks are due to the Honorary and Resident Staff of the Jessop Hospital for their valued help during that period.

Even when definition is made clear, the problem of classification of the causes of foetal death will still be hedged about with difficulties, not only because of great ignorance on such subjects as the transference of disease to the foetus in utero, but also because of the many factors brought into play in any one case. For example, at an autopsy, it may be found that the foetus has extensive intra-cranial hæmorrhage, which is the ultimate cause of death. But we may find that the child was premature and the mother had placenta prævia, for which version was performed. It was presumably during these manipulations that the child was injured. It has been considered in such cases that the actual cause of death was the cerebral hæmorrhage, the predisposing factors being the premature separation of the placenta and the prematurity of the child, which rendered its tissues more likely to damage. In any classification, therefore, there must be considerable overlapping of the various categories, maternal and placental states predisposing in many instances to the foetal condition found at autopsy.

In the present series of cases there were, in all, 160 post mortem examinations made. They form a consecutive series of still-born viable foetuses, or children dying within seven days of birth (Table I.).

TABLE I.

CAUSES OF FŒTAL DEATH IN 160 CONSECUTIVE AUTOPSIES ON STILL-BORN VIABLE FŒTUSES.

MATERNAL STATES.	No. of Cases.
Albuminuria	7
Eclampsia	9
Nephritis and Endocarditis	1
PLACENTAL STATES.	
Placenta prævia	22
Retro-placental hæmatoma	10
Placental infarction	3
FŒTAL STATES.	
Traumatism at birth	105
Foetal infections	14
Foetal anomalies	10

In this classification, there is an overlap of 21 cases where morbid maternal or placental conditions caused or complicated the state of the foetus found at autopsy.

THE STUDY OF THE CAUSATION OF FŒTAL DEATH 257

FŒTAL DEATH DUE TO FŒTAL CONDITIONS.

It is proposed to omit the first two categories, as here set out, and to deal only with the fœtal conditions found. These may be sub-divided into three groups:—

- (I) TRAUMATISM AT BIRTH. (1) Craniotomy. (2) Hæmorrhages of the new-born.
- (II) FŒTAL INFECTIONS.
- (III) FŒTAL ANOMALIES.

I.—TRAUMATISM AT BIRTH.

It is an aphorism that many more fœtuses owe their death to the complications of delivery than to maternal or fœtal disease. Figures here (105 out of 160 or 66%) speak for themselves. It must be remembered, however, that in this classification we are dealing only with the ultimate cause of death, and the predisposing causes (as, for example, placenta prævia, or contracted maternal pelvis) must be brought into consideration.

A second aphorism may be propounded: the fatal results of traumatism are due to hæmorrhage. It is, therefore, with the pathology and ætiology of hæmorrhages of the new-born that we are now concerned.

1. CRANIOTOMY.

I do not propose to deal with these cases in detail. In the majority of cases, the gross trauma has destroyed those parts of the fœtus which are most interesting from a pathological standpoint, namely, the cranium and its delicate contents. In most cases petechial hæmorrhages were met with in other parts of the body. Nothing else of interest has emerged during the examination of the bodies of these fœtuses. Reduction in the number of cases of craniotomy can come by no help from the pathologist, but only from improvement of what Holland calls 'the strategy and tactics of obstetrics.'

2. HÆMORRHAGES OF THE NEW-BORN.

The pioneer work on this subject was published by Spencer⁽¹⁸⁾ in 1892, following the writings of Sarah McNutt⁽¹¹⁾. He gave a detailed account of a consecutive series of 130 autopsies on fresh, mostly still-born fœtuses, and drew some new and remarkable conclusions. Amongst these he stated that visceral hæmorrhages more commonly followed breech delivery than cephalic, and that cerebral hæmorrhage is more frequently found in still-born fœtuses delivered by forceps than in those born naturally by the head or breech. He realised that these hæmorrhages, when not immediately fatal, may be followed by the gravest consequences. He found intra-cranial hæmorrhages in 40·7 per cent. of cases, and noted that, in a certain number of these, bleeding had occurred into the falx and tentorium, but actual tears of the meninges were not mentioned. This omission was due presumably to the method of opening the skull, for it was not till, in 1910, Professor Beneke⁽³⁾ introduced a new method that it was possible to expose the tentorium and falx adequately.

Classification of Hæmorrhages.

In all but 6·9% of cases, hæmorrhage of some degree had occurred. This is an astounding figure until we realise that in 50% hæmorrhage was petechial in type and not *per se* the cause of death:—

TABLE II.
TYPES OF HÆMORRHAGE FOUND.

	Cases.	Per cent.
(i) Petechial hæmorrhage only ...	80	50
(ii) Gross intra-cranial hæmorrhage ...	60	37·5
(iii) Gross visceral hæmorrhage ...	9	5·6
(iv) No hæmorrhage ...	11	6·9
	<hr/> 160 <hr/>	<hr/> 100 <hr/>

(i) Petechial Hæmorrhages.

It is seen that these were almost constant in still-born fœtuses. They varied considerably in number and in size; in some cases they were a most obvious lesion when the body was opened, in others they were found only on careful examination. They were much commoner in the thoracic viscera than in the rest of the body, the usual sites being (a) beneath both visceral and parietal pleura, and (b) beneath the visceral pericardium, especially that covering the upper part of the right ventricle. Frequently also they may be found in the cranium, between the two layers of the falx and of the tentorium. Less commonly, they are also seen in other viscera, as the thymus, liver, and kidneys. In most of the cases, marked congestion of the viscera is associated with them.

These hæmorrhages are found in widely different circumstances, the two cases where the hæmorrhages were most numerous occurring respectively in placenta prævia and prolapse of the cord. The only factor that the cases show in common is the degree of anoxæmia, and these small hæmorrhages may be presumed to correspond to the Tardieu's spots found in true asphyxia.

(ii) Gross Hæmorrhages (Visceral and Intra-Cranial).

Gross hæmorrhages are an extraordinarily common cause of fœtal and neo-natal death. It is probable, too, that the effects do not end here, but are the cause of disabilities which may persist through life.

We have met with gross hæmorrhages 69 times, and of these, 60 showed the results of intra-cranial trauma. The evidence of trauma is found in tears of the meninges, and the devastating effects are due to hæmorrhage. In a certain proportion of these there were tears of the falx cerebri and gross hæmorrhages of the thorax and abdomen, but compared in

frequency with tears of the tentorium, these are rare. Tears of the tentorium were found in 58 cases (exclusive of craniotomies)—a percentage of 36.25 of the total number in the series.

The results of other observers are as follows:—Warwick⁽²¹⁾, in a series of 36 still births and neo-natal deaths, found 50 per cent. showed intra-cranial hæmorrhage. Schafer⁽¹³⁾ found this percentage to be 20 in a series of 680 autopsies upon newly-born infants. Ehrenfest⁽⁷⁾ claims that tentorial ruptures occur in 30 to 40 per cent. of cases of foetal death and deaths occurring soon after delivery. Holland⁽⁹⁾ ⁽¹⁰⁾ found tears of the tentorium to be present in 48 per cent. of fresh still-born fœtuses.

Tearing of the tentorium alone cannot cause death, but in 57 cases out of the 58 autopsies where this type of meningeal tear was found, hæmorrhage had occurred. For purposes of comparison, the amount of blood extravasated was graded empirically as 'little,' 'moderate' or 'much.' Much blood was found in 30 cases, a moderate amount in 19, and little in 8, but it is to be remembered that in many of the cases where the bleeding has been designated as 'little' in quantity, there was accumulation of blood beneath the tentorium, where even a small increase of pressure may be fatal.

It is seen, then, that tears of the tentorium may be regarded as an index of intra-cranial hæmorrhage, and their ætiology must be the same. The same force which tears the meninges produces the hæmorrhage which so often proves fatal. It follows that the anatomy of the dural septa and the part that they play during labour is of outstanding importance.

Anatomy of the Dural Septa.—The tentorium (according to Cunningham) is a crescentic partition of dura mater which forms a membranous covering to the posterior cranial fossa. The cranial dura mater is composed of two layers intimately connected with each other in most regions of the cranium. Along certain lines these layers separate so as to form channels lined with endothelium. These are the blood sinuses. Thus the posterior border of the tentorium from the internal occipital protuberance to the postero-inferior angle of the parietal bone encloses the lateral sinus. Along the superior border of the petrous bone it encloses the superior petrosal sinus. At the junction of the tentorium and falx, the two membranes blend to form a strong band which Holland has called the 'white line.' Along this attachment is enclosed the straight sinus, and at the level of the internal occipital protuberance, where the superior longitudinal sinus ends in joining the lateral sinus, is the torcula herophili, the greatest sinus of all.

The Dural Septa during delivery.—The dural septa are normally in a state of rest, but when labour begins, alterations in their tension are inevitable owing to the nature of their attachment to the cranial bones.

Holland has put forward the theory that the septa should be regarded 'as stays to the cranial bones, like stays to a mast,' exercising a protective function against excessive and dangerous alterations in shape of the foetal head. He believes that the special strengthening bands contained in the septa are specially arranged so as to withstand the stress likely to fall upon cranial contents during moulding of the head. The arrangement of these fibrous bands is interesting; when the foetal head is opened by Beneke's method, they can be seen shining through the dural septa radiating 'like spokes of a wheel.'

Tears of the Dura and their Ætiology.—The site of predilection for dural tears occurs on either side the tentorium just where the fibres of the falx diverge to form its upper blade. The tear may be complete or incomplete,

unilateral or bilateral. I found the falx torn in 12 instances only, always in association with tears of the tentorium.

There are two factors which predispose to tears of the dural septa during delivery: (1) Prematurity; (2) Syphilis. Of the 58 fœtuses exhibiting torn septa, 17 or 29·3% were premature. F. Browne⁽⁴⁾ states that 'The tissues of fœtuses of 7, 7½ and 8 months, are too fragile to pass uninjured through the birth canals,' and agrees that syphilis renders the fœtal tissues more friable.

The exciting cause, however, may be found in cranial stress during delivery, so that the method of delivery in these cases is of importance.

TABLE III.

PRESENTATION IN 58 CASES OF TORN DURAL SEPTA.

HEAD PRESENTATION:						No. of Cases.
(1) Without forceps	...	...	...	...	...	13
(2) With forceps	...	...	...	...	...	17
BREECH PRESENTATION:						
(1) Primary breech	...	...	...	...	...	11
(2) After version	...	...	...	...	...	17

Of the 30 cases of head presentation, it is seen that 13 times forceps were not applied, but only twice can intra-cranial trauma be said to have been caused during normal delivery of a full-term child. An analysis of the 13 cases is given (Table IV.)

TABLE IV.

Prematurity of the fœtus	...	...	...	...	...	4 cases.
Prolonged labour	...	...	...	...	...	2 „
Precipitate labour	...	...	...	...	...	2 „
Contracted pelvis	...	...	...	...	...	3 „
Normal delivery—full-term child	...	...	...	...	...	2 „
—						13 cases.
—						

Forceps were applied 17 times, the indications for their use being as in Table V.

TABLE V.

Contracted pelvis	...	...	...	...	...	5 times.
Placenta Prævia	...	...	...	...	...	2 „
Prolapsed cord	...	...	...	...	...	2 „
Occipito-posterior presentation	...	...	...	...	...	6 „
Primary uterine inertia	...	...	...	...	...	2 „
—						17 times.
—						

Breech deliveries resulted in tearing of the tentorium in 28 cases. In 11 of them the presentation was primarily by the breech, and in 17 breech delivery was the result of version (for placenta prævia 13, for transverse lie 4).

From these figures it is seen that, though tears of the tentorium may occur during the course of a normal vertex delivery, they are uncommon. Prematurity predisposes to the fœtal danger, but the determining causes may, for practical purposes, be divided into (a) excessive stress applied to the fœtal head during delivery (as in cases of contracted pelvis, forceps delivery), and (b) normal stress applied too rapidly (as in precipitate labour).

E. Holland, in considering his series of torn tentoria, came to the conclusion that the particular force required to tear the tentorium was applied in the antero-posterior diameters, *e.g.*, sub-occipito bregmatic. Decrease in this direction necessitates compensatory increase in the vertical height. That part of the cranial vault which becomes elevated and bent, gives attachment to the middle two-thirds of the falx, which is, therefore, pulled upwards by the vault. The falx is attached below to the tentorium, so that this becomes stretched, and if stress is excessive, or suddenly applied, the tentorium tears.

As has been already said, it is not the torn dura mater which is the cause of fœtal death, but the hæmorrhage, which in the majority of cases accompanies the tear, and results from the same stress.

Source of the Hæmorrhage.—In only five cases in this series was it found that the meningeal tear had involved one or other of the sinuses. Some other site and cause of the bleeding must be sought in the majority of cases. In one, the vein of Galen was torn, but in no other case than these six could the source of bleeding be ascertained at autopsy. It was hoped that the exact location of the blood might be helpful in this determination. To this end, following the suggestion made by Holland, ten fœtal heads were injected with strong formalin, and allowed to soak in 10% formalin for 14 days. In five of these, hæmorrhage was found associated with tears of the tentorium. The hæmorrhage in one case was very extensive, but in the remaining four was localised within certain definite limits on the upper surface of the brain between the transverse fissure in front and the torn tentorium behind. The veins in this region are those of the velum interpositum and their tributaries. Each vein of Galen begins at the apex of the velum interpositum, by the union of the vena terminalis with the choroid vein. They terminate by union beneath the splenium of the corpus callosum to form the great vein of Galen, which passes backwards and slightly upwards to end in the anterior extremity of the straight sinus. In addition to the two veins of Galen from which it is formed, it receives tributaries from (1) the gyrus cinguli, (2) the pineal and quadrigeminate bodies, (3) the medial and inferior surfaces of the occipital lobes of the brain,

and (4) the upper surface of the cerebellum. In the foetus, the veins of Galen can readily be distinguished, especially in cases of torn tentorium, where they are found to be engorged with blood; but their smaller tributaries cannot be differentiated. In one case only was there actual rupture of the vein of Galen, therefore in the majority of cases it is the smaller tributary veins which are torn. Holland believes that the commonest source of bleeding in these cases is from the small veins from the mid-brain and cerebellum that enter the vein of Galen, and also the cerebellar veins which enter the straight sinus direct. This would explain why the smaller hæmorrhages that I have seen have been localised to this area.

Mechanism of the Venous Rupture.—The mechanism which leads to rupture of these veins has been fully considered by Holland and may be briefly summarised.

Any movement of the tentorium is necessarily transmitted to the vein of Galen, whose fixed point is its entry into the straight sinus. When the tentorium is unduly stretched, the line of the vein is altered, with the result that kinking occurs at its entrance into the sinus. There is consequently enormous engorgement of the vein and its tributaries, and, if further stretching occurs, they rupture.

Intra-cranial Hæmorrhage without Meningeal Tears.—In two cases only has intra-cranial hæmorrhage been met with where there was no demonstrable tear of the meninges. In both cases, the labour was premature, and the child born alive; in each the intra-cranial findings at autopsy were identical. The left lateral ventricle was filled with blood, and the clot extended through the foramen of Monro into the third ventricle, and from there, through the aqueduct of Sylvius, into the fourth ventricle. Much blood was found beneath the cerebellum and surrounding the medulla, extending along the cervical canal.

The actual source of the bleeding in these cases is a controversial point. Most writers seem to assume that it comes from the choroid plexus, but its profuseness seems rather to point to the rupture of larger veins than those of the plexus.

In these two cases hæmorrhages in other parts of the body were associated with the cranial lesion. They are of interest in regard to the 'Hæmorrhagic Diathesis of the New-Born,' which will be discussed later in the consideration of the general ætiology of hæmorrhages of the new-born.

(iii) *Gross Visceral Hæmorrhages.*

Supra-Renal Hæmorrhage.—Eight cases of hæmorrhage into one or both supra-renal glands have been found; all but three of these were associated with intra-cranial hæmorrhage. Spencer in his monumental work records

THE STUDY OF THE CAUSATION OF FETAL DEATH 263

hæmorrhage into the supra-renals as occurring 26 times, but in some of these cases the hæmorrhage was small in amount, or it is noted as occurring round the organ and not actually into it. Extensive hæmorrhage with rupture of the capsule occurred in three cases of his series, and he gives notes of delivery in two. Both were large fœtuses delivered by the breech, the second after version and through a contracted pelvis. He points out that 'delivery by the lower pole (especially when traction is employed) greatly favours the production of this injury.'

It is interesting to compare Spencer's results with those of more recent work, when it is found that they are strikingly similar. The 34 years of obstetrical technique cannot have influenced to any marked degree the factors which are responsible for the production of these lesions.

TABLE VI.
ANALYSIS OF EIGHT CASES OF SUPRA-RENAL HÆMORRHAGE.

No.	Age of fetus.	Right or left supra-renal.	Associated hæmorrhages or tears.	Mode of Delivery.	Notes.
2	F.T.	L. & R.	Torn Tentorium	Breech	Mother toxæmic
5	F.T.	L. & R.	—	Breech	Lived 3 days
10	8 m.	L. & R.	Torn Tentorium	Breech	—
21	F.T.	L. & R.	Torn Tentorium	Transverse lie, version, breech delivery	—
33	8 m.	R.	Torn Tentorium Ruptured Liver	Placenta prævia, version, breech delivery	—
37	7 m.	R.	—	Vertex presentation; no forceps	Lived 4 days
100	F.T.	L. & R.	Ruptured Liver	Placenta prævia, version, breech delivery	—
126	F.T.	L. & R.	Torn Tentorium	Transverse lie, version, breech delivery	—

It will be seen that the truth of this observation of Spencer's has been fully proved in the present investigation, for, out of the eight cases, seven or 87.5% occurred in breech deliveries. Of these seven, three were in

primary breech deliveries, and four in breech after version. The indication for version is given in Table VI.

Hæmorrhages of the Liver.—Five times have hæmorrhages of large size been met with in the liver, and in four of these the liver has been ruptured, giving rise to excessive intra-peritoneal hæmorrhage. A short analysis of the five cases is given:—

TABLE VII.
ANALYSIS OF FIVE CASES SHOWING HÆMORRHAGE IN LIVER.

No.	Capsule of Liver.	Associated Hæmorrhages and tears.	Method of Delivery.
33	Unruptured	R. Supra-renal Tentorium torn	Placenta prævia, version, breech delivery
34	Ruptured	Tentorium torn	Transverse lie, version, breech delivery
100	Ruptured	R. & L. Supra-renal	Placenta prævia, version, breech delivery
113	Ruptured	Tentorium torn	Transverse lie, version, breech delivery
117	Ruptured	None	Placenta prævia, version, breech delivery

It is very noticeable that in every case where the liver was ruptured, version had been performed. The association of other hæmorrhages in all but one case is also of note. The one case (No. 117) where the only hæmorrhage found had occurred in the liver is of interest. The mother was admitted with accidental hæmorrhage and version was performed. Natural delivery of a breech followed, but the cord was found to be tightly stretched between the legs. The fœtus was still-born, well-developed and obviously at full term. At autopsy the peritoneal cavity contained much blood, and it was found that the bleeding was coming from the under surface of the liver. Here the capsule was ruptured. The tear had occurred at the insertion of the round ligament, and was a slit about $\frac{3}{4}$ in. long, parallel with the hepatic fissure. This appearance suggested the possibility that in this instance, at any rate, the rupture of the liver might have been caused by undue traction on the cord.

Other Visceral Hæmorrhages.—Large hæmorrhages have also been seen into pleural cavities, extensively into the lung substance, into the perirenal

tissue, and into the kidney itself. In two cases hæmorrhages into many organs were met with.

ÆTIOLOGY OF HÆMORRHAGES OF THE NEW-BORN.

Such a record of fatal bleeding occurring during delivery or soon afterwards is of obvious significance, and it is not to be wondered at that during the past few years an extensive literature dealing with its causation has accumulated. Two predisposing ætiological factors and three immediate causes of hæmorrhage emerge from a mass of speculation.

Predisposing Factors.—These consist of (a) Prematurity, and (b) Syphilis, and of them little need be said. What has been stated of the tissues generally applies particularly to the vessels; imperfectly formed structures are more liable to damage than those whose development is complete.

It may be suggested here that syphilis does not necessarily act in damaging the fœtal tissues, but possibly by its effect in causing prematurity.

Immediate Causes.—Here have to be considered (a) Congestion, (b) Hæmorrhagic Diathesis of the New-Born, and (c) Trauma.

(a) *Congestion.*—The influence of asphyxial congestion as a predisposing cause of hæmorrhage was first pointed out by Ashley in 1890; Cruikshank⁽⁶⁾, however, goes further and states that congestion *per se* is the essential element. He says that 'with regard to the ætiology of hæmorrhage the factor of greatest importance appears to be the passive congestion produced by pre-natal and intra-natal asphyxia.' The degree of congestion depends chiefly on the length of time which elapses between the interruption of the exchange of gases in the placenta and the establishment of pulmonary respiration. The next most important element he believes to be the increase in congestion brought about by the passage through the birth canal; while injuries due to abnormalities of presentation or to operative interference occupy the third place in the ætiology of these lesions.

(b) *Hæmorrhagic Diathesis of the New-Born.*—Townsend in 1891 read a paper before the Boston Medical Society in which the term 'hæmorrhagic disease of the new-born' was applied to a condition in which effusions of blood occur, apparently spontaneously, in various parts of the body. He was inclined to favour the view that these were in their origin infective. He was probably the first to make a clear differentiation between such spontaneous hæmorrhages and those due to trauma.

The importance of the hæmorrhagic diathesis was not realised until Robert M. Green suggested in 1914 that birth hæmorrhages might be solely the local manifestation of a hæmorrhagic tendency. Warwick, Munro and Frostis, and many others emphasised the importance of this contention. Rodda's work on the blood-coagulation time in the new-born is of great

value in this connection. The normal coagulation time he found ranged at birth from five to nine minutes. On the second day of life the blood-coagulation time was slightly increased, and the time lengthened until it reached its maximum on the fifth day. Then a return to the average of the first day was reached by the tenth day. Evidence of hæmorrhage appeared to be a prolonged bleeding time with a delayed coagulation time. He concluded that there were factors other than trauma concerned with cerebral hæmorrhage, and that the obstetrician was carrying burdens unduly laid on him. Other observers, grouping these two factors of congestion and foetal disease together, have come to the conclusion that the number of cases of hæmorrhage which might properly be classed as traumatic in origin is quite small.

(c) *Trauma*.—Spencer, in his pioneer work in 1891, laid stress on the mechanical traumatic factor in the ætiology of hæmorrhage, and the exact mechanism of intra-cranial trauma was definitely established by Beneke. It has been fully confirmed by many other investigators, especially by Holland, Capon and Ehrenfest.

Ætiological Factors in the Present Series.

From a consideration of this series of 160 cases, the dominance of the traumatic factor is made evident. In the great majority of cases of intra-cranial hæmorrhage, other signs of injury are manifest, such, for example, as tearing of the meninges, or in rarer cases fracture of the skull.

It is certain that venous congestion must be taken into account, but I believe as a predisposing, not as an immediate, cause of hæmorrhage. The wall of a distended vein will rupture much more easily than the wall of one less completely filled with blood, and the resulting hæmorrhage will be much greater in amount in the former case.

Anoxæmia may cause capillary oozing and petechial hæmorrhages, but trauma is responsible for those gross hæmorrhages which either kill the foetus or cripple the child.

Trauma producing cerebral hæmorrhage is the root cause of many deaths clinically ascribed to asphyxia. The asphyxia is really due to mechanical pressure on the respiratory centre in the medulla by extravasated blood.

With regard to the 'Hæmorrhagic Disease of the New-Born,' it seemed to me, early in the work, that in certain instances, the traumatic element was insufficient of itself to cause the enormous amount of hæmorrhage found at autopsy. With Rodda's work in view, I undertook a series of investigations as to the blood-clotting time in these children and their mothers, using the apparatus consisting essentially of a small glass tube containing a lead shot. One interesting fact became obvious at once—that the blood-coagulation time of the child on the first day is that of the mother. After

THE STUDY OF THE CAUSATION OF FŒTAL DEATH 267

the first day, the time was found to be a little prolonged until the end of the first week, returning to the normal at the end of the second week. In only four cases was there noticeable delay in the clotting time; these cases are, however, extremely interesting in that they are of the type in which a mild degree of trauma gave rise to severe hæmorrhage. Two of these cases have been described above under the heading, 'Intra-cranial hæmorrhages apart from tears of the meninges.'

The two other cases in which delay in the clotting time was noted were both full-term children, born alive. Each lived for several days, and it was possible to record the alterations in the coagulation time. At autopsy numerous large hæmorrhages were found as follows:—In one, hæmorrhage into the skin, sub-peritoneal tissue, under the capsule of the liver, and into the peri-nephric tissues. In the other, extensive hæmorrhage into both lungs, sub-capsular hæmorrhage in the liver, and hæmorrhages between the layers of the falx and tentorium.

TABLE VIII.

A RECORD OF THE BLOOD-COAGULATION TIME IN FOUR CASES.

No.	Hæmorrhage.	Blood-clotting time.			Age at death.
		1st day.	3rd day.	7th day.	
93	Multiple	11 mins.	13 mins.	—	3 days
134	Intra-ventricular	22 mins.	—	—	56 hours
141	Multiple	9 mins.	11 mins.	13 mins.	7 days
150	Intra-ventricular	29 mins.	—	—	12 hours

For normal infants, it was found that the blood-clotting time on the first day was less than five minutes, and this time was not exceeded in the great majority of the children in which gross hæmorrhages at autopsy were found. There appears to be, however, a small group of cases in which a hæmorrhagic tendency exists. Evidently, in these instances even a mild degree of trauma may lead to severe and progressive hæmorrhage.

There must be no exaggeration of this factor. Stress should rather be laid on the importance of trauma during delivery as an ætiological factor in the causation of fœtal death. Consequently it may be hoped that the improvement of obstetrical technique will avert many of these fœtal tragedies of delivery.

Conclusions on the Ætiology of Hæmorrhages of the New-Born.

The following classification of ætiological factors is suggested:—

PREDISPOSING FACTORS:

- (1) Prematurity (with which *syphilis* is possibly included).
- (2) Venous Congestion.
- (3) Hæmorrhagic Disease of the New-Born.

IMMEDIATE FACTOR:

Mechanical Trauma.

II.—FŒTAL INFECTIONS.

Under this heading are included two different types of cases: one in which the cause of death is a true fœtal infection, and one in which the infective agent attacks the new-born child. In the latter class the disease does not differ much from that occurring in later infancy, and I do not propose to discuss these cases.

An infective process which attacks a fœtus must be modified by the intra-uterine environment. The disease may begin early in the course of fœtal development, and the fœtus, protected as it is by the amniotic fluid, and provided with an adequate blood supply, yet may be able to live. But, after birth, when the maternal blood supply is cut off and the protection is removed, the condition may prove fatal to the child. It seems possible, too, that there are infections which gain access to the fœtus during its passage through the birth canal. Of this group pneumonia forms an example.

PNEUMONIA.

F. Browne^(4a) has said that pneumonia is the commonest cause of neo-natal death. In my experience this is true only after the exclusion of the complications of delivery, which, though they may be insufficient to cause still-birth, render the child unfitted to meet the stress and strain of neo-natal life.

I have seen nine cases of pneumonia; in seven the child was born alive and died of pneumonia before the end of the first week. In this series of 160, 43 were autopsies performed on children who had lived, so that the percentage of deaths from pneumonia is 16·3. In each case the infection has run a very rapid course, and the symptoms have been so few that only once was pneumonia suspected, the diagnosis being based on the cyanosis of the child and its rapid and wheezing respirations. The histories have, in all except one case which I believe has a different ætiology, one point of similarity, namely, prolongation of the time between the rupture of the membranes and the birth of the child.

Sections of the lung tissue have been made in every instance, and in all except one, the condition has been that of broncho-pneumonia of catarrhal-celled type. The sections show a patchy consolidation, the

SAGITTAL SECTION OF FETAL HEAD.

Case No. 108.

Showing sub-tentorial hæmorrhage.
Note the shape of the head after
moulding.

FETAL HEAD.

Case No. 126.

Opened after the Beneke method, showing a tear of the right side of the tentorium, and sub-tentorial hæmorrhage.

THE STUDY OF THE CAUSATION OF FŒTAL DEATH 269

consolidated patches being confined to the areas surrounding bronchi. In these regions, the alveoli contain many large catarrhal cells, a few

Case No. 8.

Micro-photographs of section of the Lung.

Low power.

High power.

Showing consolidation of the lung tissue with accumulation of catarrhal cells polymorphonuclear leucocytes, a few red blood cells and a fluid containing fibrin. The vessels are much congested, and in most cases the areas surrounding the pneumonic patches show emphysema.

Cultures were made from the lung substance in every instance; three times a pure growth of *B. Coli Communior* was obtained, once a pure growth of *B. Coli Communis*; three times coliform bacilli were found, but the cultures were contaminated by diphtheroid and other organisms. In one, cultures from the lung grew a staphylococcus and a long chained diplo-streptococcus, and in another a pure growth of pneumococci was obtained. It has been put forward that in these cases of pneumonia, especially where the causal organism is of the coliform group, the bacillus enters the ruptured amniotic sac from the vagina of the mother and so reaches the nose and mouth of the fœtus. This being so, a delayed period between the rupture of the membranes and the birth of the child would be expected as an ætiological factor. It has been shown that in this series the only factor that could be found in common to all cases, with the one exception mentioned, was this delay in the second stage of labour. It is of interest in this connection to note that Browne in 1922 recorded a case where cultures were made from the throat of an apparently healthy infant immediately after birth in which the membranes had been ruptured 60 hours before delivery. Here cultures yielded a growth of *Bacillus Coli* and diphtheroid bacilli.

In some of the earlier cases of the series, it seemed to be that the pneumonic condition must have been existing, and even well advanced before birth, for, in the body of one child who lived ten hours, extensive hepatisation and fibrinous pleurisy were present. Two cases occurring later in the series, have made it possible to place the contention beyond doubt.

One of these was that of a full-term female foetus, whose mother was admitted with a right occipito-posterior presentation. Manual rotation was effected, and the foetus was extracted with difficulty by means of forceps. The membranes had been ruptured about 48 hours before delivery. At the autopsy on the child, extensive patches of pneumonia in the stage of grey hepatisation were found in both lungs. There was no pleurisy. Cultures from the lung surface gave a pure growth of *B. Coli Communis*.

The other was also a full-term female foetus delivered by craniotomy after unsuccessful applications of forceps. Here delivery was effected more than 60 hours after the rupture of the membranes. The lungs of the foetus showed a patchy consolidation with broncho-pneumonia of catarrhal-celled type. Cultures from the lung yielded a pure growth of *B. Coli Communis*.

It is evident from these cases and from similar cases recorded, that broncho-pneumonia may be present and even well advanced before birth.

A further case already mentioned is of considerable interest from the point of view of the possibility of the transference of disease from the mother to the foetus. It has been said already that this case cannot be brought into line with the other cases of pneumonia, either with regard to its ætiology or to its bacteriology.

A full-term female child was delivered by Cæsarian section, the membranes being unruptured before the operation, which was performed on March 24th, 1924. The child was born alive, but had a cyanotic appearance from birth and never attained a healthy colour. Its breathing was shallow and more rapid than normal, and it died on April 1st, 1924. The mother died on the 26th March, 1924, and autopsies were performed on both. The mother had an acute bronchitis with thick purulent exudate into the bronchi, and areas of broncho-pneumonia in both lungs. Sections from the lung showed areas of consolidation surrounding the bronchioles, with much polymorphonuclear emigration into the alveoli where also a few red blood cells and entangled threads of fibrin were found. The vessels in the vicinity showed a moderate degree of acute congestion. The bronchi were seen in sections to be the subject of an acute purulent inflammation, the changes being more marked in the smaller branches of the bronchial tree.

Autopsy on the child revealed a similar condition of pneumonia. The whole of the base of each lung was consolidated and in the stage of grey hepatisation. Sections confirmed the diagnosis of pneumonia, which was certainly broncho-pneumonia by its distribution, but, unlike the other cases of neo-natal pneumonia, was not catarrhal-celled in type. The predominating cell, instead of the catarrhal cell of the other pneumonias in the series, was the polymorphonuclear leucocyte, which was present in great numbers in the affected alveoli and in the bronchioles in the centre of the consolidated areas. The vessels showed much congestion and polymorphonuclear leucocytic migration into the surroundings. Instead of the coliform bacilli, which were the predominating organisms in the other cases, cultures from the lung yielded a pure growth of pneumococci.

It seems feasible that in this case infection was conveyed from the mother to the foetus, either by means of the placental circulation or indirectly. Direct transmission to the child is unlikely, as, though the child lived seven days, the mother lived for

THE STUDY OF THE CAUSATION OF FŒTAL DEATH 271

two days only after delivery, and during this time there was no direct communication. It must be remembered, too, that from its birth, the child showed symptoms of pulmonary embarrassment.

One case of pneumonia is of interest from a medico-legal aspect. The child was three days old and appeared quite healthy till two hours before its death, which was dramatically sudden, and might, in other circumstances, have given rise to the suspicion of foul play, or of overlying. At the autopsy, small areas of broncho-pneumonia were found. The outstanding feature, both macroscopically and microscopically, was the amount of hæmorrhage which had occurred into the lung substance. Cultures from the lung were sterile. It has been said that the ætiological feature in such cases as this is usually organismal, but Browne states that 'there is evidence to support the theory that it may occasionally be of the nature of an anaphylaxis.'

III.—FŒTAL ANOMALIES.

This heading includes the cases of mal-development, and the morbid states which cannot, at present, be assigned to any known infective agent.

A list of those exemplified in the series has been given, and some are singled out for further reference.

GENERAL ŒDEMA OF THE FŒTUS.

This is a rare condition of which I have seen one case. Ballantyne⁽¹⁾ ⁽²⁾ defines the condition as 'a rare condition of the fœtus, characterised by general

Case No. 136.

Micro-photographs of sections of the Liver.

Low power.

High power.

Showing the aggregation of hæmo-poietic cells, the separation of liver cells, and their pigmentation.

anasarca, by the presence of fluid effusions in the peritoneal, pleural and pericardial sacs, and usually by œdema of the placenta; and resulting in the death of the fœtus or infant before, during, or very soon after birth.'

It will be seen that the case to be described fulfils the conditions necessary to be placed in this category.

CASE.—The mother was 36 years old, tall and well built. She had had six previous children, all being normal deliveries of full-term healthy infants born alive, and during the pregnancies she had been well. The œdematous foetus was born at 7½ months. During the last month of the pregnancy the mother noticed coloured spots before her eyes at intervals, and she also had vague abdominal pains. Six days before delivery her feet and ankles became so swollen that she could not get her shoes on. Before delivery there was extensive hæmorrhage, the estimated loss being about one pint.

A catheter specimen of urine was obtained on the day of delivery. The specific gravity was found to be 1010 and the reaction acid. A heavy cloud of albumen was present, but chemical tests for other abnormal constituents were negative. On microscopical examination of the centrifuged deposit, a few pus cells were seen but no red blood cells and no casts or crystals. Stained films showed an absence of organisms.

Clinically, the mother showed a severe anæmia due to the blood loss, but no other lesion was demonstrable.

Under general anæsthesia, the membranes were ruptured and a leg brought down, and later a still-born foetus was delivered by the breech.

The foetus was a male, obviously premature, its age corresponding apparently with the estimated 7½ months gestation. The weight was 1680 gms. and the length 15½ ins. The abdomen was protuberant, and there was generalised œdema, especially of the face, chest, abdomen and external genitals. There was a clean cut wound of the skin and subcutaneous tissues along the fold of the left groin due presumably to forcible extension of the thigh during delivery. Clear yellow fluid flowed from any incision made into the subcutaneous tissues.

Post mortem examination showed small hæmorrhages between the scalp and the skull. There were no meningeal tears, but small hæmorrhages were present between the two layers of the falx cerebri and between the two layers of the tentorium cerebelli. No intra-cerebral hæmorrhage was found. The brain was œdematous and the ventricles were slightly distended with clear yellow fluid. The pleural cavities each contained about 10 ccs. of similar clear yellow fluid; the lungs had not been ærated but were apparently normal, the thymus was not enlarged, but was softer and more friable than usual; the thyroid and lymph glands were not enlarged. The pericardial sac contained about 10 ccs. of clear yellow fluid; the heart showed no congenital or other abnormalities. The peritoneal cavity contained about 200 ccs. of clear yellow fluid, and there was a little fibrinous deposit on the surface of the viscera in places, and on the peritoneal aspect of the abdominal wall. The liver was normal in appearance and consistency, and weighed 150 gms.; the umbilical vein and ductus venosus were patent and the gall bladder was distended with bile. The spleen weighed 40 gms., was deep red in colour and firm in consistency; pancreas, mesentery, stomach and intestines appeared healthy. The supra-renals showed many sub-capsular hæmorrhages, but otherwise appeared normal. Each kidney weighed 22 gms. and both appeared normal; there was no evidence of foetal nephritis, macroscopical or microscopical. The ureters and urethra were proved to be patent, and the bladder contained a little straw-coloured urine. There was much œdema of the scrotum. The left testis was still in the abdominal cavity at the level of the brim of the pelvis, the right testis being in the scrotum.

The placenta weighed 1210 gms., was bulky and very friable, dense white in colour. Much blood clot was present on the maternal side. The umbilical cord was 19 inches in length, ½ inch in diameter, and Wharton's jelly appeared very translucent. A drop of foetal urine was examined for albumen, and a faint cloud was found to be present.

THE STUDY OF THE CAUSATION OF FŒTAL DEATH 273

The ætiological factors of general œdema on which stress has been laid in the literature may be classified as (i) maternal, (ii) paternal and (iii) fœtal.

(i) Four maternal causes have been suggested.

(a) Ballantyne laid emphasis on the advanced age of the mother in these cases of fœtal œdema. This was in 1902, and in Schumann's⁽¹⁴⁾ list compiled in 1915, only 13·3% of cases occurred in mothers over 35 years of age. (In the present case the mother was 36.)

(b) In the great majority of the recorded cases there is some evidence of toxæmia in the mother, and this usually takes the form of albuminuria, œdema or excessive vomiting.

In the case under consideration now, the urine of the mother, on admission, contained a dense cloud of albumen; no blood, and casts were not found. There was marked œdema of the lower extremities and abdominal wall. She complained of 'coloured spots' before her eyes; there was no history of excessive vomiting, of headache, or of fits. She was admitted to hospital with accidental hæmorrhage.

(c) Spirochætes have once been found in the tissues of an œdematous fœtus, and in a few instances the maternal blood has given a positive Wassermann reaction, but in the great majority of cases, there has been no evidence of syphilis either clinically in the mother or found at autopsy in the child. In our case there was no evidence of syphilis, and the maternal Wassermann reaction was negative.

(d) Ballantyne first drew attention to the fact that the mother of an œdematous fœtus is often anæmic. The mother, in our case, was very pale owing to loss of blood, but there was no previous history of anæmia.

In weighing the importance of maternal morbid states as a causative factor, it must be remembered that while such maternal conditions are, unfortunately, all too common, fœtal œdema is, as Ballantyne states, a rare condition. It is reasonable to assume that there is some other factor as yet unknown. This may be directly causative or acting by means of the maternal condition.

(ii) Some writers have suggested various paternal disorders as a possible causation, *e.g.*, œdema in the father, lead poisoning, etc. The father in our case was not examined except during the taking of blood for a Wassermann reaction, no gross lesions were obvious; his size and physique were normal; there was no history of working in lead and his Wassermann reaction was negative. It seems that little reliable evidence is forthcoming on the importance of such paternal disease in fœtal morbidity.

(iii) In certain cases described, actual mechanical obstruction to the vascular or lymphatic system of the fœtus has been found. Such structural defects as, for example, complete absence of the lymphatic system as recorded by Smith and Birningham⁽¹⁷⁾ in 1899, might be reasonably brought

forward as the cause of generalised œdema, but in some cases it is difficult to ascertain the actual way in which the lesion produced the œdematous state. It has been said that in all cases, a mechanical cause can be found if a thorough post mortem examination is made, but this contention has not been supported by later experience. In a certain number of cases, congenital abnormalities have been found associated with the generalised œdema, but congenital abnormalities occur by no means infrequently in any series of stillbirths (*e.g.*, 9% of the present 160 cases), and it is possible that the associated conditions may be due to one and the same cause.

I examined the œdematous fœtus as carefully as was possible with a view to discovering any manifest mechanical cause for the œdema, but I was unable to demonstrate any obstructive lesion, either cardiac as recorded, *e.g.*, by Tait⁽⁴⁹⁾ in 1895, or umbilical (*see* Crozier and Fischer).

The case under consideration, then, comes under the heading of non-obstructive causation, and many other examples are to be found in the writings.

Schridde⁽⁴⁵⁾ in 1910 first drew attention to certain histological features, and the cases reported since have shown similar pathological findings.

It would, at this stage, be as well to describe the histological appearances as I saw them. The most active changes were found in the liver. Here normoblasts, megaloblasts and myeloid cells were seen in enormous aggregations so that groups of liver cells were cut off from one another by the hæmopoietic cells. The latter were showing well-marked karyokinetic figures. This accumulation of cells did not affect any particular part of the liver tissue more than another nor was there any demonstrable increase in the connective tissue of the liver. This does not agree with some of the described cases where it is stated that the periportal connective tissue is increased in quantity and that the hæmopoietic cells tend to accumulate in the region of the vessels within this periportal tissue. The liver cells themselves contained a golden brown pigment which, by means of potassium ferrocyanide and hydrochloric acid was shown to contain iron. In the spleen, a similar change was found, but not so advanced in degree. An accumulation of myeloblasts, megaloblasts and normoblasts had to a certain extent replaced the normal spleen tissue. Again karyokinetic figures were evident, and in the lymphoid cells of the spleen a small amount of an iron-containing pigment was found. The kidneys showed a similar accumulation of cells, but in much less degree than in liver and spleen. There was no evidence of fœtal nephritis, and no pigment was demonstrated. Search for the *spirochæta pallida* proved futile.

In assessing the value of this evidence, I have borne in mind the fact that we were dealing with a premature fœtus, and that during later fœtal life, the liver and spleen, along with the bone marrow, are the sources of the solid elements of the blood. To this end, we have sectioned the livers of premature children, normal, and abnormal of approximately the same age as the œdematous fœtus. Here we have found small clusters of nucleated red blood cells, normoblasts with a few megaloblasts always placed in definite situations between the columns of liver cells. The livers of syphilitic infants are said to contain many of these cell clusters; this is assumed to be a response to the destruction of red cells brought about by

THE STUDY OF THE CAUSATION OF FETAL DEATH 275

the circulating toxin. In the liver of an œdematous foetus, however, it is such a gross exaggeration of these processes as to show a grade of hæmopoietic response not met with in other conditions. The fact, too, that a similar response is met with, not only in the liver, but also in the spleen, and probably, though to a less extent, in the kidney, seems to show that all possible sources of blood formation are being tapped. The presence of an iron-containing pigment in the cells of the liver and spleen, points to a co-existing destruction of red blood cells.

The view appears justified that some unknown toxin is being elaborated, destroying the red blood cells and calling into being all the potential sources of blood-cell formation.

The question of the origin of the toxin of foetal, placental, and maternal elaboration, cannot in our present state of knowledge be decided.

CONGENITAL ABNORMALITIES OF THE HEART.

Four cases of congenital heart lesions occur in the series, and I am including a fifth example which I obtained at the Royal Hospital, Sheffield, from a child who died there at the age of five weeks.

Three of the five are of the type described as pulmonary stenosis. The hearts showed gross enlargement especially of the right side. The pulmonary artery was in each case small in calibre and there was varying degree of stenosis at the pulmonary orifice, though there was no evidence of fusion of the cusps. In no case was the stenosis complete, but the fifth case is of interest because the child lived for five weeks and yet the stenosis was so extreme that it is questionable if more than minute quantities of blood had been received by the lungs through the contracted orifice.

These three cases show the effects of arrest of developmental expansion of the infundibular cavity first described by Keith. There is frequently an associated defect in the interventricular septum, and this was found in two out of the three cases here described, but the foramen ovale and ductus arteriosus did not differ in any particular from those of a normal full-term still-born foetus, and the auriculo-ventricular valves and the aorta with its valves showed no lesions.

In these cases of severe pulmonary stenosis, it is difficult to understand how the pulmonary circulation is carried on especially after birth. This difficulty suggested itself to Keith, who injected the arteries in one of these cases and found that the bronchial arteries and other accessory branches derived from the intercostals were enormously enlarged in size and increased in number, and it is to be presumed that the arterial system of the lungs is filled from the systemic-circulation.

No. 4 of the series of congenital heart lesions showed a less common condition, namely, almost complete atrophy of the body of the right ventricle, with absence of the tricuspid valve. There was a large central

interventricular foramen. The foramen ovale and ductus arteriosus were patent, the fœtus being still-born. It is from cases similar to this that Keith illustrates what he calls 'the dual constitution of the right ventricle,' for here there is no stenosis of the pulmonary valve and the infundibular part is expanded, while there has been arrest of the development of the body of the right ventricle.

No. 5 of this series showed the opposite condition, namely, considerable atrophy of the left ventricle. Here, the right side of the heart appeared a little larger than usual, but the left ventricle was so small that the cavity did not admit the tip of the little finger. The mitral valve was not demonstrable, this corresponding with the absence of the tricuspid valve in No. 4. In these cases the obliteration of the valve is probably due to the fusion of the endocardial tissue out of which the valve cushions are fashioned. Here the aorta was very narrow and the ductus arteriosus was not abnormally patent and yet the child lived three days. The foramen ovale was patent, and there was a small central interventricular foramen. This case was of interest, too, because small recent and friable vegetations were found on the tricuspid and pulmonary valves, there being no evidence of stenosis of these valves. Microscope sections showed the presence of masses of fibrin with entangled red blood cells and leucocytes. No bacteria could be found.

In this instance it is obvious that the endocarditis evinced at autopsy was not the cause of the gross heart lesions found. It is possible that the same process operating in very early fœtal life and causing arrested development had, acting in the later months of intra-uterine life, given rise to a definite acute endocarditis, but it must be remembered that a damaged structure is more prone to bacterial damage than one normal and healthy.

It will be seen that in every instance but one there was defect in the interventricular septum. This is very commonly found in association with other malformations especially with pulmonary stenosis and occasionally is met with alone. It was suggested by Meckel that the interventricular foramen might be the primary lesion, the pulmonary stenosis occurring subsequently. The exact sequence of events which lead to the closure of the interventricular septum is not known, but from the fact that the foramen disappears as soon as the infundibulum becomes incorporated with the right ventricle, it would seem that the expansion of the cavity of the bulbus cordis must play an important part in its closure. Thus the work of Keith points to the view that the failure to close of the interventricular foramen is a condition secondary to other developmental errors.

The association of cardiac anomalies with other instances of faulty development is a well-established fact, for example, Vierordt found that associated defects occurred 80 times in 700 cases, while Abbott found it in

THE STUDY OF THE CAUSATION OF FŒTAL DEATH 277

20% of cases. Of the five cases recorded here, two also showed absence of the right kidney and ureter, and one of these two was, in addition, hydrocephalic.

Garrod has pointed out the association of Mongolian deficiency and congenital cardiac lesions. In two of our five cases there were definite signs of Mongolism. That is to say that in one case only was there no other defect found in association with the mal-development of the heart.

ACHONDROPLASIA.

In this series is one case of Achondroplasia.

The family history was enquired into with care, but no evidence of antecedent dwarfism was obtained. The mother was intelligent and of good education. She was 32 years of age when the achondroplasiac child was born. This was her second child, the first having been born four years before and being alive and apparently normal in all respects. The achondroplasiac child was born at full term, a normal breech delivery. The only abnormality found in the mother was an asymmetrical pelvis due to a fracture of the neck of the femur which had caused much shortening. Her Wassermann reaction was negative.

In this case at first glance it seemed apparent that the child's head was too large, while its limbs were too short. On measurement, however, it was found that only part of these assumptions could be substantiated. The size of the head was not increased; the circumference being that of a normal child of the same development. The limbs were found to be very short. When vertical, the arms reached only to a point midway between the crest of the ilium and the great trochanter. The lower limbs were only about half the length of those of a normal child. The chief diminution in length, as in all cases of achondroplasia, was in the humerus and femur respectively. Taking the length of the humerus in the normal infant as 6 cms. both were excessively shortened, the right humerus measured 3.2 cms. and the left 3.3 cms., the radius and ulna were of equal length, each measuring 2.9 cs. The femora were of equal length each measuring 6 cms., and there was on each side a slight concavity inwards of the shaft. The fibulæ measured 4.4 cms. each, and the contour was normal. The hands showed to a moderate degree that anomaly first pointed out by J. Thomson. The palm is flat, and the fingers, instead of being parallel, diverge, two to the ulnar and two to the radial side, forming the so-called 'trident' hand. The ribs were very short (the maximal length being three-quarters of an inch) while the costal cartilages were long. A deep depression was seen at the junction of the ribs with the costal cartilages. The trunk was of normal size and shape.

Morbid Anatomy.—Microscopic sections were made through the lower end of the right femur, but nothing was seen except an alteration in the normal parallel arrangement of the cartilage cells and a comparative dearth of vessels. In some cases described the epiphysis has been separated from the diaphysis by an ingrowth of connective tissue derived from the periosteum.

The internal organs showed no pathological change, and no other congenital abnormalities found, though these were sought, remembering that such had been recorded. For example, Nathan⁽¹²⁾ described congenital hernia in association with achondroplasia, and Kassowitz congenital dislocation of the hip.

The thyroid was sectioned in view of the case described by Symington & Thomson, but no lesion found. The pituitary gland, macroscopically appearing normal was, unfortunately, not examined microscopically.

CONGENITAL GOITRE.

One case of congenital enlargement of the thyroid has been met with.

This was in a still-born $7\frac{1}{2}$ months foetus, that presented by the breech. The mother had no swelling of the thyroid, and had had two children previously born alive and normal in all respects. In this foetus there were many associated abnormalities. Spina bifida was present in the lumbo-sacral region with meningocele, and there was a large umbilical hernia which contained not only many coils of small intestine, but also the gall bladder and a tongue of liver substance derived from the right lobe.

The thyroid itself formed a large mass in the front of the neck occupying the whole region between the chin and the manubrium sterni, the tissues being oedematous over it. The gland, when dissected free, was found to be $3\frac{1}{2}$ inches long, about 2 inches across, and, in places, over 1 inch in thickness. The whole of the gland was affected and it surrounded the trachea almost entirely.

Sections showed increase in the number of acini, and hypertrophy of the lining cells. No attempt to form colloid had yet been made.

CONCLUSIONS.

In this study of the causation of foetal and neo-natal death in a series of 160 cases the following conclusions are reached:—

(1) More foetuses die as the result of excessive trauma during delivery than from maternal or foetal disease.

(2) The fatal results of excessive trauma are due to hæmorrhage.

(3) Certain maternal morbid states predispose to foetal injury during delivery.

(4) Factors predisposing to hæmorrhage are prematurity, excessive venous congestion, and the ' hæmorrhagic diathesis.'

(5) Pneumonia is the commonest infection causing neo-natal death.

(6) The term ' congenital heart disease ' is frequently a misnomer.

REFERENCES.

1. Ballantyne, J. W., *The Diseases and Deformities of the Fœtus*, Edin., 1892.
2. Ballantyne, J. W., *Ante-natal Pathology and Hygiene*, Edin., 1902.
3. Beneke, *Munch. Med. Woch.*, 1910, LVII, 41.
4. Browne, F. J., *Edin. Med. Journ.*, 1921, N.S. XXXVII, 590; and 4A, *Lancet*, 1922, II, 592.
5. Capon, N. B., *Journ. Obstet. and Gyn.*, 1922, XXIX, 239.
6. Cruikshank, J. N., *Lancet*, 1923, I, 836.
7. Ehrenfest, H. E., *Birth Injuries of the Child*, N. Y., 1922.
8. Graham, E. A., *Journ. Exper. Med.*, XV, 307.
9. Holland, E., Report on the Causation of Foetal Death, Ministry of Health Report, 1922.
10. Holland, E., *Journ. Obstet. and Gyn.*, Brit. Emp., Winter, 1922.
11. McNutt, S., *Amer. Journ. Obstet.*, N. Y., 1885.
12. Nathan, *Amer. Journ. Med. Sci.*, 1904, CXXVII, 690.
13. Schafer, *Zeitschr. fur Geburtsch. u. Gynakol*, 1921, LXXXIV, 239.
14. Schumann, F. A., *Amer. Journ. Obstet.*, 1915, LXXII, 961.
15. Schridde, H., *Munch. Med. Woch.*, 1910, LVII, 397.
16. Schridde, H., *Deutsh. Med. Woch.*, 1911, XXXVII, 432.
17. Smith, A. J., and Birmingham, F., *Journ. Anat. and Physiol.*, 1889, XXIII, 532.
18. Spencer, Herbert R., *Trans. Obstet. Soc., London*, 1891, XXVIII, 203.
19. Tait, L., *Trans. Obstet. Soc., London*, 1895, XXXVII, 5.
20. Thomson, J., *Edin. Med. Journ.*, 1893, XXVII, 1112.
21. Warwick, M., *Amer. Journ. Med. Sci.*, 1919, CLVIII, 95.