Millennium Development Goals: progress in Oceania

John D Vince

Correspondence to

Dr John D Vince, Clinical Sciences Division, School of Medicine and Health Sciences, University of Papua New Guinea, PO Box 5623, Boroko, National Capital District, Port Moresby 111, Papua New Guinea; johndvince@gmail.com

Received 16 April 2014

Revised 20 May 2014

Accepted 24 May 2014

For the purposes of this commentary, Oceania is understood to encompass the Pacific Island states of Micronesia, Polynesia and Melanesia including Papua New Guinea (PNG) (but excluding the western half of mainland New Guinea, politically a province of Indonesia), Timor Leste, Australia and New Zealand. In geopolitical Oceania (which excludes Timor Leste), 25 countries, 12 with a population <100 000, are scattered over an area of 8 525 989 km² (figure 1). The total population is 35 670 000, 27.5 million of whom are in Australia and New Zealand. PNG accounts for the large majority of the remainder. The great majority of the population in Australia and New Zealand are urban dwellers, while the majority of the population in the other countries is rural, the people often living in villages with very difficult access on small islands or atolls or in remote mountain ranges.

The region includes countries with widely differing economies, infrastructure development, education, health service provision, and health indices. Table 1, based on data from Unicef's State of the World's Children and the United Nations Development Programme reports, indicates some of the important indices relating to child health from a selection of the countries.^{1 2} Only one country, Timor Leste, had reached the Millennium Development Goal (MDG)-4 target by 2012, but from a very high under 5 mortality rate in 1990. Australia, New Zealand and Vanuatu may reach the target, but most of the countries are unlikely to do so.

Mortality rates and economic indicators in Australia and New Zealand are similar to those throughout the rest of the Organization for Economic Cooperation and Development (OECD) countries. However, the overall figures mask, as they do in most countries, mortality rates for their disadvantaged population: in the case of Australia, the Aboriginal and Torres Strait Island population, which are more than double the national figures. Nevertheless, mortality rates in this population have been reduced by almost 50% since 2002.³

Good child health is dependent on many inter-related factors, the most important of which are living conditions, the level of female literacy, and financing for, provision of, and accessibility to preventative and curative health services. Between 2007 and 2011, the proportion of the gross national product allocated to health varied from 3% in PNG and Fiji to 14% in Kiribati, with the majority of countries in the 5–8% range. In PNG, this translates to US\$35–40\$ per capita.

Almost all the countries with low mortality have near 100% primary education, with many having high levels of secondary and tertiary education. Countries with higher mortality have poorer education levels, and even then the figures hide

important gender disparities. There is often a higher proportion of male than female children enrolled, and girls are less likely than boys to complete primary education. The standard of primary education in a city school in Australia with state-of-the-art internet technology is vastly different from that in a remote village school in PNG, Solomon Islands or Vanuatu, where books and blackboards may be non-existent.

Maternal mortality rates in some of the countries are high, and in PNG, where only 52% of women deliver in health facilities, and in Timor Leste, the rates are alarmingly high. Neonatal mortality rates are correspondingly high, accounting for 50% of infant mortality. The common causes of neonatal death are infection, birth asphyxia and problems associated with low birth weight. The death of a mother puts her child at risk not only of neonatal but also of later infant and child mortality.

Transport systems to provide ready access to health services are essential. However, in many of the smaller island countries, transport is by dinghy or air. In the larger islands, road transport may simply not exist in some areas, and access is either by air to small airstrips or by foot. Where there are roads, they are likely to be unmetalled and at times unusable by vehicles. Security concerns may also hamper the delivery of services in some countries and some areas. Unfortunately, run down, inadequately staffed and inadequately supplied health facilities are common in some of the countries in the region, preventing health service delivery even when access is possible.

For some of the region's countries such as PNG, Solomon Islands and Vanuatu, population growth rates of >2.4% pose a major challenge to the provision of health, education and other services. Some of the smaller countries have much lower growth rates—in some, reflecting outward migration to countries such as Australia and New Zealand.

The leading causes of child mortality in the low-income countries in the region are communicable diseases: pneumonia, meningitis, diarrhoeal disease, malaria and, in some of the countries, tuberculosis (TB). Many deaths can be avoided with simple preventative and curative interventions. EPI programmes (immunisation programmes) are well established in all countries, but achieving high vaccination coverage has proved difficult in some. In several of the countries, fewer than 75% of children are protected by DPT3 (third dose of diphtheria, tetanus and pertussis vaccine, either as 'triple vaccine' or incorporated into other multiantigen vaccines), and, although routine measles-containing vaccine coverage at the 95% level required has been reached in a few countries including Fiji and Samoa, it is well below this in

CrossMark

To cite: Vince JD. *Arch Dis Child* 2015;100(Suppl 1):s63–s65.

Regional updates

Figure 1 Map of Oceania (Timor Leste not marked). Cartographer: Daniel Feher. <http://www.freeworldmaps.net/oceania/>

several others. Conjugate *Haemophilus influenza* type b vaccine has been introduced to all the countries, but pneumococcal conjugate vaccines, which have been part of the programme in Australia and New Zealand for some years, have yet to be introduced in most of the Pacific Island countries.

Safe water supplies are still not available for many communities, and the understanding of the importance of hygiene practices is limited in some population groups.

TB is a significant problem in several of the region's countries, with prevalences ranging from 1.7/100 000 in Fiji to 541/100 000 in PNG.⁴ Mortality rates from TB in children are not

easy to find, but, in high-prevalence countries, TB is a significant cause of death. In PNG hospitals, TB accounts for between 7% and 8% of admissions to the children's wards, has disproportionately high bed occupancy, and carries a case fatality of 13%. Multidrug-resistant TB is well established in PNG, presenting additional challenges.

Malnutrition is a significant problem in some of the Oceania countries, with undernutrition estimated to affect 18% of children in PNG, 12% in Solomon Islands and 11% in Vanuatu. As elsewhere in the world, malnutrition contributes both directly and indirectly to mortality.

Table 1 Mortality rates and related indices in some Oceania countries

Rank by U5MR	Country	U5MR		Infant mortality		Neonatal mortality 2012	% change in U5MR since 1990	Adjusted maternal mortality rate 2010	Population (1000 s)	GNI per capita (US\$)	Education status		Human development index
		1990	2012	1990	2012						ALR	1 ⁰ enrol	
41	Papua New Guinea	89	63	65	48	24	29	230*	7167	1790	62		0.466
43	Kiribati	94	60	68	46	22	36		101	2260			0.629
48	Timor Leste	171	57	129	48	24	67	300	1114	3670	58	91	0.576
66	Marshall Islands	49	38	39	31	16	23	140 ⁺	53	4140		99	
72	Solomon Islands	39	31	31	26	14	19	93	550	1130		88	0.53
88	Fiji	31	22	26	19	10	27	26	875	4200		99	0.702
100	Vanuatu	35	18	29	15	9	48	110	247	3080	83		0.626
100	Samoa	30	18	25	15	7	41	100	189	3220	99	93	0.702
125	Tonga	23	13	20	11	7	44	110	105	4240	90		0.71
157	New Zealand	11	6	9	5	3	49	15	4460	30 620		99	0.919
161	Australia	9	5	8	4	3	47	7	23 050	59 570		97	0.938

Data from Unicef State of the World's Children Report (2014) and United Nations Development Programme Human Development Report (2013).

*Probably higher. DHS survey of 2006 reported a figure of 730/100 000.

⁺Reported rate 2008–2012.

1⁰, primary school; ALR, adult literacy rate; GNI, gross national income; U5MR, under 5 mortality rate.

Countries in the Oceania region report low prevalences of HIV. PNG has a reported prevalence of 0.5% in the reproductive age group, and HIV accounted for 6.3% of the deaths in the paediatric ward at Port Moresby General Hospital in 2013.

While some of the countries in the region have reached or may well reach the MDG target, others have a long way to go. However, it is important to appreciate that progress has been made in all of the countries—often in spite of formidable constraints.⁵ The introduction and tighter control of the use of effective artemether combination antimalarials in malaria-affected countries and the introduction of insecticide-treated bed nets has led to a reduction in malarial morbidity and mortality. The introduction and appropriate use of oxygen concentrators for provision of oxygen has significantly reduced mortality in children with pneumonia and other severe illnesses. Some countries have well-established child health policies and strategic implementation plans, focusing on common problems and achievable goals. Many countries are focusing attention on reduction of neonatal mortality by simple and practical means. However, for some of the Oceania region countries, there are clear deficiencies in the provision of primary healthcare, and further progress will come not from focusing on sophisticated hospital care but simply by getting the basics of disease prevention and adequate management of the common illnesses right. A functioning health system that is adequately funded, equipped and staffed will form the basis for continuing progress, but the wider aspects of health—such as education and infrastructure development—are of equal importance.

There are many challenges to continuing progress in many of the Oceania countries. These include dealing with the unmet need for family planning to slow the population growth rate, planning for the effects of climate change (being felt most acutely in the Pacific atolls), and ensuring the rights of women, children and disadvantaged communities. Much will depend on the quality and the ability of governance at all levels to ensure equitable distribution of resources and sustainable development, and to minimise damaging environmental and social effects of mineral, petroleum, gas, forestry and other industries.

Paediatricians in the Oceania region have demonstrated commitment to child health in the widest sense—and it is imperative that they continue to do so.

Acknowledgements I am grateful to Professor Trevor Duke of the Centre for International Child Health, University of Melbourne for his helpful comments.

Competing interests None.

Provenance and peer review Commissioned; externally peer reviewed.

REFERENCES

- 1 State of the World's Children 2014. Basic Indicators Tables 1, Nutrition Table 2, Economic Indicators Table 7, Women Table 8 and Rate of Progress Table 10. Unicef New York. <http://www.unicef.org/sowc2014/numbers/>
- 2 United Nations Development Programme Human Development Report. 2013. <http://hdr.undp.org/en/content/human-development-report-2013>
- 3 Australian Bureau of Statistics 3302.0—Deaths, Australia, 2012. <http://www.abs.gov.au/ausstats/abs@.nsf/mf/3302.0/>
- 4 Tuberculosis Country Profiles. WHO Geneva. <http://www.who.int/tb/country/en/>
- 5 Lagani W, Mokela D, Oswyn G, *et al.* Papua New Guinea: real progress towards MDG-4 and real challenges. *Int Health* 2010;2:186–96.

Millennium Development Goals: progress in Oceania

John D Vince

Arch Dis Child 2015 100: S63-S65

doi: 10.1136/archdischild-2013-305618

Updated information and services can be found at:
http://adc.bmj.com/content/100/Suppl_1/S63

These include:

References

This article cites 1 articles, 0 of which you can access for free at:
http://adc.bmj.com/content/100/Suppl_1/S63#BIBL

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Topic Collections

Articles on similar topics can be found in the following collections

[Epidemiologic studies](#) (1818)
[Injury](#) (437)
[Trauma](#) (434)

Notes

To request permissions go to:
<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:
<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:
<http://group.bmj.com/subscribe/>