(67%) reported securing large furniture items in place within their home. Many parents were aware of the hazards highlighted (average 1–10 scale rating, 8.2), and found the campaign useful (average 1–10 scale rating, 7.3). In 50% of cases the potential to alter current practices perceived by carers was encouraging (average 1–10 scale rating, 5.3).

Conclusion A poster campaign highlighting hazards implicated in local deaths is deemed useful by parents. Further work through the use of focus groups and parental communication is required to identify how best to promote safety practices for future campaigns.

G459(P) IMPROVING THE INVESTIGATION OF UNEXPECTED INFANT DEATHS

¹J Garstang, ²C Ellis, ³F Griffiths, ¹P Sidebotham. ¹Division of Mental Health and Wellbeing, University of Warwick, Warwick, UK; ²Faculty of Health and Life Sciences, Coventry University, UK; ³Division of Health Sciences, University of Warwick, Warwick, UK

10.1136/archdischild-2015-308599.413

Aims Since 2008, in England, all unexpected infant deaths must be investigated jointly by police, health and social services. This study aims to learn of bereaved parents' and professionals' experiences of this joint agency approach (JAA) as well as assess the effectiveness of the JAA in determining causes and risk factors for deaths and use this knowledge to improve professional practice.

Methods

1. A mixed methods study of JAA investigation of SUDI cases in one English region; involving case note analysis, questionnaires and in-depth interviews with bereaved parents and the relevant professionals.
2. A descriptive study of outcomes of JAA SUDI investigation using Child Death Overview Panel (CDOP) data.

Results 23/111 families were recruited giving theoretical saturation. The median time between infants' deaths and parental study participation was 33 weeks; data collection took place between 2011–3. 25 professionals were interviewed. CDOP Form Cs were obtained for 65/70 (93%) SUDI cases dying during 2010–2.

Non-specialist police often arrived at the parents' home along with the ambulance; increasing parental distress.

Parents felt that the JAA provided information about why their baby died but offered minimal emotional support.

The joint home visit by police and paediatrician is a key investigative process and most parents found this helpful. Final case discussions were used to discuss relevant risk factors but not to determine the cause of death; in nearly all cases the final cause of death relied on post-mortem examination alone ignoring findings of death scene examinations. Many deaths fitted the diagnostic criteria for SIDS but despite this were labelled as unascertained.

Social care were only involved in 13/23 JAA investigations, in two cases without involvement there were safeguarding concerns.

Some Coroners were reluctant to share post-mortem examination reports with paediatricians preventing effective JAA investigations.

Conclusion Ideally, SUDI investigations should be carried out only by specialist clinicians who do this work frequently and the JAA fully integrated with social care and Coroners' investigations. There needs to be a clearer system for classifying unexplained SUDI. Police should reconsider their immediate response

to SUDI; parents would like more follow-up and bereavement support from professionals.

G460(P) NEUROCOGNITIVE DEFICITS AND COMORBIDITIES IN NEUROFIBROMATOSIS 1 – EXPERIENCE FROM A TERTIARY SERVICE

R Mithyantha, H Royden, J Acharya, Z Bassi. Department of Community Paediatrics, Alderley Children's Hospital NHS Foundation Trust, Liverpool, UK

10.1136/archdischild-2015-308599.414

Aims Neurofibromatosis type 1 (NF1) is a common genetic condition with multisystem involvement. Clinical presentation of this condition can be varied, requiring clinicians to be aware of associated comorbidities and complications. We relate our experience of children seen within a tertiary centre.

Methods We analysed 100 consecutive patients with an established diagnosis of NF1 seen in a tertiary NF1 clinic. Records of patients were reviewed to determine clinical presentation and associated range of comorbidities.

Results Of the 100 patients (M:F = 55:45; median age 11.6 years (range 9 months–21 years)), 57 had at least one neurodevelopmental concern ranging from Learning difficulties 32%, specific learning difficulties 2%, social communication difficulties including ASD in 12%, attention difficulties including ADHD 18%, significant behavioural difficulties 11%, coordination difficulties 8%, sleep problems 13%, anxiety and low self esteem 3%, epilepsy in 4%. Majority of these children were in mainstream schools, so appropriate support would need to be provided for.

7/100(7%) of patients developed intracranial tumours, 14% optic pathway gliomas, 8 had non-tumour cranial abnormality identified on imaging. 10 children required neurosurgical intervention. 30% of children developed scoliosis, 5/30 required surgical fixation. 4 children had pseudoarthrosis, including a late presentation at 10 years. 17% had refractory errors. 14% of children underwent investigations for poor growth, 2 required hormone replacement therapy. 6 patients underwent investigations for concerns relating to puberty (2 precocious puberty, 2 premature adrenarche, 2 delayed puberty). 3 patients were on treatment for hypertension (Renal artery stenosis 2, essential hypertension 1). Plexiform neurofibromas, which can cause pain and affect cosmetic appearance, were present in 27/100 patients (9 required surgery).

Conclusions Under recognition of associated neurocognitive deficits and neuropsychiatric conditions in children with a diagnosis on NF1, can have significant impact on their educational outcome. The varied spectrum of clinical presentation warrants a comprehensive assessment for this group of patients. Establishing networks between local services and specialist centres will assist in early identification and management of Neurofibromatosis 1 related complications.

G461(P) CHARACTERISTICS OF HAEMOPHAGOCYTIC LYMPHOHISTIOCYTOSIS AT A TERTIARY PAEDIATRIC CENTRE

F Welsh, R Hague. Royal Hospital for Sick Children, NHS Greater Glasgow and Clyde, Glasgow, UK

10.1136/archdischild-2015-308599.415