



OPEN ACCESS

Developmental and behavioural associations of burns and scalds in children: a prospective population-based study

Alan Emond,¹ Clare Sheahan,² Julie Mytton,³ Linda Hollén¹

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/archdischild-2016-311644>).

¹Scar Free Foundation Children's Burns Research Centre, University of Bristol, Bristol, UK
²Centre for Child and Adolescent Health, University of Bristol, Bristol, UK
³Centre for Child and Adolescent Health, University of the West of England, Bristol, UK

Correspondence to

Professor Alan Emond, Scar Free Foundation Children's Burns Research Centre, Oakfield House, Oakfield Grove, Bristol BS8 2BN, UK; alan.emond@bristol.ac.uk

Received 18 July 2016

Revised 15 October 2016

Accepted 23 October 2016

Published Online First

14 November 2016

ABSTRACT

Objective To investigate child developmental and behavioural characteristics and risk of burns and scalds.

Design Data on burns in children up to 11 years from 12 966 participants in the Avon Longitudinal Study of Parents and Children were linked to developmental profiles measured before the burn injury.

Measures Preinjury profiles of the children derived from maternal questionnaires completed in pregnancy, and at 6, 18, 42, 47 and 54 months. Injury data collected by questionnaire at 6, 15 and 24 months and 3.5, 4.5, 5.5, 6.5, 8.5 and 11 years of age.

Results Incidence: Burn rates were as follows: birth–2 years 71.9/1000/year; 2–4.5 years 42.2/1000/year; 5–11 years 14.3/1000/year. Boys <2 years were more likely to sustain burns, and girls had more burns between age 5 and 11 years. Medical attention was sought for 11% of burn injuries. *Development*: Up to age 2 years, burns were more likely in children with the most advanced gross motor developmental scores and the slowest fine motor development. Children with coordination problems at 4.5 years of age had increased risk of burns between 5 and 11 years. No associations were observed with cognitive skills. *Behaviour*: At 3.5 years, the Strengths and Difficulties Questionnaire scores and reported frequent temper tantrums predicted subsequent burns in primary school age. After adjustment for confounders, burns in the preschool period were related to gender and motor development, and in school-aged children, to frequent temper tantrums, hyperactivity and coordination difficulties.

Conclusion Child factors associated with increased risk of burns were male gender in infancy and female gender at school age, advanced gross motor development, coordination difficulties, hyperactivity and problems with emotional regulation.

BACKGROUND

The importance of burns and scalds as a cause of morbidity and mortality in young children is well documented.^{1–3} Every year over 50 000 children in the UK visit hospitals as a result of burns,⁴ and although many of these are minor injuries, 45% of all severe burns or scalds occur in children under 5 years of age. Most of the data regarding paediatric burn injury rates come from series of children who have attended emergency departments (EDs)⁴ or have been admitted to hospital.⁵ These data are biased by the health seeking behaviour of the parents, and social factors such as distance from home to the nearest ED, or whether the family have access to a car. Many burn incidents are

What is already known on this topic?

- Burns and scalds are common injuries to young children, and the environmental and family risk factors for burn injury are well described.
- Few studies have investigated the developmental and behavioural characteristics of the child related to the risk of sustaining a burn.
- Most published studies of paediatric burn injury come from series of children who have attended emergency departments or been admitted to hospital.

What this study adds?

- Analysis of parental reports of 12 966 participants in a large representative population-based birth cohort showed peak burn incidence rates in the home of 72/1000 children/year, with only 11% being medically attended.
- The child's gender affected burn risk differently at different ages: boys <2 years were more likely to sustain burns, and girls had more burns between 5 and 11 years of age.
- Developmental and behavioural profiles of the children collected *before* the burn injury demonstrated that motor development, coordination difficulties, hyperactivity and problems with emotional regulation had consistent independent effects on burn injury risk.

entirely managed at home, but very few population studies have considered these more minor injuries.⁶

Factors related to paediatric injuries include individual characteristics such as younger age and male sex,^{6–8} family factors such as single parent and number of other children in the home^{9 10} and social factors such as area deprivation.¹¹ Few studies^{12 13} have investigated the developmental and behavioural characteristics of the child related to the risk of sustaining a burn. This information is important to understand why children have such high rates of burn injury, and to highlight which children at which ages are particularly vulnerable, to inform preventative strategies.



CrossMark

To cite: Emond A, Sheahan C, Mytton J, et al. *Arch Dis Child* 2017;**102**:428–433.

The objectives of this study are to describe the incidence and distribution of burns sustained at home in children up to 11 years in a representative population cohort, and to define developmental and behavioural factors at different ages affecting the likelihood of sustaining a burn.

METHODS

Sample

The Avon Longitudinal Study of Parents and Children (ALSPAC) is an ongoing population-based cohort study. Pregnant women resident in the former Avon Health Authority with an estimated date of delivery between 1 April 1991 and 31 December 1992 were invited to take part, resulting in a cohort of 14 062 live births and 13 988 children alive at 12 months.¹⁴ Ethical approval for the ALSPAC study was obtained from the local research ethics committees and this study was monitored by the ALSPAC Law and Ethics Committee. More detailed information on the ALSPAC study is available on the website: <http://www.bristol.ac.uk/alspac>, which contains details of all the data that are available through a fully searchable data dictionary.¹⁵

Social and demographic background

Child and family features were prospectively collected before the burn injury. Maternal education and marital status were derived from questionnaires completed in pregnancy, and maternal parenting score (higher is better) was derived from questionnaires at 6 and 42 months. The Indices of Multiple Deprivation (IMD) were used based on the address of the family at different injury questionnaire time points: higher IMD scores indicate greater social deprivation. The ALSPAC Family Adversity Index (FAI, a derived index including 18 items reflecting family socioeconomic status) was used: the higher the FAI score, the greater the family adversity.

Developmental measures

The child's developmental progress was measured by the ALSPAC Development Scale (a parental questionnaire derived from the Denver Developmental Screening Test¹⁶); we used the gross motor and fine motor scales at 6, 18 and 42 months. A variable for handedness, based on six questions, was derived from parental questionnaire at 42 months. Coordination problems were reported by parents in a questionnaire at 54 months. The child's IQ and selective attention were measured at 8 years in a research clinic, using a short version of the WISC III¹⁷ and the Tests of Everyday Attention for Children.¹⁸ The child's behavioural profile was measured using the Strengths and Difficulties Questionnaire (SDQ)¹⁹ completed by a parent at 47 months. We used the hyperactivity and conduct problems subscales, as externalising behaviours have been shown to be associated with increased risk of injury.²⁰ Children were classified as having problems if categorised as 'abnormal' based on cut-offs for each SDQ subscale suggested by Goodman and Goodman.²¹ At 18 and 42 months, parents were also asked to rate how frequently their child had temper tantrums. The child's anxiety level at 7 years was reported by the parents using the Developmental and Well Being Assessment.²²

Outcome data

Burns were reported in parental questionnaires when the child was 6, 15 and 24 months, and 3.5, 4.5, 5.5, 6.5, 8.5 and 11 years of age. Parents were asked if their child had been burnt or scalded since the last questionnaire, and if so how many times. Further details were collected on the action taken by the

person with the child (burn treated by themselves or sought medical attention).

Statistical analyses

Analyses were conducted in STATA V.14 (StataCorp. 2015. *Stata Statistical Software: Release 14*. College Station, TX: StataCorp LP) for three different age periods: infants aged 0–2 years, pre-school children aged 2–4.5 years and primary school-aged children aged 5–11 years. Controls were those reported as having had no burn on all questionnaires within an age period.

Period prevalence, that is, the proportion of children sustaining a burn at some time during the specified age period (including multiple questionnaires), and incidence rates, defined as the number of new burns/person-time at risk, were calculated based on cases that responded to every questionnaire within each age period (complete follow-up). As follow-up time varied between questionnaires, we calculated person-time at risk separately for each questionnaire within an age period and then summed them to calculate the total. Prevalence rates are reported as percentages with 95% CIs. Incidence rates are reported as events/1000 children per year with 95% CI.

Using multivariable logistic regressions, we adjusted for confounders in a stepwise procedure: (1) adjusted for all child factors, (2) adjusted for all child factors and maternal factors, and (3) adjusted for child factors, maternal and household/socioeconomic factors. Gross motor and fine motor skills were entered as continuous predictors, and adjusted for age at questionnaire completion. For anxiety levels at age 7, IQ at age 8 and selective attention at age 8, we used a subset of burns between 9 and 11 to ensure that predictors were collected before the injury.

Multiple imputation (N=20 imputations) with chained equations using the 'mi' command was used to minimise bias due to missing data. Variables included in the imputation model were all of those included in the final logistic regression models, including the outcome, and those that predicted missingness in the predictors and confounders. We imputed our data up to the number of participants out of the total cohort that answered at least one of the burns questionnaires (N=12 996). More details are provided in online supplementary material S1.

Analyses were performed using all available data, complete case dataset and multiple imputation. Regression results are presented using the imputed results only. Univariate and multivariable tables using all available data and complete case analyses are supplied in online supplementary tables S2 and S3.

RESULTS

Sample

In total, 12 966 participants answered at least one of the burns questionnaires. The number of participants answering questions about burns varied from 11 433 at 6 months to 7415 at age 11 (see online supplementary table S4). As controls were those reported as having had no burn on all questionnaires within an age period, overall data were available for 9558 participants between birth and 2 years, 9039 between 2 and 4.5 years and 5906 between 5 and 11 years. Multiple burn injuries (>1 per age period) were reported for 8% (121/1484) of children with burns between birth and 2 years, 8% (75/992) between 2 and 4.5 years and 11% (86/795) between 5 and 11 years. The proportion of burn injuries in which medical attention was sought was 13.7% (189/1379) between birth and 2 years, 11.2% between 2 and 4.5 years, 7.7% (58/754) between 5 and 11 years.

Missing data

Between 16% and 22% of participants had missing data on predictors and confounders between birth and 11 years, and 27%–55% were missing outcomes across the age groups (see online supplementary materials S1 and S5). The proportion with burns in the imputed dataset was slightly lower compared with that in the all available data analysis (14.2% vs 15.5%). Factors that strongly predicted missingness in the outcome were low maternal education, higher family adversity (FAI) and high socioeconomic deprivation (IMD). Other variables predicting missingness in predictors and confounders (all $p < 0.05$) were young maternal age at delivery, high levels of antenatal depression, presence of domestic violence in pregnancy and non-white ethnicity.

Prevalence

Period prevalence (table 1) between birth and 2 years was significantly higher than that between 2–4.5 and 5–11 years, particularly in boys. Overall prevalence between 2 and 4.5 years was no different to that at 5–11 years, but prevalence in girls aged 5–11 years was slightly higher than that in boys aged 5–11 years. No gender difference was seen between 2 and 4.5 years.

Incidence

The rates of burns were highest between birth and 2 years with an overall incidence of 71.9/1000/year, dropping to 42.2 between 2 and 4.5 years, and to 14.3 between 5 and 11 years. Girls had lower rates than boys before 2 years, but slightly higher rates after 5 years (table 1).

Maternal and socioeconomic factors

Online supplementary material S2 contains tables showing proportions and means of the maternal and socioeconomic factors included in our analyses. Multivariable analysis (table 2) showed that the maternal factors most consistently associated with increased risk of child burn injury in infancy and the preschool period were higher maternal education, never having been married and high family adversity as measured by the FAI. Better maternal parenting score at 6 m showed a protective effect on risk of burns between birth and 2 years, but not between 2 and 4.5 years. Between 5 and 11 years, the only confounder that showed any association with burns was maternal education, as mothers with a degree were more likely to report

burns to their children. IMD scores were not associated with burn injury at any age.

Child factors

The results of multivariable analyses using the imputed dataset of child factors associated with burn injury at the different ages are shown in table 2. Results were similar using all available data and the complete case dataset (see online supplementary tables S3 and S6).

Gender: Boys were more likely to sustain a burn between birth and 2 years, whereas there was no difference between boys and girls in the age period of 2–4.5 years. Girls were slightly more likely to sustain burns in the school age period.

Motor development: Children with the most advanced gross motor developmental scores reported at 6 months were more likely to sustain burns between birth and 2 years (table 2 and figure 1). The odds of a burn increased by 3% for every one-point increase in motor scores. Fine motor development showed the opposite pattern, with the most advanced fine motor development being associated with a decrease in odds of a burn, but this effect was weaker and only apparent when adjusting for gross motor scores (figure 2). The effect of gender and gross motor scores remained after adjustment for confounders, whereas adjustment for family and socioeconomic factors attenuated the association with fine motor scores. Children with reported coordination problems at 4.5 years were much more likely to sustain burns between ages 5 and 11. Left-handed children did *not* have any increased risk of a burn between 5 and 11 years.

Cognitive development: No associations were observed between total IQ scores (OR (95% CI): 1.01 (1.00 to 1.01)) and attention scores at 8 years (1.12 (0.79 to 1.59)) with burns in the subset of children aged 9–11 years.

Emotional regulation: Frequent temper tantrums at 18 months were not associated with burns in the preschool period, but reported tantrums at 42 months predicted an increased risk of burns between ages 5 and 11. Maternal report of general child anxiety symptoms at age 7 showed no association with increased risk of burns at ages 9–11 (OR: 0.74 (0.33 to 1.69)).

Behavioural profile: The SDQ completed by the parent at 47 months was predictive of subsequent burns: hyperactivity scores showed consistent associations with injury between ages 5 and 11. Conduct difficulties were also associated with increased risk of burns, but this association attenuated after adjustment.

Table 1 Period prevalence and incidence rate of burns at the three different age periods, using cases with complete follow-up*

Period (months)	Population at risk	New cases	Period prevalence % (95% CI)			Incidence rate (per 1000/year (95% CI))		
			Total	Boys	Girls	Total	Boys	Girls
0–6	9339	142	13.5 (12.9 to 14.3)	15.2 (14.2 to 16.2)	11.8 (10.9 to 12.8)	71.9 (68.0 to 75.9)	81.1 (75.4 to 87.1)	62.1 (57.0 to 67.6)
6–15	9197	542						
15–24	8655	581						
24–38	8933	602	9.9 (9.3 to 10.6)	10.3 (9.4 to 11.2)	9.6 (8.7 to 10.5)	42.2 (39.5 to 45.1)	43.7 (39.9 to 47.8)	40.6 (36.8 to 44.7)
38–54	8331	284						
54–65	5658	132	9.7 (8.9 to 10.5)	8.8 (7.8 to 9.9)	10.6 (9.5 to 11.8)	14.3 (13.1 to 15.5)	12.9 (11.4 to 14.6)	15.7 (13.9 to 17.5)
65–77	5526	88						
77–103	5438	117						
103–140	5321	210						

*Complete follow-up refers to those participants responding to every questionnaire within each of the three age periods: 0–24 months=9339, 24–54 months=8933, 54–140 months=5658.

Table 2 Multivariable results from imputed data set (N=12 996)

Measure (non-reference group)	Age outcome (years)	Age at measure (months)	Unadjusted		Adjusted 1*		Adjusted 2†		Adjusted 3‡		p Value
			OR	95% CI	OR	95% CI	OR	95% CI	OR	95% CI	
Gender (male)	0–2	Birth	1.34	1.19 to 1.50	1.32	1.18 to 1.48	1.33	1.18 to 1.49	1.32	1.18 to 1.49	<0.001
Gross motor score	0–2	6	1.03	1.02 to 1.04	1.03	1.02 to 1.04	1.03	1.02 to 1.04	1.03	1.02 to 1.03	<0.001
Fine motor score	0–2	18	0.98	0.96 to 1.00	0.98	0.96 to 0.99	0.98	0.96 to 1.00	0.99	0.97 to 1.01	0.13
Gender (male)	2–4.5	Birth	1.11	0.98 to 1.27	1.10	0.97 to 1.26	1.10	0.97 to 1.26	1.10	0.97 to 1.26	0.15
Gross motor score	2–4.5	6	1.01	1.00 to 1.02	1.01	1.00 to 1.02	1.01	1.00 to 1.02	1.01	1.00 to 1.02	0.11
Fine motor score	2–4.5	18	0.98	0.96 to 1.01	0.98	0.96 to 1.00	0.98	0.96 to 1.00	0.98	0.96 to 1.01	0.16
Temper tantrums (often)	2–4.5	18	1.26	0.99 to 1.61	1.26	0.99 to 1.60	1.27	0.99 to 1.61	1.16	0.91 to 1.48	0.23
Gender (male)	5–11	Birth	0.78	0.64 to 0.95	0.78	0.64 to 0.95	0.78	0.64 to 0.95	0.78	0.64 to 0.95	0.02
Gross motor score	5–11	42	0.98	0.95 to 1.01	0.98	0.95 to 1.00	0.98	0.95 to 1.01	0.98	0.95 to 1.01	0.11
Fine motor score	5–11	42	1.00	0.98 to 1.03	1.02	1.00 to 1.04	1.02	1.00 to 1.04	1.02	1.00 to 1.04	0.05
Temper tantrums (often)	5–11	42	1.60	1.20 to 2.14	1.43	1.06 to 1.93	1.44	1.07 to 1.95	1.41	1.04 to 1.92	0.03
Hyperactivity (abnormal)	5–11	47	1.36	1.11 to 1.67	1.25	1.01 to 1.54	1.26	1.02 to 1.55	1.24	1.01 to 1.54	0.04
Conduct problems (abnormal)	5–11	47	1.40	1.16 to 1.69	1.24	1.01 to 1.54	1.23	0.99 to 1.53	1.20	0.96 to 1.50	0.10
Coordination problems (yes)	5–11	54	1.86	1.31 to 2.62	1.70	1.22 to 2.37	1.71	1.23 to 2.38	1.69	1.21 to 2.35	0.002
Handedness (left/mixed)	5–11	42	0.95	0.78 to 1.16	0.93	0.75 to 1.15	0.94	0.76 to 1.16	0.93	0.76 to 1.15	0.51

Full model 0–2 years, OR (95% CI) for confounders; maternal education (degree): 1.34 (1.14 to 1.58), maternal parenting score (worst decile): 1.19 (1.00 to 1.41), marital status (never married): 1.38 (1.20 to 1.60), FAI: 1.11 (1.07 to 1.15), IMD (worst quintile): 1.09 (0.91 to 1.31).

Full model 2–4.5 years, OR (95% CI) for confounders; maternal education (degree): 1.35 (1.12 to 1.62), maternal parenting score (worst decile): 0.89 (0.69 to 1.15), marital status (never married): 1.45 (1.22 to 1.72), FAI: 1.06 (1.02 to 1.10), IMD (worst quintile): 1.08 (0.86 to 1.35).

Full model 5–11 years, OR (95% CI) for confounders; maternal education (degree): 1.36 (1.12 to 1.65), maternal parenting score (worst decile): 1.26 (0.99 to 1.60), marital status (never married): 1.06 (0.83 to 1.34), FAI: 1.05 (1.00 to 1.11), IMD (worst quintile): 0.98 (0.73 to 1.30).

Motor scores are entered as continuous variables.

*Adjusted 1: all child factors.

†Adjusted 2: adjusted 1 + maternal education, maternal parenting score.

‡Adjusted 3: adjusted 2 + marital status, FAI and IMD. p Values reported for adjustment 3 only.

FAI, Family Adversity Index; IMD, Index of Multiple Deprivation.




Figure 1 Probability predictions of gross motor scores affecting risk of burns between birth and 2 years.




Figure 2 Probability predictions of fine motor scores affecting risk of burns between birth and 2 years.

DISCUSSION

This prospective study is unique because it has captured burn injuries in the home environment, with information on the child's development and behaviour collected before the injury occurred. Burns to children aged below 11 years in the home environment were commonly reported, with a peak in the age period of 15–24 months. The child's gender affected burns risk differently at different ages, and motor development, coordination difficulties and problems with hyperactivity and emotional regulation had consistent independent effects on burn injury risk.

The strengths of this study are that it uses a large representative population-based birth cohort, with burns prospectively

captured by parental report, thereby avoiding biases associated with attending hospital. The developmental and behavioural characteristics of the children were collected before the burn injury, and a range of family and social covariates were available for adjustment. There are few population-based studies of burns in children,^{23–25} most are based on hospital attendance,⁴ and none have prospectively collected information on development and behavioural profile before the injury.

Limitations are that the parental reports were not validated, and the recall period was variable for different questionnaires. Since parents are more likely to recall severe events than minor ones with the elapse of time,²⁶ recall bias could make the results

for minor burns less accurate, and mothers who were better educated reported more injuries. Overall, the study is likely to have underestimated the true incidence of paediatric burns. There were limited data on the severity of the burn, so that seeking medical attention was used as a proxy for injury severity. No information was available on whether the burns were intentional or related to neglect or maltreatment.^{27–28} The most important limitation is missing data due to loss to follow-up, which is inevitable in a study like this, so multiple imputation was used to correct for any biases introduced by attrition.

The peak incidence in burns in the second year of life is well documented, but the continuing injury rates reported through school age were higher than would be predicted from ED attendances.⁴ The gender differences changed noticeably with increasing age, and could be a result of girls being more exposed to thermal hazards in the kitchen than boys.⁵ The associations of burn injury with motor development may not surprise clinicians, but there is little literature on the connection between motor development and burns. The peak incidence of burns around 15 months, particularly in those with advanced gross motor skills, may reflect that this is the age when exploratory behaviour is greatest.²⁹ In contrast, poor coordination (perceived clumsiness) was related to burns risk in the older children. Hertog *et al*⁷ attributed clumsiness as being contributory in 57% of scald injuries in their cohort of children aged 0–4 years, although individual differences in motor ability were not analysed. Activity levels and emotional regulation were the most important behavioural factors affecting burn risk at later ages. We showed consistent effects of hyperactivity and to a lesser extent conduct difficulties on burn risk during school age, consistent with studies showing that activity levels in children with attention deficit hyperactivity disorder are related to higher injury levels,²⁰ although not all burn studies have shown an effect of hyperactivity.³⁰

CONCLUSIONS AND IMPLICATIONS

Minor burns are common in young children, and over 80% do not seek medical attention, illustrating the need for promoting good first aid practices in the home.³¹ Reducing hazards by environmental modification and limiting exposure by improving supervision are crucial strategies to prevent thermal injuries, but individual child characteristics are also independent risk factors, and change through childhood. As the child gets older, individual characteristics become relatively more influential than family or environmental factors. Anticipatory guidance is essential in educating parents about changing burn risks associated with developmental progress,³² and clinicians need to highlight to parents the increased risk of burns in those children with difficulties in motor coordination, emotional regulation and behavioural control.

Acknowledgements We are extremely grateful to all the families who took part in this study, the midwives for their help in recruiting them, and the whole ALSPAC team, which includes interviewers, computer and laboratory technicians, clerical workers, research scientists, volunteers, managers, receptionists and nurses. We thank Tom Norris for support in multiple imputation techniques.

Contributors AE conceptualised and designed the study, wrote and revised the manuscript, and approved the final manuscript as submitted. CS drafted sections of the initial manuscript, and approved the final manuscript as submitted. JM reviewed and revised the manuscript, and approved the final manuscript as submitted. LH undertook the statistical modelling and analysis, reviewed and revised the manuscript, and approved the final manuscript as submitted. LH had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Funding The UK Medical Research Council and the Wellcome Trust (Grant ref: 102215/2/13/2) and the University of Bristol provide core support for ALSPAC. This

research was funded by the Children's Burns Research Centre, which is part of the Burns Collective, a Scar Free Foundation initiative with additional funding from the Vocational Training Charitable Trust and the Welsh Government. The views expressed are those of the authors, and not necessarily those of the Scar Free Foundation or other funding bodies.

Competing interests None.

Patient consent Parental/guardian consent obtained.

Ethics approval Bristol LREC.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement Information about the ALSPAC study dataset is available on the website: <http://www.bristol.ac.uk/alspac>, which contains details of all the data that are available through a fully searchable data dictionary.

Open Access This is an Open Access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) license, which permits others to distribute, remix, adapt and build upon this work, for commercial use, provided the original work is properly cited. See: <http://creativecommons.org/licenses/by/4.0/>

REFERENCES

- 1 Agran PF, Anderson C, Winn D, *et al*. Rates of pediatric injuries by 3-month intervals for children 0 to 3 years of age. *Pediatrics* 2003;111:e683–92.
- 2 Hjern A, Ringbäck-Weitoff G, Andersson R. Socio-demographic risk factors for home-type injuries in Swedish infants and toddlers. *Acta Paediatr* 2001;90:61–8.
- 3 UK accident statistics; Home and Leisure Accident Surveillance System. <http://www.hassandlass.org.uk/reports/2002data.pdf>
- 4 Kemp AM, Jones S, Lawson Z, *et al*. Patterns of burns and scalds in children. *Arch Dis Child* 2014;99:316–21.
- 5 Brusselselaers N, Monstrey S, Vogelaers D, *et al*. Severe burn injury in Europe: a systematic review of the incidence, etiology, morbidity, and mortality. *Crit Care* 2010;14:R188.
- 6 Kohen DE, Soubhi H, Raina P. Maternal reports of child injuries in Canada: trends and patterns by age and gender. *Inj Prev* 2000;6:223–8.
- 7 den Hertog PC, Blankendaal FA, ten Hag SM. Burn injuries in the Netherlands. *Accid Anal Prev* 2000;32:355–64.
- 8 Hippisley-Cox J, Groom L, Kendrick D, *et al*. Cross sectional survey of socioeconomic variations in severity and mechanism of childhood injuries in Trent 1992–7. *BMJ* 2002;324:1132.
- 9 Wadsworth J, Burnell I, Taylor B, *et al*. Family type and accidents in preschool children. *J Epidemiol Community Health* 1983;37:100–4.
- 10 Scholer SJ, Hickson GB, Ray WA. Sociodemographic factors identify US infants at high risk of injury mortality. *Pediatrics* 1999;103:1183–8.
- 11 Reading R, Langford IH, Haynes R, *et al*. Accidents to preschool children: comparing family and neighbourhood risk factors. *Soc Sci Med* 1999;48:321–30.
- 12 Farmer JE, Peterson L. Injury risk factors in children with attention deficit hyperactivity disorder. *Health Psychol* 1995;14:325–32.
- 13 Bijur P, Golding J, Haslum M, *et al*. Behavioral predictors of injury in school age children. *Am J Dis Child* 1988;142:1307–12.
- 14 Boyd A, Golding J, Macleod J, *et al*. Cohort Profile: The 'Children of the 90s'—the index offspring of the Avon Longitudinal Study of Parents and Children. *Int J Epidemiol* 2013;42:111–27.
- 15 <http://www.bris.ac.uk/alspac/researchers/data-access/data-dictionary/>
- 16 Frankenburg WK, Dodds JB. The Denver Developmental Screening Test. *J Pediatr* 1967;71:181–91.
- 17 Wechsler D. *Wechsler Abbreviated Scale of Intelligence III*. San Antonio, TX: Harcourt Assessment Inc, 1999.
- 18 Manly T, Nimmo-Smith I, Watson P, *et al*. The differential assessment of children's attention: the Test of Everyday Attention for Children (TEA-Ch), normative sample and ADHD performance. *J Child Psychol Psychiatry* 2001;42:1065–81.
- 19 Goodman R. The strengths and difficulties questionnaire: a research note. *J Child Psychol Psychiatry* 1997;38:581–6.
- 20 Brehaut JC, Miller A, Raina P, *et al*. Childhood behavior disorders and injuries among children and youth: a population-based study. *Pediatrics* 2003;111:262–9.
- 21 Goodman A, Goodman R. Strengths and difficulties questionnaire as a dimensional measure of child mental health. *J Am Acad Child Adolesc Psychiatry* 2009;48:400–3.
- 22 Goodman A, Heiervang E, Collishaw S, *et al*. The 'DAWBA bands' as an ordered-categorical measure of child mental health: description and validation in British and Norwegian samples. *Soc Psychiatry Psychiatr Epidemiol* 2011;46:521–32.
- 23 Pickett W, Streight S, Simpson K, *et al*. Injuries experienced by infant children: a population-based epidemiological analysis. *Pediatrics* 2003;111:e365–70.
- 24 Faelker T, Pickett W, Brison RJ. Socioeconomic differences in childhood injury: a population based epidemiologic study in Ontario, Canada. *Inj Prev* 2000;6:203–8.

- 25 Fernández-Morales E, Gálvez-Alcaraz L, Fernández-Crehuet-Navajas J, *et al.* Epidemiology of burns in Malaga, Spain. *Burns* 1997;23:323–32.
- 26 Harel Y, Overpeck MD, Jones DH, *et al.* The effects of recall on estimating annual nonfatal injury rates for children and adolescents. *Am J Public Health* 1994;84(4):599–605.
- 27 Chester DL, Jose RM, Aldlyami E, *et al.* Non-accidental burns in children—are we neglecting neglect? *Burns* 2006;32:222–8.
- 28 Hobson MI, Evans J, Stewart IP. An audit of non-accidental injury in burned children. *Burns* 1994;20:442–5.
- 29 Warrington SA, Wright CM, ALSPAC study team. Accidents and resulting injuries in premobile infants: data from the ALSPAC study. *Arch Dis Child* 2001;85:104–7.
- 30 Petridoua E, Trichopouloua D, Mera E, *et al.* Risk factors for childhood burn injuries: a case-control study from Greece. *Burns* 1998; 24: 123–8.
- 31 British Burns Association. First Aid Position statement. <http://www.britishburnassociation.org/resources>
- 32 Gielen AC, Modena EH, Wilson MEH, *et al.* Randomized Trial of Enhanced Anticipatory Guidance for Injury Prevention. *Arch Pediatr Adolesc Med* 2001;155:42–9.