- Birth weight varies from 1.01 kg to 2.80 kg
- Out of 20 preterm babies 17(85%) had some form of pain relief.
- 3(15%) of babies had received no pain control measures.
- Non pharmacological measures were used in 15(75%) of babies.
- Pharmacological measures were used in 13(65%) babies.
- Oral sucrose was used in 10(50%)babies.
- 3(15%) babies received topical analgesia for ROP screening.
- Combined methods of pharmacological and non pharmacological pain relief were used in 13(65%)babies.

Conclusions Overall pain management was was not satisfactory in our unit.

Oral sucrose was used only in 50 per cent of the babies and also it was used inappropriately. In most of the babies it was given just before procedure or during the procedure. There were no records found for how many doses of oral sucrose given to any neonate in twenty four hours.

While giving vitamin K injection to babies they received no pain relief.

The use of combination methods of pain relief was relatively low.

Recommendations and action plan

Oral sucrose should be given at least 2 minutes before the procedure as per recommendations.

We can do better by following the use of combination methods.

While doing any procedure there should be documentation of pain relieving methods followed along with procedure notes.

It is also better to keep record of number of doses of sucrose given to any pre term baby in 24 hours.

P461 THE DIGITS DON'T ADD UP: EXOMPHALOS AND DIGITAL DYSPLASIA

Zaheera Yusuf*, Michael Boyle. *Rotunda Hospital, Dublin, Ireland*

10.1136/archdischild-2019-epa.797

Baby Girl S was born at term by elective caesarean section for an antenatal diagnosis of exomphalos. At delivery she was noted to have exomphalos major with distal dysplasia of the second, third, and fourth digits on her left hand. X-ray confirmed the second and third digit to have only two ossified phalanges, and the fourth digit to have only one ossified phalanx. On review of digital anomalies and exomphalos we found that digital dysplasia had not been previously described in infants with exomphalos.

Exomphalos is a midline abdominal wall defect occurring at the base of the umbilical cord affecting about 1 in 5000 pregnancies.¹ The defect contains abdominal viscera and is covered by a membrane of amnion, peritoneum, and Wharton's jelly.² Exomphalos can be further classified as exomphalos minor, where the opening is less than 4 cm and only contains the intestine, and as exomphalos major, where the opening is greater than 4 cm and/or with the liver inside the cord.³ Unlike gastroschisis, exomphalos has a membrane covering the abdominal contents, and is more likely to be part of a syndrome, or have associated anomalies.² Approximately half of infants have associated anomalies, including Trisomy 13 and

18, syndromes such as Beckwith-Wiedemann and Pentalogy of Cantrell, cardiac anomalies, pulmonary hypoplasia, VACTERL anomalies, and nervous system abnormalities such as holoprosencephaly and anencephaly.^{1,2,3}

Baby Girl S did not require resuscitation at delivery and the sac was intact. She was transferred to a tertiary paediatric hospital for surgical management. Her neonatal course was uneventful except for three moderate hypoglycaemic episodes. She had imaging, a cardiology review, and a genetics review. Her microarray and karyotype were normal, and she was believed to be non-syndromic with no other associated anomalies. She did not meet criteria for Beckwith-Wiedemann syndrome, despite the episodes of hypoglycaemia. Surgical management of the exomphalos involved a silastic pouch to allow gradual reduction of the defect.

This is the first documented case of digital dysplasia with exomphalos.

REFERENCES

- Southampton Children's Hospital. Exomphalos. 2014. Accessed at <http://www.uhs.nhs.uk/OurServices/Childhealth/NeonatalSurgery/ConditionsweTreat/Exomphalos.aspx>
- Neonatal Directorate Management Committee. Clinical Practice Guideline: Exomphalos (Omphalocele). Government of Western Australia North Metropolitan Health Service. 1 September 2014. <https://www.kemh.health.wa.gov.au/~media/Files/Hospitals/WNHS/For%20Health%20Professionals/Clinical%20Guidelines/NEO/WNHS.NEO.ExomphalosOmphalocele.pdf>
- Great Ormond Street Hospital for Children. Exomphalos. 2018. Accessed at <https://www.gosh.nhs.uk/conditions-and-treatments/conditions-we-treat/exomphalos>

P462 OUT OF HOUR BLOOD TRANSFUSIONS IN NICU AND CAN IT BE PREVENTED?

Navdeep Kaur Brar*, Prof Naomi McCallion. *The Rotunda Hospital, Dublin, Ireland*

10.1136/archdischild-2019-epa.798

Introduction Overnight blood transfusion carry increased risk of transfusion complication due to availability of less staff to monitor the patient and there is likely to be fewer medical and laboratory staff available to respond to the complication. As per hospital protocol non urgent red blood transfusions should be done during working hours so that there is less work load on night duty doctor, nurses and lab personnel. Other reason to order transfusion during working hours so that out of hours charges should be avoidable which is 150 Euros, as not always blood packs supply is available in the hospital. The aim of the audit to know why red blood cell transfusion top-ups were ordered out of hours and to see how can it can be avoided, thereby saving cost.

Methods Data was collected retrospectively from July 2017 till September 2017. The total of 77 transfusions took place on 26 patients as some patient had multiple transfusions. The collected data was entered directly into audit tool on excel sheet. The questionnaire includes the following questions: gestation, date of birth, weight at delivery, reason for transfusion, general status, haemoglobin level at the time of transfusion, the time of FBC taken and resulted, in working hours or out of hours, the time when blood transfusion ordered and collected, in working hours or out of hours, day of life at transfusion. The charts of patients were collected and reviewed. The guidelines used for this audit was the Neonatal administration and transfusion specific blood/blood components 2016 issued by Rotunda Hospital

Results Out of 77 transfusions, 40 (52%) were collected during the working hours while 37(48%) were collected out of hours. Nearly (40)52% patients were stable at the time of transfusion and 25(32%) were unstable and 12(16%) were not recorded. Median gestation and weight at birth was 28 +3 weeks and 1.14 kg respectively. Median Day of Life at transfusion was 22. Median Hb level before transfusion was 104.5.

Conclusion A significant proportion of out of hour's orders are collected 5 to 7pm - could there be time savings made to get these within the 5 pm cut off. More than half times when patients got transfused out of hours were stable. Could there be introduction of new guidelines in terms of transfusing treatment in stable patients. Out of hours transfusions carry risk of increased work load on staff and costs more if transfusion pack is not available in the hospital.

P463 AUDIT ON OFC MEASUREMENT IN THE ROTUNDA HOSPITAL NICU

Roisin Egan*, Sheiniv Giva, Adrienne Foran. *Dept. of Neonatology, The Rotunda Hospital, Dublin 1, Ireland*

10.1136/archdischild-2019-epa.799

Introduction Measurement of the occipital frontal circumference (OFC) is a quick and non-invasive method of monitoring head growth. Although it measures skull size, it also reflects overall brain volume and has been described as a 'widely used proxy of neural growth and brain size'. There is mounting evidence to suggest that suboptimal post natal head growth in term and preterm neonates is associated with neurodevelopmental delay. This is likely because of changes that occur during the 'critical' neonatal period [4]. Thus, serial OFC measurement not only has diagnostic but also prognostic significance.

Measurement is performed by placing a tape above the ears and midway between the forehead and the hairline to the occipital prominence. The first measurement should be performed at 36–48 hours after delivery to allow for resolution of oedema and moulding.

Methodology This review audit involved a spot review of 33 inpatient charts at the Rotunda Hospital NICU.

Exclusion criteria included patient instability at time of measurement and palliation.

The guideline of weekly OFC measurements for inpatient neonates was used as standard of care. OFC was plotted on sex specific growth charts.

Results 32 of the 33 patients met the inclusion criteria. Of these, n= 11 (34.4%) were very preterm (<31+6 weeks), n=9 (28.1%) late preterm (32–36+6 weeks) and n= 12 (37.5%) were term (>37 weeks). 50% (n=16) did not meet the weekly OFC measurement standard. Of these, 50% (n=8) were on CPAP or BiPAP at time of missed measurement. Whilst the majority (n=11) missed only one measurement, 3 missed two and 2 missed three; 4 of these were extremely premature (<28 weeks) and 2 had known abnormalities on cranial ultrasound. Four babies had no recorded OFC measurement. Furthermore the intervals between measurements varied, ranging from 1–14 days.

Conclusion The above audit demonstrates lack of adherence to our NICU guideline of weekly OFC measurements in 50% of our inpatients.

Recommendations We recommend identification of those who are at high risk for missed measurements (e.g: infants on CPAP/BiPAP) and the implementation of strategies for opportunistic measurements in this cohort. As part of the NICU team, it would be beneficial to involve other carers in this task while performing other routine measurements such as weight checks. Dissemination of the results of this audit will be important in increasing awareness and the need to reaudit once new systems have been put in place.

P464 BREASTFEEDING IN VERY LOW BIRTH WEIGHT

Hacer Ergin*, Nuran Özçiftçi Ertuğral, Özmert Ma Özdemir, Ceren Çirali. *Pamukkale University Faculty of Medicine Department of Neonatology, Denizli, Turkey*

10.1136/archdischild-2019-epa.800

Introduction Available research has shown that breastfeeding reduces the frequencies of sepsis, necrotizing enterocolitis, retinopathy of prematurity, and bronchopulmonary dysplasia (BPD), shortens reaching full enteral feeding and hospital stay, and improves neurodevelopment. Although sucking from breast prolongs breastfeeding duration in very low birth weight (VLBW, ≤1500 g) infants, breastfeeding prevalence are still below the target levels.

Methods According to unit protocol, all newborns are fed with their mothers' milk if there is no medical obstacle. Breast milk is given via orogastric tube, bottle or cup to the infants who can not suck from the breast. Orogastric feeding is accelerated through kangaroo mother care, and the application of breast milk smell and taste. Breastfeeding is supported after discharge by educated nurses at outpatient clinic. Data of VLBW infants discharged from our hospital and referred to our outpatient clinic were collected from medical records and their mothers. Feeding practices recorded according to corrected age at discharge and postdischarge were classified in three categories: exclusive breast milk (EBM), exclusive formula, and mixed (a combination of breast milk and formula). Exclusive breast milk and mixed feeding were named as breastfeeding.

Results This study included 115 VLBW infants discharged from our hospital (88) and referred to outpatient clinic (27). No significant difference was found between discharged and referred groups in terms of birth weight (1047 ± 272 vs 1046 ± 253 g), gestational age (28 ± 2.9 vs 28 ± 1.5 w), male gender (48.9% vs 51.9%), SGA rates at birth and on discharge [(15.9%, 30.7%) vs (7.4%, 40.7%)], moderate/severe BPD incidence (23.9% vs 40.7%), hospitalization duration (57 ± 27 vs 78.2 ± 31.7 d), and mother education (high school% 48.9 vs 66.7%) ($p > 0.05$).

At discharge, EBM (63.7% vs 18.5%) and breastfeeding rates (91.7% vs 77.8%) were significantly higher in discharged infants compared with referred infants ($p < 0.05$). EBM and breastfeeding rates at 0–6 months were 65.4% and 100% for discharged infants, 38.4% and 84.5% for referred infants. Breastfeeding rates were 95.5% and 88.9% in discharged infants, 81.8% and 66.6% were in referred infants at 6–12 months and 12–24 months. BPD existence reduced EBM rates at discharge (23% vs 76.8%) and 0–6 months (19.3% vs 35.3%) in discharged infants from our hospital ($p < 0.05$).

Conclusion The policies supporting breastfeeding during hospitalization and after discharge increases breastfeeding rates in VLBW infants.