anaemia, transient abnormal myelopoiesis, post-transplant lymphoproliferative disorder and Fanconi anaemia. Of the remaining 154 patients diagnosed with cancer, 88 (57.1%) were classified as 'green' according to the Royal Marsden Hospital 'Stop-light' system. 48 (31.2%) were classified as 'orange' and 18 (11.7%) were classified as red, qualifying for automatic referral to clinical genetics.

Conclusions A significant proportion of patients diagnosed with cancer in OLCHC may benefit from referral to clinical genetics and screening for underlying cancer predisposition syndromes.

P490 INCIDENCE AND 5 YEARS SURVIVAL RATES OF CHILDHOOD CANCER DIAGNOSED LESS THAN 1 YEAR OLD IN IRELAND 2007–2017

Jsun Loong Wong*, Frieda Clinton, Maria Carroll, Andrea Malone, Aengus O'Marcaigh, Owen Patrick Smith, Jane Pears, Michael Capra, Cormac Owens. *Our Lady's Children's Hospital, Dublin, Ireland*

10.1136/archdischild-2019-epa.826

Introduction Childhood cancer is the second commonest cause of death in children in developed countries. Childhood cancer survival rates has improved over the last decade with the advancement of diagnostic procedures and continuous improvement of multimodal treatment strategies. According to the National Cancer Registry of Ireland, an average of 137 cancers were diagnosed per year in children under the age of 15 between 1994 and 2014. The 5-year overall survival rate for this entire cohort was 81%. It is well documented that survival rates in paediatric oncology vary depending on specific cancer diagnosis, age of the patient at diagnosis and disease stage.

Aim We describe the incidence of cancer in very young children and the influence of age on outcome in children diagnosed with cancer in Ireland under the age of 1 between 2007–2017.

Method Data were extracted from the database of the National Children's Cancer Service (NCCS) based at Our Lady's Children's Hospital, Dublin. The data presented refer to the International Classification of Childhood Cancer (ICCC) version 3 with the inclusion of Langerhans Cell Histiocytosis (LCH).

Result 185 patients were diagnosed with paediatric cancer under the age of 1 at the time of their initial diagnosis. 159 (85.5%) patients were diagnosed with solid tumours or LCH. 86 (46.5%) were male. The average age at diagnosis was 5.24 (range 0–12) months. 19 (10.3%) patients were diagnosed following an abnormal antenatal scan. The most common cancers diagnosed were neuroblastoma (22.7%), CNS tumours (19.5%) and leukaemia (13.5%).

158 (85.5%) patients received treatment. 124 (78.4%) patients received chemotherapy as part of their treatment. 20 (12.7%) patients received radiotherapy. The mean age for radiotherapy was 1.35 years (range 0.17–4). 18 (9.7%) patients received stem cell transplant/rescue. The cumulative overall survival rate at 5 years is 80%. CNS tumours have the worst prognosis followed by leukaemias. 35 (18.9%) patients have relapsed during follow up, and 20 are alive in follow-up.

Conclusion Incidence rates of specific paediatric cancer types vary according to the age at diagnosis. There is no difference in survival rates between children diagnosed with cancer under 1 year of age and older children. Outcomes at the NCCS compare favourably with international standards.

P491 A RARE PLEURAL BASED MALIGNANCY IN CHILDHOOD

¹Thomas McGrath*, ¹Elizabeth Murphy, ¹Thara Persaud, ²Jane Pears, ¹Claire Purcell. *¹Tallaght University Hospital, Dublin, Ireland; ²Our Lady's Children's Hospital, Crumlin, Dublin, Ireland*

10.1136/archdischild-2019-epa.827

Introduction Malignant Mesothelioma (MM) is a rare invasive neoplasm arising from the mesothelial lining of several organs, primarily the pleura and peritoneum¹. A causal relationship with asbestos exposure is well established, however; MM can occur in the absence of well-defined risk factors and at an earlier age².

Case Presentation A 15 year old boy, presented to the Emergency Department with a 6-week history of cough, 5 kilogram weight loss and right sided pleuritic chest pain. He had attended the ED 4 weeks prior to this with chest pain, cough, temperature and a chest infection was diagnosed. Repeat chest x-ray demonstrated a right lower lobe consolidation with obscuration of the hemidiaphragm and a suspected pleural based opacification. Laboratory investigations showed a hypochromic microcytic anaemia, hyponatremia and raised inflammatory markers. A CT of thorax, abdomen and pelvis showed lobulated rind of pleural based soft tissue densities in the right thorax with downward invasion through the hemidiaphragm into the liver. On transfer to a tertiary Paediatric Oncology centre, A CT-guided biopsy demonstrated a malignant epithelioid neoplasm confirming a diagnosis of malignant mesothelioma with EWSR-ATF1 rearrangement. He was treated with 2 cycles of Pemetrexed and Cisplatin Chemotherapy (NCCP guideline). Follow up imaging demonstrated disease progression which coincided with clinical deterioration. NG feeding, PleuralX drain and Oxygen therapy are part of his treatment regimen. Vinorelbine monotherapy was commenced and a Palliative approach has been adopted. Currently he is stable awaiting further imaging.

Discussion MM is rare in young people. The genetic susceptibility to MM is complex and recently an EWSR1-ATF1 fusion transcript has been described in 4 younger patients without prior asbestos exposure². The prognosis in MM is poor but improved survival in younger patients has been observed¹. We wish to highlight this case because it is rarely encountered in a paediatric setting.

P492 ELECTROCLINICAL FEATURES AND SEIZURE OUTCOME OF PEADiatric PATIENTS WITH IMMUNE MEDIATED ENCEPHALOPATHY

Ilayda Uslu*, Fatih Varol, Serhat Guler, Halit Cam, Sema Saltik. *Istanbul University Cerrahpaşa-Cerrahpaşa Medical Faculty, Istanbul, Turkey*

10.1136/archdischild-2019-epa.828

Purpose Most patients with immune mediated encephalitis (IME) suffer from seizures, but data on the seizure outcomes in the paediatric population remains limited. This study was conducted to assess the clinical course and electroclinical features of paediatric populations diagnosed with seizures associated with IME.

Methods Total 15 patients with IME were identified from Cerrahpaşa Medical Faculty Paediatric Neurology Department and Intensive Care Unit from 2014 to 2018. The demographic, clinical and electrographical data of participants were recorded. Symptoms were classified as psychiatric, seizure, flu-

like symptoms, movement disorders and altered state of consciousness. Neuron specific antibodies (LG11, CASPR2, AMPAR1/2, NMDAR, GABABR) in CSF/serum, antibodies against GAD, thyroid peroxidase and paraneoplastic syndrome panel (anti-Hu, anti-Yo, anti-Ri, anti-Amphipysian, anti-Tr, anti-PCA-2, anti-Ma, anti-CV-2, anti-ANNA-3) were detected in the serum.

Results The mean onset age of patients were 8,3 years (2–13,8 years). Female and male ratio was 8:7. In all patients admitted to the hospital the most common symptom was seizure (73%). The other symptoms were psychiatric (20%) and altered state of consciousness (7%). Seven (46%) patients with acute seizures present with status epilepticus. The mean hospitalization time of patients were 38,5 days (14–70 days) and 20 days (3–50) at Intensive Care Unit. Five patients were mechanically ventilated.

Significant serum levels of antibodies were detected in 10 out of 15 patients. (GAD n = 8, anti-TPO n = 5) CASPR2 and NMDA in CSF were detected in different two patients. One patient diagnosed with FIRES. Oligoclonal banding or elevated IgG index were detected in 6 of 15 patients. Lymphocytic pleocytosis (40%) were detected in cerebrospinal fluid.

All patients had abnormal electroencephalogram findings but 35.7% of patients had abnormal brain magnetic resonance imaging findings at the time admission of the hospital.

Immunotherapy, firstline therapy (IVIG n = 14, pulse steroid n = 13, plasma exchange n = 7) and second line therapy (rituximab n = 1, cyclophosphamide n = 1) were contributed to treatment.

After the discharge of the hospital no more seizure were seen in 12 patients (80%) during follow-up period. At the last visit all of these patients had normal EEG findings. But in the other 3 patients epileptic activity was detected in their EEGs. Among them, only one patient was resistant to the antiepileptic treatment.

Conclusions In this study, we observed that seizures with IME were generally resistant to the treatment during the acute phase of the illness in pediatric population. But after proper and early immunotherapies, seizures remission was detected with good prognosis in most of them.

P493

LATE AND ATYPICAL PRESENTATION OF MECP2 MUTATION

¹Karen Kelleher*, ²Sally Ann Lynch, ¹Bryan Lynch. ¹Department of Neurology, Temple Street Childrens University Hospital, Dublin, Ireland; ²Department of Clinical Genetics, Temple Street Childrens University Hospital, Dublin, Ireland

10.1136/archdischild-2019-epa.829

Introduction Rett syndrome (RS) is an X-linked dominant neurodevelopmental disorder affecting girls, most commonly caused by MECP2 loss-of-function mutations. The typical phenotype of RS is characterised by a period of developmental stagnation in late infancy, followed by progressive dementia, developmental regression, stereotyped hand-movements, microcephaly and gait dyspraxia. There are well recognised variants such as Atypical Rett syndrome (Angelman-like), Zappella-variant (preserved speech variant), and Congenital Rett syndrome.

Objective We describe two cases of unrelated Irish females who presented towards the end of their first decade with unusual Rett phenotypes.

Results Both cases had normal early development. Mild/moderate developmental abnormalities were picked up at 2 years.

Prior to onset of epilepsy, both cases were in mainstream school with a special needs assistant and had upright and independent gait. Case 1 developed severe progressive scoliosis from preschool age.

Case 1 and Case 2 presented with severe, refractory epilepsy aged 10 and 9 respectively; consisting of a mixture of atonic and complex-partial seizures evolving to include epileptic spasms with epileptic encephalopathy in both. In both cases gait deterioration began as an increasingly crouched posture and shuffling gait progressing to wheelchair-dependence. Concurrently, was a profound regression in motor skills, cognitive abilities, very poor appetite and weight loss. Neither had features of hand-wringing or microcephaly.

Case 1 was notable for excessive drowsiness and development of progressive neuromuscular muscle weakness leading to frequent pneumonias, life-threatening choking episodes and a progressive deterioration in her respiratory function. She died, aged 13 years, of respiratory failure. Case 2, aged 13, does not have features of progressive bulbar palsy described in Case 1.

Extensive workup was unrevealing except for de-novo heterozygous mutations in their MECP2 genes. In Case 1, a novel variant of MECP2 was detected: A duplication of the majority of the protein-coding portion of exon 4. Isolated duplication of large sections exon 4 has never, to our knowledge, been described before. In Case 2 there was a deletion and seven base pair insertion in exon 4: 1006_1275delinsBATGCC(p.Leu336Aspfs*12), predicted to introduce a premature termination codon.

Conclusions Case 1 represents a unique case of MECP2 mutation, expanding our knowledge of the deleterious mutations leading to a Rett phenotype.

We suggest that young girls with late onset developmental regression, new onset refractory epilepsy and crouching gait be investigated for MECP2 gene mutations.

Early onset of scoliosis in an ambulatory child, and severe neuromuscular weakness are not previously known features of this condition.

P494

ETIOLOGY OF MICROCEPHALY: 5 YEARS EXPERIENCE

Şule Akyan*, Eda UTİNE. Hacettepe, Ankara, Turkey

10.1136/archdischild-2019-epa.830

Microcephaly is a developmental disorder in which the head circumference, measured as an occipitofrontal, is below the third circumferential, or at least 2 standard deviations (SD), relative to age, gender, and ethnic origin. The aim of this study is to evaluate the diagnostic approach of microcephaly in childhood and to determine the prevalence of various underlying causes/diseases, and to determine a standard diagnostic approach according to disease frequency. 1474 patients who had microcephaly and from prenatal 32nd gestational week to 18 years of age was admitted to any of the outpatient clinics of Hacettepe University Hospitals between July 2012 and July 2017. The patients' head circumference distribution (SDS) was -2/-3 with a rate of 24.97%, the ratio of patients with -3/-4 was 30.6%, the ratio of patients with -4/-5 was 18.3% and -5/- The rate of patients with 6 was 11.87%. According to the majority of patients, 68,72% were genetic, 4,21% were related with chronic disease, 3,73% were related to metabolic disease