Results During the study period 770 newborns were hospitalized in NICU due to various pathological conditions and 50 newborns have been selected with AKI. As the control group, 50 newborns were taken with comparable associated pathological conditions, but without kidney injury. The calculated prevalence of AKI in neonates was 6.4%. Most of involved neonates in the study in both groups (AKI and non AKI) were born at term (64% and 54%) with predominance of male (68% and 60%). The mortality rate was higher in newborns with AKI than control group (36% vs 24%). In half of newborn infants with AKI predominate severe score level, while in control group predominate median score level (42%). There is a significant difference between the mean score value in neonates with AKI and lethal outcome compared to neonates with AKI without lethal outcome (70.73 ± 18.6 vs. 40.2 ± 16.6).

Conclusion Acute kidney injury is a life threatening condition with still high mortality rate. The severity of the illness of hospitalized newborn infants in NICU is estimated by SNAPPE 2 score. The high score level is associated with the severity of the disease and higher mortality. Appropriate treatment of newborns with severe kidney injury improves the outcome and reduces the mortality of the disease.

P632 MORBIDITY AND MORTALITY OUTCOMES IN NEONATES WHO WERE TRANSFERRED FROM TO THE NEONATAL INTENSIVE CARE UNIT IN SFAX-TUNISIA

Amel Ben Hmed, Mouna Loukil, Chiraz Regaieg*, Manel Charfi, Amira Bouraoui, Nedha Hmida, Ridha Regaieg, Afef Ben Thabet, Abdellatif Gargouri. *Department of Neonatology, Hedi Chaker Hospital, Sfax, Tunisia*

10.1136/archdischild-2019-epa.963

Background In view of the multiplicity of places of birth and the scarcity of neonatal intensive care units, the transfer of sick newborns is necessary in several situations in order to provide them with the necessary care not available at the birth places. This transport exposes the newborn to some risks that may affect morbidity and mortality.

Aims of the study To study the characteristics of transported newborns and the conditions of their transfer in order to identify the factors that can influence morbidity and mortality.

Methods Retrospective chart review: the medical records of 273 newborns transported and admitted in NICU over a period of 3 years, from January 1, 2012 to December 31, 2014.

Results The transfer rate during the study period was 6.4%. The private sector contributed in 57.5% of the transfers while the rest of the cases were originated from the peripheral maternities (25.2%). The main reasons for transfer were neonatal respiratory distress (62.3%), neonatal infection (43.2%) and prematurity (24.2%). Surgical diseases (or other malformations) were addressed mainly from peripheral maternities with a significant difference compared to the private sector. 89.4% of cases were transported in an ambulance with a very limited equipment (incubator, oxygen cylinder, rarely a scope) and in the presence of a single nurse and exceptionally a doctor. Incidents occurred during the transfer and were observed in 6 cases: accidental extubation, depletion of the oxygen cylinder. Complications were observed in 48.4 of the cases, the main one was iatrogenic infection (19.75%). The death rate was 13.6%: this rate is influenced by hypoglycaemia, hypoxia, surgical diseases and the use of mechanical ventilation.

Conclusion The severity of the initial neonatal pathology conditioned the prognosis of the newborns transported which is also influenced by the conditions of the transport.

P633 NON-CATHETER-RELATED NEONATAL AORTIC THROMBOSIS REVEALING AN ASSOCIATION OF FACTOR V LEIDEN AND HYPERHOMOCYSTEINEMIA

Ben Thabet Afef, Manel Charfi*, Amel Ben Hamad, Chiraz Regaieg, Ridha Regaieg, Nedha Hmida, Amira Bouraoui, Abdellatif Gargouri. *Department of Neonatology, Hedi Chaker University Hospital, Sfax, Tunisia*

10.1136/archdischild-2019-epa.964

Introduction Non-catheter-related aortic thrombosis is a rare condition in neonates. It may be life threatening or leads to severe complications. It occurrence requires looking for an underlying congenital prothrombotic condition.

Methods We report a case of a spontaneous aortic thrombosis in a newborn revealing an association of factor V Leiden and hyperhomocysteinemia.

Results A full term male was born by c-section. He presented immediate severe respiratory distress. Echocardiography performed at the second day of life, showed persistent pulmonary hypertension. He required high frequency oscillatory ventilation and inhaled monoxide administration. Initial respiratory stabilization was noted. But at 11 days age, we noted an increase in oxygen requirement; a tachycardia, a hepatomegaly, an edema and femoral pulses were no more detected. Control echocardiography showed a left ventricular dysfunction with an ejection fraction of 30%. The abdominal doppler ultrasound found an extensive thrombosis in infrarenal abdominal aorta. Thrombolytic treatment was not administrated as a sub-arachnoid hemorrhage was found in the cerebral ultrasound. After 48 hours of mechanical ventilation and inotropic support, hemodynamic and respiratory stabilization was obtained. Control echocardiography at day 14, showed an ejection fraction of 50%. The biological assessment revealed heterozygosis R506Q mutation for the factor V (factor V Leiden) and heterozygosis MTHFR C677T mutation (hyperhomocysteinemia). The aortic thrombosis has been spontaneously lysed and disappeared within 10 months. At 4 years, his physical examination is normal.

Conclusions Till now, there are no clear established guidelines concerning the management of arterial thrombosis. We insist that the blood clotting screen must be systematic and complete to look for association of congenital prothrombotic conditions which would increase the thrombotic risk.

P634 HOSPITAL MORTALITY OF FULL-TERM NEWBORNS: FACTORS AND CAUSES

Amel Ben Hamed, Ilyes Regaieg, Mouna Loukil, Manel Charfi, Chiraz Regaieg*, Amira Bouraoui, Ridha Regaieg, Nedha Hmida, Afef Ben Thabet, Abdellatif Gargouri. *Department of Neonatology, Hedi Chaker Hospital, Sfax, Tunisia*

10.1136/archdischild-2019-epa.965

Introduction Full-term newborns mortality's is a public health issue because of its unexpected impact either on the infant's family and the medical staff, as opposed to the preterm death. Therefore, we aim to determine factors and causes associated with hospital mortality of full-term newborns.