Correction: *Overweight and obese children do not consult their general practitioner more often than normal weight children for musculoskeletal complaints during a 2-year follow-up*

In their paper 'Overweight and obese children do not consult their general practitioner more often than normal weight children for musculoskeletal complaints during a 2-year follow-up' (*Arch Dis Child* 2018;103:149–54) the authors have found an error in the odds ratios in table 2. The statistics program accidentally inverted the odds ratios. These have now been corrected in table 2 below. This error does not affect the conclusions of the paper.

Table 2 Number of patients with musculoskeletal consultations during 2 year follow-up

	Total (n=617)	OW (n=111)	NW (n=506)	Or (95% CI)	Overall p-value	Adjusted OR* (95% CI)	Adjusted P value*
Any musculoskeletal consultation, # of patients (%)	260 (42.1)	56 (50.5)	204 (40.3)	1.51 (1.00 to 2.28)	0.05	1.36 (0.86 to 2.16)	0.19
Lower extremities consultation, # of patients (%)	164 (26.6)	36 (32.4)	128 (25.3)	1.42 (0.91 to 2.12)	0.12	1.30 (0.79 to 2.16)	0.30
Upper extremities consultation, # of patients (%)	104 (16.9)	20 (18.0)	84 (16.6)	1.10 (0.65 to 1.89)	0.72	1.05 (0.58 to 1.91)	0.88
Other body parts consultation (back, neck), # of patients (%)	65 (10.5)	13 (11.7)	52 (10.3)	1.16 (0.61 to 2.21)	0.66	1.32 (0.65 to 2.70)	0.44
Unclear/not registered where complaint is located, # of patients (%)	42 (6.8)	9 (8.1)	33 (6.5)	1.27 (0.59 to 2.73)	0.55	1.28 (0.52 to 3.12)	0.59

*adjusted for gender, age, marital status, sex.
NW, normal weight; OW, overweight.

© Author(s) (or their employer(s)) 2020. No commercial re-use. See rights and permissions. Published by BMJ.
Arch Dis Child 2020;**105**:e2. doi:10.1136/archdischild-2017-313118corr1