Conclusions Post-transplant management is individualised based on multiple factors such as clinical conditions e.g. renal/liver impairment and whether other agents such as ATG or steroids are being used.¹ The lack of documentation around the treatment decisions made it difficult to explain deviations from standards in this audit. Ciclosporin standards were not met completely but were most likely unfeasible due to a narrow target range and the time between first dose and level monitoring. There did not appear to be a clear association between standards not being met and episodes of rejection. It would be beneficial to repeat this as a larger, prospective audit using revised standards.

REFERENCES

1. Costanzo MR, Costanzo MR, Dipchand A, *et al.* The international society of heart and lung transplantation guidelines for the care of heart transplant recipients. *J Heart Lung Transplant* 2010;**29**:914–56.
2. Sandimmun Concentrate for Solution for Infusion 50 mg/ml - Summary of Product Characteristics (SmPC) - (eMC) [Internet]. [cited 2018 Sep 2]. Available from: <https://www.medicines.org.uk/emc/product/1036>

P56 WITHDRAWN AS AUTHOR REQUESTED IT IS NOT PUBLISHED

P57 CONTINUOUS DRUG DELIVERY IS SIGNIFICANTLY AFFECTED BY RELATIVE HEIGHT CHANGES BETWEEN PATIENT AND SYRINGE DRIVER

¹Aisha Zahid, ²Andrew Wignell*, ²Dusan Raffaj, ²Patrick Davies. ¹School of Medicine, University of Nottingham; ²Nottingham University Hospitals NHS Trust

10.1136/archdischild-2020-NPPG.65

Aims Syringe drivers are the principle method of giving continuous infusions of important drugs to patients. Many of these drugs are critical for the maintenance of normal physiology. Anecdotal evidence abounds of severe patient instability on movement of syringe drivers during infusion. Our objective was to define the variation in drug delivery seen in three different syringe drivers, with changes in relative height between the syringe driver and the end of the giving set.

Methods Three syringe drivers (Alaris CC (Becton Dickinson), Perfusor Space (B Braun), and Synamed µSP6000 (Arcomed)) were analysed for reliability of flow at 0.5, 1, 2, and 5 ml/hr. A small air bubble was introduced into the giving set, and the progression of this was documented before and after a vertical movement of the syringe driver by 25 or 50 cm upwards or downwards relative to the delivery port.

Results For all pumps, delivery was interrupted on movement of the pumps downwards, and a bolus was given with movement of the pump upwards. Delivery halted at lower pump speeds for longer than higher pump speeds. The maximum delivery interruption was 11.8 minutes. Boluses given on moving the pump up were calculated as the equivalent number of minutes needed to deliver the bolus volume at steady state. The maximum bolus given was equivalent to 15.8 minutes of delivery. We were unable to eliminate the effects seen by very slow, steady movement of the pumps up or down. Static height differences made no difference to delivery.

Conclusions Syringe drivers should not be moved vertically in relation to the patient. Critical drug delivery is interrupted for up to 12 minutes with relative downward movements, and significant boluses of drugs are given with relative upward movements. As far as possible, elimination of relative height movements is advised, and extreme caution is necessary if any movements are unavoidable.

P58 MANAGING ACCESS TO 'OUT OF HOURS' MEDICATIONS IN A TERTIARY PAEDIATRIC HOSPITAL

Mary Worrall*, Elizabeth Devlin, Moninne Howlett. *Children's Health Ireland at Crumlin*

10.1136/archdischild-2020-NPPG.66

Aims Many UK and Irish hospitals provide a Monday to Friday pharmacy service; automated dispensing cabinets and hospital-wide information systems remain uncommon. Locating and accessing out of hours (OOH) medications can be a significant workload for nursing staff. In an Irish 230-bed tertiary paediatric hospital processes involve: nursing staff contacting other wards by telephone to source items; completion of an 'Out of Hours Requisition'; and either transferring stock between wards or contacting nursing administration staff (NAS) to access stock from the Pharmacy Department. A quality improvement project was undertaken to: measure current levels of OOH medications; identify areas for improvement; implement and assess impact of new processes.

Methods 'Out of Hours Requisition' data were entered into a custom-built database and analysed for the period January - December 2018. The findings were discussed with nursing staff and NAS. Improving processes for locating medications was identified as a key area for improvement. The 'Out of Hours Requisition' form was amended to provide clearer instructions for completion. Using data from clinical area stock lists, a searchable Medicines Locator Database was developed and made accessible to all staff in the pharmacy department, clinical areas and the NAS office in December 2018 enabling staff to remotely identify the location of all medications stocked in the hospital. Data for 'Out of Hours Requisitions' for the period January - May 2019 were collated, analysed and compared with data from the same time period in 2018 (January - May). Microsoft Excel® was used for data collection, analyses and development of the Medicines Locator Database.

Results A total of 1747 OOH medications were accessed by NAS from pharmacy in 2018, 746 during the period January - May 2018. Anti-microbial agents (36%) were most common, with requests originating from 16 clinical areas. Request from the paediatric intensive care units (36%) and the surgical/orthopaedic ward (36%) were most frequent. 515 medications were accessed in the first 5 months after the introduction of the Medicines Locator Database (January - May 2019). This represents a 35% reduction in the number of medications dispensed in the same time period in 2018. No changes to the types of medications were identified, but some differences in clinical areas were found.

Conclusion This significant reduction (35%) in numbers of medications accessed out of hours, and the corresponding reduction in workload for NAS, demonstrates the benefits of reviewing medication management processes. Further substantial time savings for nursing staff locating stock at ward level are likely. Readily-available technology can be successfully