high risk for AKI to enable preventative measures to be implemented. This retrospective audit aimed to describe the epidemiology of AKI at a tertiary paediatric centre in order to prioritise interventions.

Methods All inpatients with an AKI stage 2 or 3 alert between 1st March and 31st May 2017 were included in this audit. The electronic healthcare record for each patient was accessed and data extracted.

Results Over a 3 month period there were 354 AKI stage 2 or 3 alerts accounting for 125 AKI episodes affecting a total of 101 patients. Males (53.5%) and females (46.5%) were evenly represented. Children aged between 1 month and 2 years were most frequently affected (37%). 22% of patients were under the care of cardiology, 12% under general paediatrics, and 11% under respiratory. The majority of episodes lasted only 1 day, with 55 episodes (44%) lasting for 2 or more days. 51% of AKI episodes were associated with exposure to at least one nephrotoxic medication during, or in the 7 days preceding, the event. The most common medications were furosemide (14%), piperacillin with tazobactam (14%) and ibuprofen (11%). Overall the cause of AKI was infrequently documented (37% of episodes), however, where documented, the leading causes were cardiac surgery (28%) and nephrotoxic medications (20%).

Conclusion Cardiac surgery represents a significant risk factor for the development of AKI in children. In addition, over half of children who developed AKI were exposed to nephrotoxic medications, and this was the second most common identified cause. Future AKI management strategies should focus on delivering improvements for these groups of patients to reduce the impact of AKI.

G328(P) ABSTRACT WITHDRAWN

G329(P) INCIDENCE OF AMPHOTERICIN-INDUCED ACUTE KIDNEY INJURY IN A PAEDIATRIC POPULATION: A SYSTEMATIC REVIEW

¹D Mistry, ^{2,3}L Oni, ^{3,4}SJ McWilliam. ¹Medical School, University of Liverpool, Liverpool, UK;

²Department of Paediatric Nephrology, Alder Hey Children's NHS FT, Liverpool, UK;

³Department of Women's and Children's Health, University of Liverpool, Liverpool, UK;

⁴MRC Centre for Drug Safety Science, University of Liverpool, Liverpool, UK

10.1136/archdischild-2020-rcpch.284

Aims The aim of this project was to perform a systematic literature review to determine the incidence of Amphotericin-induced Acute Kidney Injury (AKI) in the paediatric population.

Methods A systematic review was performed to obtain articles until October 2018 using Scopus and Medline, with additional articles obtained using the bibliographies from selected studies. Studies were chosen if they included an intervention of Amphotericin with monitoring of renal function and a declaration of how AKI would be defined. All types of studies were considered and those retrievable in full text and English were included.

Results Of sixty-three articles identified, seventeen were available in full text. A further ten studies were rejected due to not mentioning dose, duration or outcomes. The remaining seven studies totalled 279 patients aging between

0–15 years. The ratio of boys to girls was 50:50 (demographic data reported in only 4 of 7 studies). Of the seven studies none were randomised controlled trials. Visceral Leishmaniasis was the indication for treatment with liposomal Amphotericin in 6 of the 7 studies. The dose of Amphotericin ranged from 15–30 mg/kg and a duration of 2 – 21 days. The total incidence of AKI was found to be 1%, with the pRIFLE criteria being the most commonly used method of defining AKI, however many of the studies did not explicitly define AKI in their outcome criteria, leading to low identification rates. The optimal dose range is 3–4 mg/kg for a regime of 5-day treatment followed by a further dose on day 10. Additional outcomes included risk factors for AKI with underlying infection being the most consistent.

Conclusion This systematic review found an incidence of Amphotericin-induced AKI in a paediatric population of 1%. Comparatively, adults show an incidence rate of around 50%, suggesting large differences between the two cohorts. There were limitations to this study, mainly due to the lack of standardisation in the definition of drug-induced AKI in children across the studies. Improved reporting of Amphotericin-induced AKI, using standardised AKI definitions, is required to confirm the true incidence in children.

G330(P) A SYSTEMATIC REVIEW OF THE RISK FACTORS ASSOCIATED WITH VANCOMYCIN-INDUCED ACUTE KIDNEY INJURY IN CHILDREN

¹C Williams, ^{2,3}L Oni, ^{3,4}SJ McWilliam. ¹Medical School, University of Liverpool, Liverpool, UK;

²Department of Paediatric Nephrology, Alder Hey Children's NHS FT, Liverpool, UK;

³Department of Women's and Children's Health, University of Liverpool, Liverpool, UK;

⁴MRC Centre for Drug Safety Science, University of Liverpool, Liverpool, UK

10.1136/archdischild-2020-rcpch.285

Aims Vancomycin has been suggested to contribute to acute kidney injury in adults, however few studies have addressed its role in the paediatric population. The primary aim of this paper is to review the evidence surrounding the risk factors involved in the development of paediatric acute kidney injury as a result of vancomycin administration.

Methods A systematic search was performed in November 2018 on PubMed, Web of Science and Scopus using ((AKI OR acute kidney injury OR nephrotoxic* OR renal injur* OR renal insufficiency OR kidney damage) AND (vancomycin OR vancocin) AND (p*ediatric OR child* OR infant* OR child* OR adolescen* OR neonat*)). Eligible studies were meta-analyses, clinical trials, observational studies or case series of vancomycin use in a paediatric population (0–18 years) reporting outcomes of nephrotoxicity.

Results Of 1021 records identified, sixteen retrospective cohort studies were eligible for inclusion in the review, totalling 10,575 patients with an overall vancomycin-induced acute kidney injury (v-AKI) incidence of 13.3%. Thirteen studies considered the effect of co-administration of vancomycin with other nephrotoxins. Use of concomitant nephrotoxic medications was associated with a significantly increased risk of AKI. Co-administration of aminoglycosides, piperacillin-tazobactam, vasopressors, furosemide (and other loop diuretics), ACE inhibitors and NSAIDs were all found to be significantly associated with increased risk of AKI in at least one study. However, there was significant variation between studies. Six

studies identified a serum trough vancomycin concentration ≥ 15 mg/L to be an independent risk factor for AKI. To a lesser extent, high dose, duration of treatment and ICU admission were also identified in some studies as risk factors for v-AKI.

Conclusion Incidence of vancomycin-induced kidney injury in children is high. Patients who are high-risk should be identified and monitored more closely in view of reducing the number of cases diagnosed. In particular, concomitant nephrotoxins should be avoided. Well-designed prospective studies are required to determine the mechanism of vancomycin-induced kidney injury and the contribution from the above risk factors.

G331(P) ARE WE MISSING PATHOLOGICAL CAUSES OF PROLONGED JAUNDICE BY STREAMLINING OUR INVESTIGATIONS?

Z Jacob, S Khan. Royal Aberdeen Children's Hospital, NHS Grampian, Aberdeen, UK

10.1136/archdischild-2020-rcpch.286

Aims Prolonged jaundice is common, occurring in 2–15% of all neonates. NICE guidelines published in 2010 on screening babies with prolonged jaundice were debated and were felt to result in an excess of unnecessary investigations in healthy babies with an otherwise unremarkable examination. These investigations included total and conjugated bilirubin, urine culture, G6PD where ethnically appropriate, full blood count, blood group determination and Coomb's test.

We initially audited 50 of our patient assessments and investigations against advised national guidelines.

Subsequently, we introduced a simple proforma to assist clinicians in assessing for red flag features on history and examination, and reduced recommended investigations to total and conjugated bilirubin only if baby did not have any concerning features on review.

The purpose of this re-audit was to assess whether this change in practice was safe, or had it hampered our ability to identify pathological jaundice, and did it result in better use of resources and improved patient/parent satisfaction.

Methods This prospective re-audit study analysed 105 proformas over a six month period for data on a number of parameters including gestation, age at referral, feeding, weight gain, urine and stool colour.

Outcomes for patients with unconjugated or conjugated hyperbilirubinaemia or those who were admitted were followed up.

The gastroenterology team was consulted for information on any patients who were screened during the audit period and subsequently diagnosed with pathological jaundice.

Results No patients with a diagnosis of pathological jaundice were 'missed' during either audit cycle.

One patient with conjugated hyperbilirubinaemia was diagnosed with alpha-1-antitrypsin deficiency having been flagged up by the streamlined screening process.

Our rate of follow-up investigations increased from 10% to 24.7% in the re-audit cycle.

3.8% of patients were screened too early.

Conclusions The proforma was effective in identifying patients with potentially pathological causes of prolonged jaundice. Reducing investigations resulted in significant

savings (potentially £107 per patient) and better use of NHS resources.

We have since updated our intranet guidelines, aiming to educate practitioners and reduce unnecessary referrals.

We plan to assess whether we can safely reduce our follow-up investigation rate.

G333(P) SERUM BIOMARKERS, BUT NOT DUAL-ENERGY X-RAY ABSORPTIOMETRY (DXA), PREDICT BONE MINERAL DENSITY IN CHILDREN WITH CHRONIC KIDNEY DISEASE

¹AD Lalayannis, ²NJ Crabtree, ³V Askiti, ³A Mitsioni, ⁴M Fewtrell, ⁵A Kaur, ⁶DV Milford, ¹R Shroff. ¹Paediatric Nephrology, Great Ormond St Hospital and UCL Institute of Child Health, London, UK; ²Paediatric Endocrinology, Birmingham Women's and Children's NHS FT, Birmingham, UK; ³Paediatric Nephrology, 'P and A Kyriakou' Children's Hospital, Athens, Greece; ⁴Childhood Nutrition Research Centre, UCL Institute of Child Health, London, UK; ⁵Paediatric Nephrology, Manchester University NHS FT, Manchester, UK; ⁶Paediatric Nephrology, Birmingham Women's and Children's NHS FT, Birmingham, UK

10.1136/archdischild-2020-rcpch.287

Introduction Currently available biomarkers and Dual-energy X-ray Absorptiometry (DXA) are thought to be poor predictors of bone mineral density (BMD). The 2017 KDIGO guidelines on CKD-MBD propose using DXA if it will affect patient management. We set out to determine the clinical utility of DXA and routine clinical biomarkers by comparing them with tibial cortical BMD measured by peripheral Quantitative Computed Tomography (pQCT), which clearly defines cortical and trabecular compartments and predicts future fracture risk.

Methods We performed a prospective multi-centre cross-sectional study in 97 children (5–19 years) with CKD stages 4–5 and on dialysis. Children underwent hip and lumbar spine DXA and tibial pQCT. All DXA and pQCT data were expressed as age corrected Z-scores, and lumbar spine DXA was additionally corrected for height (BMAD Z-score). Imaging findings were correlated with routine serum biomarkers.

Results Hip Z-score and lumbar spine BMAD Z-scores correlated significantly with each other ($p < 0.001$). Hip Z-scores and lumbar spine BMAD Z-scores showed a strong correlation with tibial trabecular BMD ($p < 0.001$), whereas they did not show any correlation with tibial cortical BMD. Neither the hip nor the lumbar spine measures correlated with any biomarkers.

PTH showed a positive correlation with tibial trabecular BMD ($p = 0.024$) and a negative correlation with cortical BMD ($p < 0.001$). Tibial cortical BMD was also associated with alkaline phosphatase ($p = 0.014$) and total calcium ($p = 0.019$).

On multivariable linear regression analysis the significant and independent predictors of tibial cortical BMD were PTH ($\beta = 0.355$, $p < 0.001$), alkaline phosphatase ($\beta = 0.238$, $p = 0.011$) and serum calcium ($\beta = 0.184$, $p = 0.052$), which together predicted 26% of variability in tibial cortical pQCT.

Conclusions Routinely used biomarkers- calcium, alkaline phosphatase and PTH- when used together are weak to moderate predictors of BMD. No associations were seen with hip or lumbar spine DXA. DXA is not a clinically useful tool and should not be performed routinely.