G531(P) A COHORT STUDY OF WHETHER PARENTAL SEPARATION AND LACK OF CONTACT WITH A PARENT PREDICTS DISEASE SEVERITY AT DIAGNOSIS IN YOUNG PEOPLES CHRONIC FATIGUE SYNDROME/MYALGIC ENCEPHALOMYELITIS

¹N Ayadi O'Donnell, ²A McCourt, ²T Segal. ¹Children and young people's services, UCL Hospital FT, London, UK; ²Medical School, UCL, London, UK

10.1136/archdischild-2020-rcpch.449

Aims Chronic Fatigue Syndrome (CFS)/myalgic encephalomyelitis (ME) is a complex condition of variable severity, affecting the physical and mental wellbeing young people. Triggers are recognised such as infection, physical illness or emotional distress. There has also been a link between adverse childhood events and likelihood of CFS/ME in later life. This cohort study looked at whether those with CFS have a higher incidence of parental separation, lack of contact with a parent and whether this sub-group have more severe disease at diagnosis.

Methods Retrospective analysis of the notes of 123 patients who attended a tertiary level adolescent CFS/ME service between April 2012 and April 2014 was conducted with further sub-group breakdown of those with parental separation and lack of contact with a parent with analysis on whether this group had an increase disease severity of CFS/ME at diagnosis.

Results Out of a total of 123 patients, 38% (n=45) had parents who were separated, compared with 10.6% of families nationally defined as a one parent household. Furthermore, 66% (n=32) with separated parents had lack of contact with at least one parent. Most patients in this cohort had moderate CFS at diagnosis (defined by NICE as having reduced mobility and restriction in all activities of daily living). Further sub-group analysis of those with separated parents showed no difference in severity compared to the non-separated group (p=0.69). There was no difference in disease severity for the group who lacked contact with at least one parent (p=0.51).

Conclusion Parental separation or lack of contact is not related to severity of CFS at diagnosis in this cohort. Larger cohort analysis of the demographics of those with CFS/ME is needed to attempt prediction of those who are more at risk of having severe CFS/ME, the impact of family history of CFS/ME and the impact on school attendance and bullying on severity. Further analysis of family structure and whether this impacts on response to treatment is also needed as this may help influence the structure and support of treatment.

G532(P) 'SCFN' (SUBCUTANEOUS FAT NECROSIS) OF THE NEWBORN – AN IMPORTANT DIFFERENTIAL DIAGNOSIS IN EVALUATION FOR CAUSES OF BRUISE-LIKE SKIN LESIONS IN A NEONATE

UA Khan, E Clifford, M Modair, R Barry. Department of Paediatrics, Mercy University Hospital, Cork, Ireland

10.1136/archdischild-2020-rcpch.450

Introduction SCFN of the newborn is an uncommon, self-limiting panniculitis characterized by development of erythematous, firm, indurated plaques or nodules commonly over the extremities, back, buttocks and thighs, mostly occurring in

term or post term babies within the first few weeks after birth. Hypercalcaemia is a life threatening complication which requires monitoring for several months. Perinatal mechanical stress, tissue hypoxia, and hypothermia may contribute to the pathogenesis. Exact incidence is unknown with a nearly equal male to female ratio.

Case Background We report a neonate presenting to emergency department by parents at 10 days of life from home with a sudden onset bruise-like linear lesion 6 cm long and 1.5 cm wide extending from the midline of upper part of the back to the right axilla. There was no history of trauma and baby was born via NVD. The skin lesion was red/purple in colour, non tender and palpable on examination. Clinical examination was otherwise unremarkable. As the marks on the back were suggestive of bruising, the baby underwent detailed Non Accidental Injury (NAI) investigations and social services assessment. Coagulation studies and FBC were normal. MRI brain reported subdural haematoma and suspected skull fracture, making NAI a strong consideration. Ophthalmology assessment and skeletal survey were normal. CT Skull was advised and this excluded skull fracture in favour of vermian skull bones. The origin of the subdural haematoma was then deemed uncertain but reported to be likely associated with vaginal delivery. Dermatology consultation confirmed the clinical diagnosis of SCFN of newborn.

Conclusion SCFN is a rare but important condition to consider in young infants who present with unexplained bruise-like lesions which require thorough investigation and consideration of possible NAI. Dermatology opinion was very helpful and should be sought if there is uncertainty about differential diagnosis.

G533(P) TIME FOR A KITKAT? AN ANALYSIS OF NIGHT SHIFT BREAK HABITS IN JUNIOR DOCTORS

¹SD Ahsan, ¹N Khanderia, ¹B Mirza, ²A Roueche, ²M Butler. ¹GKT School of Medical Education, King's College London, London, UK; ²General Paediatrics, Evelina London Children's Hospital, London, UK

10.1136/archdischild-2020-rcpch.451

Background/Introduction In the climate of increased patient safety awareness, the BMJ launched the 2019 'give us a break' campaign urging doctors to take breaks to combat fatigue. A cross-sectional survey of junior doctors found poor break uptake with further studies showing that fatigue is associated with medical errors.^{1 2}

Aim(s)/Objectives

- Establish whether night-shift doctors are taking breaks.
- Improve the percentage of doctors taking night-shift

Methods Four PDSA cycles were developed including:

- Implementing break rota charts
- Raising awareness at the Junior Doctor Forum
- Implementing a Whatsapp group for the night team
- Raising awareness about the importance of breaks with posters

Results Baseline data showed break uptake in 100% of Paediatric Nurse Practitioners (PNPs) whilst 0% of the general team doctors and 50% of specialty doctors were taking breaks. Post night-shift surveys identified common reasons. PDSA1 showed break uptake in 50% of doctors from both

teams, and after interventions by PDSA3 all night team doctors were taking breaks.

Discussion/Conclusion PDSA interventions were successful in increasing staff breaks, however sustainability of results in Junior Doctors will require a culture shift where doctors take personal responsibility for taking breaks. This is well-modelled by the PNPs who effectively arrange cross cover for breaks. A culture shift to improve patient safety overall can be aided by nominating a Junior Doctor break 'Champion'. Encouraging break taking in staff on night shifts will not only increase morale and wellness in the workplace but also contribute to improved patient care.

REFERENCES

1. Jackson E and Moreton A. Safety during night shifts: a cross-sectional survey of junior doctors' preparation and practice: Table 1. *BMJ Open* 2013(9);3:e003567.
2. Murray A, Pounder R, Mather H and Black D. Junior doctors' shifts and sleep deprivation. *BMJ* 2005(7505);330:1404.
3. West C. Association of Resident Fatigue and Distress With Perceived Medical Errors. *JAMA* 2009(12);302:1294.

G534(P) TRUST PROVIDED BREAKFAST FOR ALL NIGHT STAFF IN THE CHILDREN'S HOSPITAL

S Mullen, M Kearney, B Bartholome. *PED, RBHSC, Belfast, UK*

10.1136/archdischild-2020-rcpch.452

Aims Staff wellness is a fundamental pillar of the modern service. In the current climate, staff are under increasing stress with the potential to feel undervalued. The overarching aim of this project was to try to give something back to the work force to recognise the incredible service they provide.

Methods Following a multi-disciplinary discussion, the option of a complimentary post night shift breakfast was trialled. We all acknowledging that post night shift staff can be tired, hungry, emotionally fatigued and stressed. This was available to all staff within RBHSC and not limited to medical or nursing staff. Promotion occurred in the form of posters, emails and word of mouth.

Results An emailed survey was distributed to all staff to gain feedback with 49 responders. 79% were aware of the project was running with 93% strongly agreeing or agreeing that this was a great policy. 92% felt the addition of a snack would be beneficial. When questioned if this helped staff get home safe, 50% agreed or strongly agreed, 38% neutral and 12% disagreeing. In the free text box comments included; 'I believe this would do some way of showing a caring work force they are cared for too', 'It's lovely for staff to bump into others post night and chat/de-stress', 'Kindness is magic' and 'I think it is a lovely gesture to show that management are trying to support all staff working overnight'. The project started with offering a hot beverage and following feedback was extended to include water, fruit juice, light snacks as well as hot beverages.

Conclusion The project was promoted on Twitter with excellent feedback and multiple re-tweets, generating a lot of positive dialogue. This project demonstrates how we can better look after our work force with small improvements aiming to improve morale, make staff feel more valued and prompting a positive culture in our work place. A key component was ensuring this was available to all staff who work nights, not just medical and nursing.

G535(P) DO TREATMENTS FOR PAEDIATRIC CHRONIC FATIGUE SYNDROME IMPROVE PAIN? A SYSTEMATIC REVIEW

¹C Ascough, ¹H King, ¹T Serafimova, ¹S Jackson, ¹L Beasant, ²JCW Brooks, ³AE Pickering, ¹E Crawley. ¹Centre for Academic Child Health, University of Bristol, Bristol, UK; ²School of Psychology, University of Bristol, Bristol, UK; ³School of Physiology, Pharmacology and Neuroscience, University of Bristol, Bristol, UK

10.1136/archdischild-2020-rcpch.453

Aims Paediatric chronic fatigue syndrome/myalgic encephalomyelitis (CFS/ME) is common (prevalence 1–2%). Two thirds of children experience moderate or severe pain, and this is associated with increased fatigue and poorer physical function. However, current treatment approaches do not target pain.

This systematic review aims to identify whether specialist treatment of paediatric CFS/ME improves pain.

Methods We conducted a detailed search in MEDLINE, EMBASE, PsycINFO, and the Cochrane Library. Two researchers independently screened texts published since 1994 with no language restrictions. We searched trial registration websites for unpublished trials and hand searched reference lists of all included studies. Inclusion criteria were (1) RCTs & observational studies; (2) Participants aged <19 years with CFS/ME; (3) Measure of pain (quantitative, qualitative, or mixed methods) before and after an intervention.

Results Of 1898 papers screened, 27 studies investigated treatments in paediatric CFS/ME, 20 of which did not measure pain. Only 5 treatment studies measured pain at baseline and follow-up and were included in this review. Four of the studies investigated behavioural interventions and one study investigated a pharmacological intervention (low dose clonidine). None of the interventions were specifically targeted at treating pain.

Pain measures used were heterogenous and included a Visual Analogue Scale, CHQ-87 Bodily Pain Subscale, Brief Pain Inventory and a mean Daily Observed Pain score. Of the included studies 2 showed no improvement in pain scores with treatment, 1 small study described an improvement in pain in one subgroup, and 2 studies identified improvements in pain measures in 'recovered' patients compared to 'non-recovered patients'.

Conclusion Despite the prevalence and impact of pain in children with CFS/ME surprisingly few treatment studies measured pain. In those studies that did measure pain, there is limited evidence that treatment helps improve pain scores. However, patients who recover appear to have less pain than those who do not recover. More studies are needed to determine if pain in paediatric CFS/ME requires a specific treatment approach, with particular focus on patients who do not recover following initial treatment.

G536(P) SEAMLESS YOUNG PEOPLE'S SERVICES AND RARE DISEASES: THE BEST OF BOTH WORLDS?

¹S Rasul, ^{1,2,3}R Tattersall, ¹M Dockery, ¹D Hawley, ¹S Maltby, ¹A McMahon. ¹Paediatric Rheumatology Department, Sheffield Children's Hospital, Sheffield, UK; ²Adolescent Rheumatology Department, Royal Hallamshire Hospital, Sheffield, UK; ³Adult Rheumatology Department, Royal Hallamshire Hospital, Sheffield, UK

10.1136/archdischild-2020-rcpch.454

Transition from paediatric to adult services is a challenging process and developing rare, life-threatening, lifelong illness