


## Lockdown during the COVID-19 pandemic: impact on infants with pyloric stenosis

The COVID-19 pandemic has posed challenges for the delivery of healthcare for infants with disruption to 6-week health checks and health visitor services.<sup>1</sup> An area of particular concern is late presentation to the hospital.<sup>2</sup> However, current data do not offer an objective picture of how significant a problem this may be, with other reports showing low rates of delays in presentation.<sup>3</sup> Infantile hypertrophic pyloric stenosis (IHPS) is a common, non-infective infantile condition with a predictable clinical course and therefore a good indicator condition to assess for delays in presentation. We aimed to assess whether infants with IHPS presented later during 'lockdown' compared with the same period the preceding year.

Ten centres within the UK (England, Scotland and Northern Ireland) contributed data from babies with IHPS via a website (covidinchildren.co.uk) between 23 March 2020 and 31 May 2020 (the COVID-19 lockdown period) and between 23 March and 31 May 2019 (controls). A total of 87 eligible infants were included, comprising 40 controls (46%) and 47 cases (54%). The demographic and baseline characteristics of the two groups were similar (table 1 and figure 1).

Median age and weight at presentation in the control group were 31 days (24–41) and 3600 g (3190–4081), and those in the lockdown group were 34 days (26–41) and 3580 g (3120–4085). These differences were not statistically significant ( $p=0.64$ ,  $p=0.84$ ) (figure 1). The change in standardised weight loss was also comparable. (table 2). Patients requirement for preoperative intensive



**Figure 1** Comparison between the age at presentation (A) and admission weight (B) of infants with IHPS in the control period (2019) and the lockdown period. No significant difference is seen between the two groups (age at admission  $p=0.64$ , admission weight  $p=0.84$ ). IHPS, Infantile hypertrophic pyloric stenosis.

care and serum biochemistry was also similar except the lockdown group had a statistically but not clinically significant higher serum potassium (4.16 vs 4.5 mmol/L,  $p=0.04$ ) (table 2).

As an indicator condition, we have some reassurance that infants with IHPS have not had a significantly delayed presentation due to the COVID-19 lockdown. A recent objective study looking at paediatric presentations to emergency departments found very low numbers of delayed presentations to the hospital, with minimal associated morbidity.<sup>3 4</sup> Prompt, proactive changes to National Health Service 111 algorithms, guidance for parents by the Royal College of Paediatrics and Child Health<sup>5</sup> and the rapid uptake of virtual general practice and health visitor consultations may have avoided morbidity. Further work, focusing on different types of conditions, or different subsections of society will help provide useful information relating to the impact of societal lockdown on healthcare-seeking behaviour

in the UK and will enable more effective delivery of healthcare provision and public messaging in the event of further lockdowns.

Paediatric Surgery Trainee Research Network,<sup>1</sup> Felicity Arthur ,<sup>2</sup> Rachel Harwood ,<sup>3</sup> Benjamin Allin,<sup>4</sup> George Stephen Bethell ,<sup>5</sup> Tristan Boam,<sup>6</sup> Sumita Chhabra,<sup>7</sup> Mary Patrice Eastwood,<sup>8</sup> Kathryn Ford,<sup>9</sup> Hannah Rhodes,<sup>10</sup> Sesi Hotonu ,<sup>11</sup> Aileen Rooney,<sup>12</sup> Jennifer Billington,<sup>13</sup> Joseph Rutherford Davidson ,<sup>14</sup> Greg Dewar,<sup>10</sup> Ciaran Durand,<sup>8</sup> Nick Engall,<sup>15</sup> Patrick Green,<sup>3</sup> Matthew Jobson,<sup>16</sup> Anthony Starr,<sup>17</sup> Lucinda GC Tullie,<sup>18</sup> PSTRN Collaborative Group

<sup>1</sup> Collaborative Trainee Research Group, Paediatric Surgery Trainees, UK, UK

<sup>2</sup> Paediatric Surgery, Royal Hospital for Children, Glasgow, UK

<sup>3</sup> Paediatric Surgery, Alder Hey Children's NHS Foundation Trust, Liverpool, UK

<sup>4</sup> Oxford Radcliffe Hospitals NHS Trust, Oxford, UK

<sup>5</sup> University Surgery Unit, University of Southampton, Southampton, UK

<sup>6</sup> Norfolk and Norwich University Hospital NHS Trust, Norwich, UK

<sup>7</sup> Paediatric Surgery, Alder Hey Children's Hospital, Liverpool, UK

<sup>8</sup> Paediatric Surgery, Royal Belfast Hospital for Sick Children, Belfast, UK

<sup>9</sup> Paediatric Surgery, John Radcliffe Hospital, Oxford, UK

<sup>10</sup> Department of Paediatric Surgery, Bristol Royal Hospital for Children, Bristol, UK

<sup>11</sup> Edinburgh Royal Infirmary, Edinburgh, UK

<sup>12</sup> Paediatric Surgery, Glasgow Children's Hospital Charity, Glasgow, UK

<sup>13</sup> St George's University Hospitals NHS Foundation Trust, London, UK

<sup>14</sup> Paediatric Surgery, Guy's and St Thomas' Hospitals NHS Trust, London, UK

<sup>15</sup> Paediatric Surgery, Manchester Children's Hospitals, Manchester, UK

<sup>16</sup> Paediatric Surgery, Chelsea and Westminster Healthcare NHS Trust, London, UK

<sup>17</sup> Paediatric Surgery, Morecambe Bay Health Authority, Manchester, UK

<sup>18</sup> Specialist Neonatal and Paediatric Surgery, Great Ormond Street Hospital NHS Foundation Trust, London, UK

**Correspondence to** Miss Felicity Arthur, Royal Hospital for Children, Glasgow G51 4TF, UK; felicity.arthur2@ggc.scot.nhs.uk

**Twitter** Rachel Harwood @RachelHarwood10 and Joseph Rutherford Davidson @PaedSurgJoc

**Collaborators** PSTRN Collaborative Group: writing group: BA, FA, GB and RH. PSTRN Committee: BA, FA, GB, TB, SC, MPE, KF, RH, HR, SH and AR.

**Contributors** Writing group: BA, FA, GSB and RH. PSTRN committee: BA, FA, GSB, TB, SC, MPE, KF, RH, HR, SH and AR. Data contributors: FA, GSB, JB, TB, JRD, CD, GD, MPE, NE, PG, SH, MJ, AS and LGCT.

**Funding** The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

**Competing interests** None declared.

**Patient consent for publication** Not required.

**Provenance and peer review** Not commissioned; internally peer reviewed.

This article is made freely available for use in accordance with BMJ's website terms and conditions for the duration of the covid-19 pandemic or until

**Table 1** Characteristics of control (2019 patients) and lockdown (2020) patients

	Control (n=40)	Cases (n=47)	P value
	n (%)*	n (%)*	
Sex			
Female	4 (10)	7 (15)	0.53
Male	36 (90)	40 (85)	
Comorbidities			
Major	2	4	0.79
Minor	6	6	
None	32	37	
	Median (IQR)	Median (IQR)	
Gestational age at birth (completed weeks)	39 (38–40)	39 (37–40)	0.49
Birth weight (g)	3600 (3190–4083)	3580 (3120–4085)	0.84

\*% of those with complete data.

**Table 2** Comparison of the primary and secondary outcome measures for infants presenting during the lockdown and control periods

	Control (n=40)	Lockdown (n=47)	P value
Preoperative PICU admissions	2 (5)*	2 (4.2)*	>0.99
	<b>Median (IQR)</b>	<b>Median (IQR)</b>	
Age at admission (days)	31 (24–41)	34 (26–41)	0.64
Admission weight (g)	3600 (3190–4083)	3580 (3120–4085)	0.84
Preoperative admission to PICU, n (%)	2 (5)	2 (4.2)	0.99
Admission serum pH	7.47 (7.43–7.51)	7.47 (7.42–7.51)	0.89
Serum bicarbonate (mmol/L)	33 (27.35–38.27)	29.1 (25.3–34.0)	0.15
Serum chloride (mmol/L)	97 (90.5–101.0)	100 (94–108)	0.08
Serum potassium (mmol/L)	4.16 (3.7–4.7)	4.5 (4.1–5.0)	0.04
Standardised weight change—birth to admission, z score	−1.03 (−1.61 to −0.64)	−1.21 (−1.44 to −0.92)	0.55

\*% of those with complete data.

PICU, paediatric intensive care unit.

otherwise determined by BMJ. You may use, download and print the article for any lawful, non-commercial purpose (including text and data mining) provided that all copyright notices and trade marks are retained.

© Author(s) (or their employer(s)) 2021. No commercial re-use. See rights and permissions. Published by BMJ.



**To cite** Research Network PST, Arthur F, Harwood R, *et al.* *Arch Dis Child* 2021;**106**:e32.

Accepted 15 November 2020  
Published Online First 1 December 2020

*Arch Dis Child* 2021;**106**:e32.  
doi:10.1136/archdischild-2020-320544

#### ORCID iDs

Felicity Arthur <http://orcid.org/0000-0002-7983-2592>

Rachel Harwood <http://orcid.org/0000-0003-3440-3142>

George Stephen Bethell <http://orcid.org/0000-0002-1302-0735>

Sesi Hotonu <http://orcid.org/0000-0003-4649-7318>

Joseph Rutherford Davidson <http://orcid.org/0000-0002-3303-6320>

#### REFERENCES

- 1 NHS England. COVID-19 prioritisation within community health services 2020.
- 2 Lynn RM, Avis JL, Lenton S, *et al.* Delayed access to care and late presentations in children during the COVID-19 pandemic: a snapshot survey of 4075

paediatricians in the UK and Ireland. *Arch Dis Child* 2021;**106**:e8.

- 3 Roland D, Harwood R, Bishop N, *et al.* Children's emergency presentations during the COVID-19 pandemic. *The Lancet Child & Adolescent Health* 2020;**4**:e32–3.
- 4 Roland D, Nijman R, Ponmani C, *et al.* Arriving late, delayed, or not at all—presentations to paediatric emergency departments during covid-19 pandemic. *The BMJ Opinion* 2020 <https://blogs.bmj.com/bmj/2020/08/15/arriving-late-delayed-or-not-at-all-presentations-to-paediatric-emergency-departments-during-covid-19-pandemic>
- 5 RCPCH. Advice for parents during coronavirus, 2020. Available: [https://www.rcpch.ac.uk/sites/default/files/2020-04/covid19\\_advice\\_for\\_parents\\_when\\_child\\_unwell\\_or\\_injured\\_poster.pdf](https://www.rcpch.ac.uk/sites/default/files/2020-04/covid19_advice_for_parents_when_child_unwell_or_injured_poster.pdf)