Quality Improvement and Patient Safety

1787

'SKILLS AND DRILLS': STAFF TRAINING SESSIONS IN THE PAEDIATRIC EMERGENCY DEPARTMENT OF A MAJOR TRAUMA CENTRE

Michael Burns, Alice Taylor, Andrew Tester, Lailah Peel, David Baker, Dominique Kampman, Steve Foster. *Royal Hospital for Children, Glasgow, UK*

10.1136/archdischild-2021-rcpch.848

Background The Scottish Trauma Network (STN) was set up to form four new Major Trauma Centres across Scotland.

To prepare our paediatric ED for this a series of 'Skills and Drills' sessions were created by ED staff to cover a range of topics related to the practical care of a patient in a major trauma setting.

Objectives The aim was to create safe, socially distanced teaching sessions with pre-reading materials that would enable staff to fulfil the Key Performance Indicators of a Major Trauma Centre in the midst of the global pandemic.

Methods An initial six week pilot round of sessions was delivered and feedback sought from staff who had attended. This was taken into account and used to update the sessions.

Staff chose from a selection of specific words to describe the sessions and were encouraged to add their own words that they perceived to be relevant. A 'word cloud' was created that visually highlighted the most commonly selected words.

Staff were asked to provide a number out of 10 that described their knowledge of the topic covered in the session both before and after the session was delivered. Averages were then visualised in bar charts to represent the change in knowledge levels following sessions.

Results A word cloud was created. The more times a word was chosen, the larger it was represented on the chart. Sessions were widely found to be practical, interactive and supportive.

Average values for self-estimated knowledge levels out of 10 were represented in bar charts and are shown:

Conclusions Common descriptors of the sessions included: 'practical', 'interactive', and 'informative'. This suggests that

sessions reduced cognitive load leaving more time for staff to assess patients in front of them. The bar charts show a subjective estimation of increased knowledge in every topic covered during the sessions

Given the positive responses prompted by these initial sessions, it is hoped that similar sessions may be delivered by other departments in the hospital. The success of these trial sessions is an example of the benefits of an interactive, structured, face to face teaching programme that is achievable within the current constraints of social distancing.

Association of Paediatric Emergency Medicine

1788

THE PEM SPECIALIST: APPLYING AND DEVELOPING THEIR UNIQUE SKILLSETS IN THE EARLY MANAGEMENT OF PIMS-TS

¹Susannah Pye, ²Elise Randle, ²Padmanabhan Ramnarayan. ¹*Children's Acute Transport Service;* ²*Great Ormond Street Hospital*

10.1136/archdischild-2021-rcpch.849

Background Between March 2020 and February 2021, 160 patients with acute covid infections and the associated paediatric inflammatory disorder, PIMS-TS, were transferred to paediatric intensive care units by the largest regional transport service in Europe.

Local paediatric emergency departments have become attuned to the presentation of these patients, implementing early monitoring, workup and treatment in line with evolving national guidance. We have identified there has been reduction in the time taken to identify PIMS-TS, to initiate treatments and referral to PICU transport teams, potentially improving patient outcomes.

The role and skillset of the PEM specialist is vital in providing early identification and management of this patient group. We make the case that point of care ultrasound

Abstract 1787 Table 1

Pre-Alert	Airway Setup/RSI setup	Respiratory Support	Ventilator	Major Haemorrhage Protocol	Patient Transfer onto bed	Patient Transfer from Resus	Haemorrhage Control	
Knowledge Level Before Teaching	5.9	5.7	5.6	7	3.5	4.5	4.5	4.7
Knowledge Level After Teaching	8.5	8.4	8.4	8.5	6.8	8.1	7.8	7.3

Abstract 1787 Table 2

Pre-Alert	Patient Transfer onto bed	Airway Setup and Respiratory Support	Ventilator use	Cardiac arrest and use of Defibrillator	Cardiac arrest and use of Defibrillator (repeat session)	Paediatric Emergency Drugs	Central venous access/IO access	
Knowledge Level Before Teaching	6.8	6.6	6.8	4.5	5.9	5.7	4.5	5
Knowledge Level After Teaching	9.2	9.2	8.8	8.5	8.7	7.9	7.7	7.4

scanning (POCUS) and applied PICU clinical skills make this a subset of specialists with potential to further improve PIMS-TS outcomes.

Objectives This retrospective review compares two cohorts of PIMS-TS patients, the first and second covid waves, who required transport to paediatric intensive care units (PICU). The objective is to:

1. Describe any differences in presenting phenotypes of the two cohorts
2. Review timeliness of investigations and treatment interventions
3. Analyse changes in patient outcomes
4. Highlight where further improvement could be made, specifically around hypotension assessment and management in paediatric ED

Methods A retrospective review was performed for all patients with suspected PIMS-TS and COVID-19 infection transferred to PICUs by the regional transport team between 1st March 2020 and 28th February 2021. A cross-reference of local hospital, CATS and PICU notes were used to obtain necessary clinical data. This service evaluation project was registered with the trust and local GDPR guidelines were followed.

Results Data analysis is ongoing. The following areas will be reported on:

Signs, symptom & clinical assessment	-Patient demographics
Interventions	-Presentation phenotype
	-Management of hypotension
	-Medications
Patient outcomes (PICU)	-Length of PICU stay and total hospital LOS
	-Duration of vasoactive agent support
Incidental findings	-Arterial line placements at local hospital
	-Choice of vasoactive agents + familiarity at local hospitals

Conclusions Data analysis is ongoing. Early findings suggest local hospitals are more familiar with the presenting features of PIMS-TS and the spectrum of conditions it has phenotypical overlap with. In the first wave there were delays in diagnosis, in the second wave there is a suggestion of potential over diagnosis - attention and caution around this is vital.

Teams were vigilant around hypotension, with early use of vasoactive agents rather than large fluid resuscitation volumes. The number of ECHOs (particularly POCUS assessments) performed increased guiding decisions around transfer and vasoactive choice.

Frontline PEM clinicians have an important skillset in early detection and management of PIMS-TS patients, especially those needing cardiovascular support. Applying POCUS training for clinical assessment of cardiac function and inserting peripheral arterial lines under local anaesthetic to obtain accurate BP recordings can guide prompt and appropriate treatment for patients with PIMS-TS. This has wider-reaching implications on PEM training and their evolving role within the paediatric service.

Paediatric Mental Health Association

1789

CHANGES TO THE COMMUNITY CHILD HEALTH SERVICES FOR CHILDREN AND ADOLESCENTS WITH ADHD IN THE UK DURING THE COVID-19 PANDEMIC

Michael Ogundele. *Bridgewater Community Healthcare NHS Foundation Trust*

10.1136/archdischild-2021-rcpch.850

Background ADHD is the most commonly diagnosed neuro-behavioural disorders among children and young people (CYP). Given that patients with ADHD are already experiencing significant functional impairment in several domains, they are particularly vulnerable to the distress caused by the COVID-19 pandemic and social distancing measures. It is therefore important to understand changes and helpful adaptations to the ADHD service provision during the pandemic in order to learn lesson that could be applied during preparations for any future pandemic or further lockdowns.

Objectives We aimed to ascertain how clinicians have adapted their practices in response to the pandemic and their awareness of Covid-19-related professional guidelines in the management of CYP with ADHD.

Methods An online custom-designed questionnaire was approved by the Executive committee members of the National ADHD Network, George-Still Forum (GSF) and emailed to all the delegates who have attended the latest GSF annual scientific meeting.

Results Sixty-two of the 100 eligible respondents completed the survey (53 online and 9 as email attachments). Two thirds of the respondents were seeing 76% to 100% of their follow-up (FU) patients remotely, of which 98% involved telephone calls. Almost half of the clinicians (47%) were not offering clinic appointments for new referrals.

About a quarter of the respondents (24%) were able to obtain patients' Blood Pressure and Pulse, and growth measurements in accordance with the NICE recommendations. Locations for obtaining the physical measurements for patients with ADHD included specialist health centres (25%), GP surgery or other primary care locations (10%) and homes (13%).

Forty-two percent of respondents worked from home, 43% used video calls and 75% were able to signpost patients with ADHD and their families to various online resources for self-help and non-pharmacological behaviour management strategies.

Thirty-eight percent of the respondents reported being redeployed to other high-demand clinical areas during the pandemic including the acute general paediatrics (19%), statutory safeguarding work (6%) and adult medical care (2%).

In relation to awareness of professional guidelines, 32% of the respondents were aware of the general Covid-19 recommendations from the RCPCH and 33% knew about the European ADHD Guidance Group (EAGG) Covid-19 recommendations for the management of CYP with ADHD. However, 8% of the clinicians indicated no awareness of any Covid-19 specific guidance.

Conclusions Clinicians made several modifications in order to maintain ADHD services during the pandemic. The