The prevalence of the classical form was 58.3% in group 1 and 66.7% group 2 ($p=0.736$); atypical CD 41.7% vs 31.1% ($p=0.509$). Moderate and severe epigastric pain was common in both groups: 50.0% vs 31.1% ($p=0.309$). Significantly elevated level of anti- tTG antibodies ($>100\text{RU/ml}$) was detected equally in groups: 33.3% vs 47.7% ($p=0.516$). The prevalence of elevated level of antiparietal antibodies in groups was the same: in group 1- anti- H+/K+ ATPase antibodies in 2 children and in 3 children in group 2; and anti-intrinsic antibodies in 1 person in group 1 as in group 2. Thus, 25% vs 8.7% ($p=0.078$).

Conclusion There are conflicting studies results regarding the association between *H. pylori* and CD. Whether *Helicobacter pylori* triggers, doesn't affect or protects against CD is currently the subject of research. This study didn't reveal any effects of *H. pylori* on course of celiac disease in children despite the fact, that *H. pylori* is suspected as possible trigger of autoimmunity.

264 SCREENING FOR ANTIBODIES, ASSOCIATED WITH AUTOIMMUNE LIVER DISEASES IN CHILDREN WITH CELIAC DISEASE

Ekaterina Orlova*, Novikova Valeria, Shapovalova Natalia, Gurina Olga, Dementieva Elena, Kikunova Ksenia. *Institution of the presenting author*

10.1136/archdischild-2021-europaediatrics.264

Objective to determine the prevalence of autoimmune liver diseases in children with CD.

Methods We observed 45 children aged 3 to 16 years. CD in all patients was diagnosed according to the ESPGHAN criteria. Serological examination with the determination of antibodies to tissue transglutaminase (anti-tTG IgG, IgA); histological examination of the duodenal mucosa; genetic typing for HLA-DQ2/DQ8 were carried out. Duodenal histology having Marsh grade III features were eligible for the study. Antibodies to hepatic antigen cell nuclei, skeletal muscle, cell nuclei, mitochondria, smooth muscles of IgG class were determined by indirect immunofluorescence (nRIF) using reagent kits. The antigen were biochips of primate's liver, primate's musculus iliopsoas, human Hep2 epithelial cells, liver, kidney, and stomach of rats.

Normal titer $<1:80$. Anti-Parietal Cell Antibody (PCA), IgG was determined by nRIF using biochips primate's stomach as antigen. This kit of reagents detects antibodies for the diagnosis of such diseases as autoimmune hepatitis (AIH) 1, 2 and 3 types, primary biliary cirrhosis, primary sclerosing cholangitis (PSC), overlap syndrome (combination of AIH and PSC), autoimmune gastritis.

Results We didn't obtain elevated levels of antibodies associated with autoimmune liver disease in all children with CD. In 1 person we observed elevated levels of anti-parietal cell antibodies. It was a 15-year-old girl with a typical form of celiac disease, additionally suffering from primary oligomenorrhea, autoimmune diabetes mellitus (type 1). Further examination revealed non- *Helicobacter pylori* atrophic gastritis. Thus, autoimmune gastritis was diagnosed.

Conclusion Antibodies, associated with autoimmune liver diseases were uncommon in children with CD. Probably, due to insufficient amount of participants of the study. On the other hand, anti-parietal cell antibodies have been found.

265 SCFA (SHORT CHAIN FATTY ACIDS) PROFILE IN CHILDREN WITH COELIAC DISEASE

Tilen Seničar*, Andraž Kukovičič, Jernej Dolinšek, Maja Rupnik. *Faculty of Medicine University of Maribor*

10.1136/archdischild-2021-europaediatrics.265

The intestinal microbiota ferments complex carbohydrates that have not been broken down or absorbed during digestion into short-chain fatty acids (SCFA). SCFA are involved in different metabolic processes and the functioning of the immune system. Recent studies have shown changes in the SCFA profile in people with celiac disease (CD). Here we compared the composition of SCFA in the faecal water of the paediatric population with CD and the healthy children (HC).

Faecal samples were obtained from 5 individuals with CD (2 female, 3 male) on a gluten-free diet (GFD) and 5 HC (3 female, 2 male) aged 13-18 years. To determine the SCFA concentrations sterile filtered fecal water was prepared.

The SCFA profile was determined by gas chromatography.

We compared the average SCFA concentrations between groups with CD and HC, for which we used the Student's t-test. We also calculated the total SCFA and the fermentation index $FI = (\text{acetic acid} - (\text{propionic acid} + \text{butyric acid})) / (\text{the sum of SCFA})$.

The comparison of concentrations of individual SCFA fractions showed a statistically significantly lower value of acetic acid ($p = 0.04$) and a statistically marginal increase in caproic acid concentration ($p = 0.089$) in the CD group. The averages of other SCFA were higher in healthy individuals but without statistical significance. Total SCFA was statistically significantly higher ($p = 0.047$) in HC. Fermentation index was 0.092 in HC and 0.079 in CD patients.

The literature regarding the profile of SCFA in the paediatric population CD is scarce. Our results differ from the current literature, which reports a significant increase of acetic acid, total SCFA, and fermentation index in the paediatric population with CD compared to the control group. An increase in caproic acid in CD has not been reported yet. Recent studies have suggested that the study groups' age difference or the amount of gluten in the diet could explain the discrepancies in our results.

266 EFFECT OF DIETARY PROTEIN INTAKE ON BODY MASS INDEX AMONG PRIMARY SCHOOL CHILDREN

Ana Ilić*, Lucija Marić, Martina Bituh, Ivana Rumbak, Tea Karlović, Ružica Brečić, Irena Colić Barić. *Faculty of Food Technology and Biotechnology, University of Zagreb*

10.1136/archdischild-2021-europaediatrics.266

Recent studies have suggested that excessive dietary protein intake, especially animal-source protein, in infancy could affect health outcomes (e.g., obesity) in childhood. However, the effect of dietary protein intake in school-age children on growth and development is still unclear. The aim of this study was to assess dietary protein intake and its association with body mass index (BMI) in primary school children in Croatia.

Anthropometric measurements of children ($n=156$; 50% boys) aged 8.3 ± 0.5 years from primary schools in Zagreb City were performed according to standard protocols. Sex- and age-standardized BMI z-scores were obtained using AnthroPlus software. A dietary record for 3 non-consecutive