Results We performed 606 ileocolonoscopies in the above mentioned period due to suspicion of chronic inflammatory bowel disease or a bleeding polyp, and *E. vermicularis* was identified macroscopically in 10 patients (1.65%), of which 7/10 boys (70%). At the time of diagnosis, the median age was 10.99 years (3–17.5), and the median BMI was 18.62 kg/m² (13–22.06). Five patients (50%) were referred for diagnostic processing due to chronic inflammatory bowel disease suspicion, and five patients (56%) due to haematochezia and suspected bleeding polyp. The patients presented with various complaints of which: abdominal pain 4/10 (40%); blood in stool 6/10 (60%); diarrhea 3/10 (30%); and significant weight loss and perianal abscess in one patient. All patients had normal inflammatory parameters (CRP <2.8) and normal hemoglobin values for age (Hb 116–142). The mean duration of symptoms was 6.3 months (0.25 – 18). Parasite analysis of stool was performed in 4 patients (40%) and was negative in all, and perianal impression with adhesive cellophane tape was performed in only one patient, also negative. In all patients, *E. vermicularis* was confirmed macroscopically. In one patient, vulnerable mucosa reminding of ulcerative colitis (UC) was found along with enterobiasis, and in one patient we found non-specific terminal ileitis. Eosinophilia was found in four patients in intestinal biopsy samples. Following anthelmintic therapy with mebendazole, there was no recurrence of pinworm. One patient was later diagnosed with UC in the follow-up.

Conclusion *Enterobius vermicularis* is generally considered an innocent parasite, which in most cases causes perianal pruritus, and although it matures and lives in the intestines, severe gastrointestinal symptoms have rarely been reported. *E. vermicularis* infestation should be considered in patients with hematochezia, the screening of choice is perianal impression with cellophane tape, and the choice of treatment is mebendazole.

277

ASSESSMENT OF NUTRITIONAL STATUS OF CHILDREN WITH A NEWLY DIAGNOSED COELIAC DISEASE – A 10-YEAR TERTIARY CENTRE EXPERIENCE

Ivana Trivić*, Mario Mašić, Oleg Jadrešin, Iva Hojsak, Sanja Kolaček, Zrinjka Mišak. *Referral Centre for Paediatric Gastroenterology and Nutrition, Children's Hospital Zagreb, Zagreb, Croatia*

10.1136/archdischild-2021-europaediatrics.277

Introduction and Aims Coeliac disease is an immune-mediated systemic disease caused by gluten consumption that occurs in children and adults with a hereditary predisposition. The clinical features of the disease are diverse, and the most common symptoms in children are chronic diarrhoea, failure to thrive, vomiting, abdominal pain and chronic constipation. In some children, if cases when there is increased risk for coeliac disease despite the absence of symptoms and signs of the disease, targeted serological screening can be undertaken. The aim of the study was to assess and compare the nutritional status in symptomatic patients and children in whom the diagnosis of coeliac disease was made based on screening.

Methods This was a single-centre, retrospective, observational study involving all children diagnosed with coeliac disease at the Referral Centre for Paediatric Gastroenterology and Nutrition, at Children's Hospital Zagreb, in the period from 1 January 2010 to 1 January 2020. Nutritional status data were obtained from a systematic review of patients' electronic data.

Results During the study period, coeliac disease was diagnosed in 166 children (70 boys, mean age 7.47 years, age range 6 months to 17.5 years). In 28 children (16.37%) the disease was detected by screening. Data on body weight (BW), body height (BH) and body mass index (BMI) were recorded at the time of diagnosis in 137 children (82.5%) and based on the available data, Z-scores corrected for age and sex were determined for BW, BH and BMI for age and sex. The mean BW Z-score was -0.40 (-0.47 in symptomatic children vs -0.05 in children identified by screening, $p < 0.05$), the Z-score was <-2.00 in 17 children (12.41%) children. The mean BH Z-score was 0.17 (0.13 in symptomatic children vs 0.04 in children identified by screening, $p > 0.05$), a delay in linear growth (Z-score <-2.00) was noted in two children. The mean BMI Z-score was -0.60 (-0.67 in symptomatic vs -0.23 in children identified by screening, $p < 0.05$). The BMI Z-score was <-2.00 in 22 children (16.05%), of whom only two were children in whom coeliac disease was detected by screening. Only one child, with symptoms of the disease, was obese.

Conclusion At the time of diagnosis, BW and BMI of symptomatic children were significantly lower than in children with coeliac disease identified through screening, a total of 16.05% of children were malnourished. Normal linear growth was recorded in both groups. Most of the symptomatic children had normal nutritional status at the time of diagnosis, and the nutritional status in almost all children detected by screening was normal. In conclusion, normal nutritional status does not rule out the disease and should not deter a physician from conducting a targeted screening for coeliac disease.

278

PLANNING OF GLUTEN FREE DIET USING NUTRITIONAL SYSTEMS BIOLOGY APPROACH

¹Kristina Tušek*, ²Sandra Remenar, ²Ana Karatović, ²Ana Jurinjak Tušek, ²Maja Benković, ²Tamara Jurina, ²Davor Valinger, ²Jasenska Gajdoš Kljusurić. ¹Health Centre Krapina-Zagorje County, Mirka Crkvenca 1, 49000 Krapina, Croatia; ²University of Zagreb, Faculty of Food Technology and Biotechnology, Pierottijeva 6, 10000 Zagreb, Croatia

10.1136/archdischild-2021-europaediatrics.278

Introduction Celiac disease is an autoimmune disease that occurs in people with genetic predisposition, where gluten ingestion causes damage to the small intestines. When people with celiac disease eat gluten (a protein found in wheat, rye and barley), their body has an immune response that attacks the small intestine resulting in damage to the villi in small intestine. When the villi get damaged, nutrients cannot be properly absorbed into the body. So it is very important to know what happens in the cells after introducing gluten into organism. Lately, it has been recognised that systems biology tools have potential to increase understanding of how nutrition influences metabolic pathways and homeostasis. In this work the effect of the diet of paediatric patients on the celiac disease immune response was analysed using nutritional system biology approach.

Materials and Methods Celiac disease immune response mathematical model was constructed and analysed using CellDesigner 4.0 (Systems Biology Institute (SBI), Tokyo, Japan). Analysed mathematical model in the form of ordinary differential equations describes processes that take place in two intestinal compartments (lumen and lamina propria) and incorporates 16 variables and 34 processes, which correspond to 34 reaction rates. The effect of the different concentrations of gluten daily intake between paediatric patients on the