and parasitological processings were all negative. Bone marrow analysis revealed predominance of mature eosinophils without elements of malignant proliferation. Thus, diagnosis of chronic IEP was made. Oral administration of prednisone (0.7 mg/kg) provided a dramatic clinical improvement in two days. The prednisone dose was gradually reduced at intervals of two weeks and inhaled fluticasone 500 mcg daily was added. After 6 months systemic administration of prednisone was discontinued. However, 7 months later relapse of IEP occurred. Again, the boy responded very well to resumed oral prednisone (0.35 mg/kg) which was tapered again over a next 4 months course. Conclusion: The clinical presentation, the course of the disease, the severe eosinophilia of blood and BAL samples, and the absence of other known causes of pulmonary eosinophilia, all these diagnostic clinical elements suggested the diagnosis of chronic IEP. In spite to favourable long-term prognosis further follow-up of the boy is needed because according to modest published clinical experience relapses are always threatening.

428

THE USE OF NON-INVASIVE MECHANICAL VENTILATION IN THE PATIENT WITH DEFORMITY OF THE THORACIC SPINE – CASE REPORT

Iva Hižar*, Maja Bosanac, Ivan Pavić, Jadranka Sekelj Fureš. Department of Paediatrics, Children's Hospital Zagreb, Zagreb, Croatia

10.1136/archdischild-2021-europaediatrics.428

Introduction Non-invasive ventilation (NIV) refers to the delivery of mechanical ventilation, without the need for an invasive artificial airway (endotracheal tube or tracheostomy tube). There are different types of interfaces available for use with NIV including: total face mask, oronasal mask, nasal mask, and nasal pillows. The use of NIV has been shown to be an effective in patients with acute or chronic respiratory failure. Scoliosis is a complex three dimensional deformity that leads to impaired chest wall mechanics that prevent normal inflation of the lungs resulting in decreased total lung capacity. Rotation of the chest can produce displacement of the intrathoracic structures and compression of the main bronchi, causing upper airway obstruction. As the disease progresses, due to chronic airway inflammation and and secretion retention, small airways also became obstructed. Over time, ventilation-to-perfusion mismatch leads to chronic respiratory failure. In this case, we emphasize the importance of NIV in the patient with chronic respiratory failure caused by thoracic spine deformities.

Case Presentation A 17-year-old girl has been multidisciplinary followed at our hospital since early childhood due to progressive epileptic encephalopathy that led to severe cognitive impairment. The patient does not have major motor deficits and can walk or move unassisted. Due to recurrent respiratory infections associated with bronchial obstruction, at the age of 14 she underwent diagnostic testing at The Pulmonology, and Allergology division. Allergy tests were negative, and since the patient was uncooperative, pulmonary function tests were not performed. Chest radiograph showed significant progression of the idiopathic thoracolumbar scoliosis. Initially, the patient started therapy with inhaled corticosteroids, and later continued with combination therapy of inhaled steroids and long-acting beta2-agonists. Despite therapy, the patient developed severe acute respiratory failure during acute respiratory infections (most often viral infections with secondary bronchopneumonia). In February 2018, she developed severe acute

respiratory failure, from which she was slowly recovering and was discharged home after 20 days of hospital treatment, dependent on oxygen. In September 2018 the patient started non-invasive positive pressure ventilation via oronasal mask using Trilogy100 ventilator. Despite severe cognitive impairment, the patient accepted the mask very well, and she continued to use NIV at home only during sleep, an average 6-8 hours per day. With the use of NIV, the patient achieved satisfactory oxygen saturation throughout the day. In the following two years after NIV introduction, the patient did not have any worsening of the respiratory status and her quality of life was significantly improved.

Conclusion Deformities of the thoracic spine can lead to various pulmonary disorders. In patients with chronic respiratory failure, caused by deformity of the thoracic spine, non-invasive mechanical ventilation has proven to be an effective respiratory support.

429

SPONTANEOUS ELIMINATION OF AN INHALED FOREIGN BODY THROUGH THE CHEST WALL

¹Emilija Hostička*, ¹Nevenka Cigrovski, ¹Jadranka Kelečić, ²Miram Pasini. ¹University Hospital Centre Zagreb, Department of Pediatrics, Division for Respiratory Diseases, Clinical Immunology and Allergology; ²University Hospital Centre Zagreb, Department of Surgery, Division of Pediatric Surgery, Zagreb

10.1136/archdischild-2021-europaediatrics.429

A 20-month-old boy was referred to us from another hospital due to deterioration of the respiratory function during right-sided bronchopneumonia and a suspicion of foreign body aspiration. At the admission, the boy was subfebrile, tachycardic and tachypnoic, with normal oxygen saturation in the room air, drenched in sweat and with a productive cough. Inflammatory parameters were elevated and X-ray of the lungs showed a suspected inflammatory infiltrate in the right cardiophrenic angle. Bronchoscopy did not show the presence of a foreign body in the respiratory system. During hospitalization, swelling developed in the right infrascapular region with redness, induration and with palpable foreign body in the central part. A surgical incision in the general anesthesia was performed with foreign body extraction. It was a wall barley grass inflorescence (lat. *Hordeum murinum*). Due to the characteristic structure, wall barley inflorescence tends to move in only one direction. In the case shown, after aspiration, the barley grass spikelets migrated through the tracheobronchial tree, penetrating into pulmonary parenchyma to the periphery of the lung, through the pleural layers, intercostal muscles and finally protruded on the chest wall.

430

LUNG ABSCESS AS A DIFFERENTIAL DIAGNOSIS OF CHEST PAIN

Andrea Lidmila Obralić*, Nevenka Cigrovski, Ivan Bambir, Drago Čaleta, Jadranka Kelečić. University Hospital Center Zagreb, Department of Pediatrics, Department of Clinical Immunology, Respiratory and Allergological Diseases and Rheumatology

10.1136/archdischild-2021-europaediatrics.430

Introduction Chest pain is a common diagnosis in children in puberty and adolescents. Lung abscess is a rare cause of chest pain. It occurs as a result of a lung infection that has led to suppurative necrosis and destruction of the affected lung